

Your grade: 100%

Your latest: 100% • Your highest: 100% • To pass you need at least 80%. We keep your highest score.

Next item →

1. A function which maps ___ to ___ is a value function. [Select all that apply]

1 / 1 point

☒ States to expected returns.

✓ Correct

Correct! A function that takes a state and outputs an expected return is a value function.

☒ State-action pairs to expected returns.

✓ Correct


Correct! A function that takes a state-action pair and outputs an expected return is a value function.

☐ Values to actions.

☐ Values to states.

2. Consider the continuing Markov decision process shown below. The only decision to be made is in the top state, where two actions are available, left and right. The numbers show the rewards that are received deterministically after each action. There are exactly two deterministic policies, π_{left} and π_{right} . Indicate the optimal policies if $\gamma = 0$? If $\gamma = 0.9$? If $\gamma = 0.5$? [Select all that apply]

1 / 1 point



☒ For $\gamma = 0$, π_{left}

✓ Correct

Correct! Since both policies return to the top state every two time steps, to determine the optimal policy, it suffices to consider the reward accumulated over the first two time steps. For the policy left, this is equal to 1; for the policy right, this is equal to 0.

☒ For $\gamma = 0.5$, π_{right}

✓ Correct

Correct! Since both policies return to the start state every two time steps, to determine the optimal policy, it suffices to consider the reward accumulated over the first two time steps. For the policy left, this is equal to 1; for the policy right, this is equal to 1.

☒ For $\gamma = 0.9$, π_{right}

✓ Correct

Correct! Since both policies return to the top state every two time steps, to determine the optimal policy, it suffices to consider the reward accumulated over the first two time steps. For the policy left, this is equal to 1; for the policy right, this is equal to 1.8.

☐ For $\gamma = 0.9$, π_{left}

☐ For $\gamma = 0$, π_{right}

☒ For $\gamma = 0.5$, π_{left}

✓ Correct

Correct! Since both policies return to the start state every two time steps, to determine the optimal policy, it suffices to consider the reward accumulated over the first two time steps. For the policy left, this is equal to 1; for the policy right, this is equal to 1.

3. Every finite Markov decision process has ___. [Select all that apply]

1 / 1 point

☐ A stochastic optimal policy

- ☐ A unique optimal policy
- ☒ A unique optimal value function



Correct

Correct! The Bellman optimality equation is actually a system of equations, one for each state, so if there are N states, then there are N equations in N unknowns. If the dynamics of the environment are known, then in principle one can solve this system of equations for the optimal value function using any one of a variety of methods for solving systems of nonlinear equations. All optimal policies share the same optimal state-value function.

- ☒ A deterministic optimal policy



Correct

Correct! Let's say there is a policy π_1 which does well in some states, while policy π_2 does well in others. We could combine these policies into a third policy π_3 , which always chooses actions according to whichever of policy π_1 and π_2 has the highest value in the current state. π_3 will necessarily have a value greater than or equal to both π_1 and π_2 in every state! So we will never have a situation where doing well in one state requires sacrificing value in another. Because of this, there always exists some policy which is best in every state. This is of course only an informal argument, but there is in fact a rigorous proof showing that there must always exist at least one optimal deterministic policy.

4. The ____ of the reward for each state-action pair, the dynamics function p , and the policy π is ____ to characterize the value function v_π . (Remember that the value of a policy π at state s is $v_\pi(s) = \sum_a \pi(a|s) \sum_{s',r} p(s',r|s,a)[r + \gamma v_\pi(s')]$.)

1 / 1 point

- ☒ Mean; sufficient
- ☐ Distribution; necessary



Correct

Correct! If we have the expected reward for each state-action pair, we can compute the expected return under any policy.

5. The Bellman equation for a given a policy π : [Select all that apply]

1 / 1 point

- ☐ Holds only when the policy is greedy with respect to the value function.
- ☐ Expresses the improved policy in terms of the existing policy.
- ☒ Expresses state values $v(s)$ in terms of state values of successor states.



Correct

Correct!

6. An optimal policy:

1 / 1 point

- ☐ Is unique in every finite Markov decision process.
- ☒ Is not guaranteed to be unique, even in finite Markov decision processes.
- ☐ Is unique in every Markov decision process.



Correct

Correct! For example, imagine a Markov decision process with one state and two actions. If both actions receive the same reward, then any policy is an optimal policy.

7. The Bellman optimality equation for v_* : [Select all that apply]

1 / 1 point

- ☐ Expresses the improved policy in terms of the existing policy.
- ☒ Holds for the optimal state value function.



Correct

Correct!

- ☐ Holds for v_π , the value function of an arbitrary policy π .
- ☐ Holds when the policy is greedy with respect to the value function.
- ☒ Expresses state values $v_*(s)$ in terms of state values of successor states.



Correct

Correct!

8. Give an equation for v_π in terms of q_π and π .

1 / 1 point

- ☐ $v_\pi(s) = \max_a \pi(a|s) q_\pi(s, a)$

- ☐ $v_{\pi}(s) = \sum_a \gamma \pi(a|s) q_{\pi}(s, a)$
- ☐ $v_{\pi}(s) = \max_a \gamma \pi(a|s) q_{\pi}(s, a)$
- ☒ $v_{\pi}(s) = \sum_a \pi(a|s) q_{\pi}(s, a)$

✓ Correct
Correct!

9. Give an equation for q_{π} in terms of v_{π} and the four-argument p .

1 / 1 point

- ☐ $q_{\pi}(s, a) = \sum_{s'} \sum_r p(s', r|s, a) [r + v_{\pi}(s')]$
- ☒ $q_{\pi}(s, a) = \sum_{s'} \sum_r p(s', r|s, a) [r + \gamma v_{\pi}(s')]$
- ☐ $q_{\pi}(s, a) = \sum_{s'} \sum_r p(s', r|s, a) \gamma [r + v_{\pi}(s')]$
- ☐ $q_{\pi}(s, a) = \max_{s', r} p(s', r|s, a) [r + \gamma v_{\pi}(s')]$
- ☐ $q_{\pi}(s, a) = \max_{s', r} p(s', r|s, a) [r + v_{\pi}(s')]$
- ☐ $q_{\pi}(s, a) = \max_{s', r} p(s', r|s, a) \gamma [r + v_{\pi}(s')]$

✓ Correct
Correct!

10. Let $r(s, a)$ be the expected reward for taking action a in state s , as defined in equation 3.5 of the textbook. Which of the following are valid ways to re-express the Bellman equations, using this expected reward function? [Select all that apply]

1 / 1 point

☒ $v_*(s) = \max_a [r(s, a) + \gamma \sum_{s'} p(s'|s, a) v_*(s')]$

✓ Correct
Correct!

☒ $q_*(s, a) = r(s, a) + \gamma \sum_{s'} p(s'|s, a) \max_{a'} q_*(s', a')$

✓ Correct
Correct!

☒ $q_{\pi}(s, a) = r(s, a) + \gamma \sum_{s'} \sum_{a'} p(s'|s, a) \pi(a'|s') q_{\pi}(s', a')$

✓ Correct
Correct!

☒ $v_{\pi}(s) = \sum_a \pi(a|s) [r(s, a) + \gamma \sum_{s'} p(s'|s, a) v_{\pi}(s')]$

✓ Correct
Correct!

11. Consider an episodic MDP with one state and two actions (left and right). The left action has stochastic reward 1 with probability p and 3 with probability $1 - p$. The right action has stochastic reward 0 with probability q and 10 with probability $1 - q$. What relationship between p and q makes the actions equally optimal?

1 / 1 point

- ☐ $7 + 3p = -10q$
- ☐ $13 + 2p = -10q$
- ☒ $7 + 2p = 10q$
- ☐ $7 + 3p = 10q$
- ☐ $13 + 2p = 10q$
- ☐ $13 + 3p = 10q$
- ☐ $13 + 3p = -10q$
- ☐ $7 + 2p = -10q$

✓ Correct
Correct!