

MCMC Effective Sample Size (ESS) Summary

ESS for temnospondyli_pyrate_1_G_ADE_ADE_mcmc.log
Iterations: 23140000 Sampling Rate: 10000

posterior	prior	PP_lik	BD_lik
210	305	374	43

ESS for temnospondyli_pyrate_10_G_ADE_ADE_mcmc.log
Iterations: 4910000 Sampling Rate: 10000

posterior	prior	PP_lik	BD_lik
368	225	365	127

ESS for temnospondyli_pyrate_2_G_ADE_ADE_mcmc.log
Iterations: 24820000 Sampling Rate: 10000

posterior	prior	PP_lik	BD_lik
682	512	1072	120

ESS for temnospondyli_pyrate_3_G_ADE_ADE_mcmc.log
Iterations: 23540000 Sampling Rate: 10000

posterior	prior	PP_lik	BD_lik
251	624	477	42

ESS for temnospondyli_pyrate_4_G_ADE_ADE_mcmc.log
Iterations: 8590000 Sampling Rate: 10000

posterior	prior	PP_lik	BD_lik
325	405	421	43

ESS for temnospondyli_pyrate_5_G_ADE_ADE_mcmc.log
Iterations: 5030000 Sampling Rate: 10000

posterior	prior	PP_lik	BD_lik
394	303	395	83

ESS for temnospondyli_pyrate_6_G_ADE_ADE_mcmc.log
Iterations: 4890000 Sampling Rate: 10000

posterior	prior	PP_lik	BD_lik
352	168	356	46

ESS for temnospondyli_pyrate_7_G_ADE_ADE_mcmc.log
Iterations: 6020000 Sampling Rate: 10000

posterior	prior	PP_lik	BD_lik
372	239	369	68

ESS for temnospondyli_pyrate_8_G_ADE_ADE_mcmc.log
Iterations: 7850000 Sampling Rate: 10000

posterior	prior	PP_lik	BD_lik
207	237	439	32

ESS for temnospondyli_pyrate_9_G_ADE_ADE_mcmc.log
Iterations: 5280000 Sampling Rate: 10000

posterior	prior	PP_lik	BD_lik
354	178	353	46