

Interactions

Last time

Assumptions & Diagnostics

Today

Interactions/moderation

What are interactions?

When we have two variables, A and B, in a regression model, we are testing whether these variables have **additive effects** on our outcome, Y. That is, the effect of A on Y is constant over all values of B.


- Example: Drinking coffee and hours of sleep have additive effects on alertness; no matter how many hours I slept the previous night, drinking one cup of coffee will make me .5 SD more awake than not drinking coffee.

What are interactions?


However, we may hypothesize that two variables have **joint effects**, or interact with each other. In this case, the effect of A on Y changes as a function of B.

- Example: Chronic stress has a negative impact on health but only for individuals who receive little or no social support; for individuals with high social support, chronic stress has no impact on health.
- This is also referred to as **moderation**.


Univariate regression



Multivariate regression



Multivariate regression with an interaction



Example

We have an outcome (Stress) that we are interested in predicting from trait Anxiety and levels of Social Support.

```
##          vars    n   mean      sd median trimmed   mad   min    max
## id          1 118 488.65 295.95 462.50  485.76 372.13 2.00 986.00
## Anxiety     2 118   7.61   2.49   7.75   7.67   2.26 0.70  14.64
## Stress      3 118   5.18   1.88   5.27   5.17   1.65 0.62  10.32
## Support     4 118   8.73   3.28   8.52   8.66   3.16 0.02  17.34
## group*      5 118   1.53   0.50   2.00   1.53   0.00 1.00   2.00
##          range  skew kurtosis    se
## id          984.00  0.10    -1.29 27.24
## Anxiety     13.94 -0.18     0.28  0.23
## Stress      9.71  0.08     0.22  0.17
## Support     17.32  0.18     0.19  0.30
## group*      1.00 -0.10    -2.01  0.05
```

```
i.model1 = lm(Stress ~ Anxiety + Support + Anxiety:Support,  
              data = stress.data)  
i.model2 = lm(Stress ~ Anxiety*Support, data = stress.data)
```

Both methods of specifying the interaction above will work in R. Using the `*` tells R to create both the (partial) main effects and interaction effect. Note, however that the following code *gives you the wrong results*:

```
imodel_bad = lm(Stress ~ Anxiety:Support,  
                data = stress.data) # this does not create ma  
#It is VERY WRONG
```



```
i.model1 = lm(Stress ~ Anxiety*Support, data = stress.data)
summary(i.model1)
```

```
##
## Call:
## lm(formula = Stress ~ Anxiety * Support, data = stress.data)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -3.8163 -1.0783  0.0373  0.9200  3.6109
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   -2.73966    1.12101   -2.444  0.01606 *
## Anxiety         0.61561    0.13010    4.732 6.44e-06 ***
## Support        0.66697    0.09547    6.986 2.02e-10 ***
## Anxiety:Support -0.04174    0.01309   -3.188  0.00185 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 1.462 on 114 degrees of freedom
## Multiple R-squared:  0.4084,    Adjusted R-squared:  0.3928
## F-statistic: 26.23 on 3 and 114 DF,  p-value: 5.645e-13
```

$$\hat{Y} = b_0 + b_1X + b_2Z + b_3XZ$$

You can interpret the interaction term in the same way you normally interpret a slope coefficient -- this is the effect of the interaction controlling for other variables in the model.

You can also interpret the intercept the same way as before (the expected value of Y when all predictors are 0).

But here, b_1 is the effect of X on Y when Z is equal to **0**.

$$\hat{Y} = b_0 + b_1X + b_2Z + b_3XZ$$

Lower-order terms change depending on the values of the higher-order terms. The value of b_1 and b_2 will change depending on the value of b_3 .

- These values represent "conditional effects" (because the value is conditional on the level of the other variable).
- In many cases, the value and significance test with these terms is either meaningless (if Z is never equal to 0) or unhelpful, as these values and significance change across the data.

$$\hat{Y} = b_0 + b_1X + b_2Z + b_3XZ$$

Higher-order terms are those terms that represent interactions. b_3 is a higher-order term.

- This value represents how much the slope of X changes for every 1-unit increase in Z AND how much the slope of Z changes for every 1-unit increase in X.
- Symmetric!

Conceptual interpretation

Higher-order interaction terms represent:

- the change in the slope of X as a function of Z
- the degree of curvature in the regression plane
- the linear effect of the product of independent variables

```
stress.data$AxS = stress.data$Anxiety*stress.data$Support  
head(stress.data[,c("Anxiety", "Support", "AxS")])
```

##		Anxiety	Support	AxS
##	1	10.18520	6.1602	62.74287
##	2	5.58873	8.9069	49.77826
##	3	6.58500	10.5433	69.42763

```
summary(lm(Stress ~ Anxiety + Support + AxS, data = stress.da
```

```
##
```

```
## Call:
```

```
## lm(formula = Stress ~ Anxiety + Support + AxS, data = stress.data)
```

```
##
```

```
## Residuals:
```

```
##      Min       1Q   Median       3Q      Max
## -3.8163 -1.0783  0.0373  0.9200  3.6109
```

```
##
```

```
## Coefficients:
```

```
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) -2.73966     1.12101  -2.444  0.01606 *
## Anxiety      0.61561     0.13010   4.732 6.44e-06 ***
## Support      0.66697     0.09547   6.986 2.02e-10 ***
## AxS          -0.04174     0.01309  -3.188  0.00185 **
```

```
## ---
```

```
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
##
```

```
## Residual standard error: 1.462 on 114 degrees of freedom
```

```
## Multiple R-squared:  0.4084,    Adjusted R-squared:  0.3928
```

```
## F-statistic: 26.23 on 3 and 114 DF,  p-value: 5.645e-13
```

```
summary(lm(Stress ~ Anxiety*Support, data = stress.data))
```

```
##
## Call:
## lm(formula = Stress ~ Anxiety * Support, data = stress.data)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -3.8163 -1.0783  0.0373  0.9200  3.6109
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   -2.73966    1.12101  -2.444  0.01606 *
## Anxiety         0.61561    0.13010   4.732 6.44e-06 ***
## Support        0.66697    0.09547   6.986 2.02e-10 ***
## Anxiety:Support -0.04174    0.01309  -3.188  0.00185 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 1.462 on 114 degrees of freedom
## Multiple R-squared:  0.4084,    Adjusted R-squared:  0.3928
## F-statistic: 26.23 on 3 and 114 DF,  p-value: 5.645e-13
```

They're the same!!

The regression line estimated in this model is quite difficult to interpret on its own.

- **Simple slope**: the equation for Y on X at different levels of Z; but also refers to only the coefficient for X in this equation
- **Conditional effect**: the slope coefficients in the full regression model. These are the lower-order terms associated with a variable. E.g., X has a conditional effect on Y.

Which variable is the "predictor" (X) and which is the "moderator" (Z)?

The conditional nature of these effects is easiest to see by "plugging in" different values for one of your variables. Return to the regression equation estimated in our stress data:

$$\hat{Stress} = -2.74 + 0.62(Anx) + 0.67(Sup) - 0.04(A \times S)$$

Set Support to 5

$$\begin{aligned}\hat{Stress} &= -2.74 + 0.62(Anx) + 0.67(5) - 0.04(Anx \times 5) \\ &= -2.74 + 0.62(Anx) + 3.35 - 0.2(Anx) \\ &= 0.61 + 0.42(Anx)\end{aligned}$$

The conditional nature of these effects is easiest to see by "plugging in" different values for one of your variables. Return to the regression equation estimated in our stress data:

$$\hat{Stress} = -2.74 + 0.62(Anx) + 0.67(Sup) - 0.04(A \times S)$$

Set Support to 10


$$\begin{aligned}\hat{Stress} &= -2.74 + 0.62(Anx) + 0.67(10) - 0.04(Anx \times 10) \\ &= -2.74 + 0.62(Anx) + 6.7 - 0.4(Anx) \\ &= 3.96 + 0.22(Anx)\end{aligned}$$

Interaction shapes

Often we graph the simple slopes as a way to understand the interaction. Interpreting the shape of an interaction can be done using the numbers alone, but it requires a lot of calculation and mental rotation. For those reasons, consider it a requirement that you graph interactions in order to interpret them.


Interaction shapes

Ordinal interactions




Interaction shapes


Cross-over (disordinal) interactions




```
library(sjPlot)
plot_model(imodel, type = "int")
```




```
plot_model(imodel, type = "int", mdrt.values = "meansd")
```



```
plot_model(imodel, type = "pred", terms = c("Support", "Anxie
```




```
plot_model(imodel, type = "pred", terms = c("Support", "Anxie
```



Pop Quiz

You are interested in the effects of "brain games" and sleep on the development of Alzheimer's disease. You hypothesize that games might aid in slowing the progression of AD if participants are well rested.

- Write out the regression equation. Which variable is the moderator?
- Interpret the coefficients.
- Sketch out a plot of the simple slopes. There are different ways of doing this reflecting different hypotheses, but make sure the hypothesis stated above is somehow shown.

The slope at any particular value is a combination of both b_1 and b_3

$$\hat{Y} = (b_0 + b_2Z) + (b_1 + b_3Z)X$$

$$se_{b@z} = \sqrt{se_{b_1}^2 + (2 * Z * cov_{b_1b_3}) + (Z^2 se_{b_3}^2)}$$

$cov_{b_1b_3}$ refers to the covariance of the coefficients, not the covariance of the variables. This may seem a strange concept, as we only ever have one value for b_1 and b_3 -- the covariance of these coefficients refer to idea that if we randomly sample from a population, estimate the coefficients each time, and then examine the covariance of coefficients across

```
round(vcov(imodel),4)
```

```
##              (Intercept) Anxiety Support Anxiety:Support
## (Intercept)          1.2567 -0.1330 -0.0966           0.0100
## Anxiety              -0.1330  0.0169  0.0110          -0.0015
## Support              -0.0966  0.0110  0.0091          -0.0011
## Anxiety:Support       0.0100 -0.0015 -0.0011           0.0002
```

We can use the standard error of the slope to estimate whether or not it is significantly different from 0.

$$\hat{Y} = (b_0 + b_2 Z) + (b_1 + b_3 Z)X$$

$$t = \frac{(b_1 + b_3 Z)}{se_{b@z}}$$

$$df = N - p - 1$$

$$\hat{Stress} = -2.74 + 0.62(Anx) + 0.67(Sup) - 0.04(A \times S)$$

We want to know whether anxiety is a significant predictor of stress at different levels of support.

```
library(reghelper)
simple_slopes(imodel, levels = list(Support = c(4,6,8,10,12)))
```

##	Anxiety	Support	Test	Estimate	Std. Error	t value	df	Pr(> t)	Sig.
## 1	sstest	4		0.4486	0.0886	5.0617	114	1.610e-06	***
## 2	sstest	6		0.3652	0.0733	4.9791	114	2.289e-06	***
## 3	sstest	8		0.2817	0.0654	4.3095	114	3.488e-05	***
## 4	sstest	10		0.1982	0.0674	2.9424	114	0.003946	**
## 5	sstest	12		0.1147	0.0786	1.4600	114	0.147036	

If you don't list levels, then this function will test simple slopes at the mean and ± 1 SD around the mean.

Simple slopes - Significance tests

What if you want to compare slopes to each other? How would we test this?

The test of the interaction coefficient is equivalent to the test of the difference in slopes at levels of Z separated by 1 unit.

```
coef(summary(imodel))
```

##	Estimate	Std. Error	t value	Pr(> t)
## (Intercept)	-2.73966246	1.12100519	-2.443934	1.606052e-02
## Anxiety	0.61561220	0.13010161	4.731780	6.435373e-06
## Support	0.66696689	0.09547464	6.985802	2.017698e-10
## Anxiety:Support	-0.04174076	0.01309328	-3.187954	1.849736e-03

Centering

The regression equation built using the raw data is not only difficult to interpret, but often the terms displayed are not relevant to the hypotheses we're interested.

- b_0 is the expected value when all predictors are 0, but this may never happen in real life
- b_1 is the slope of X when Z is equal to 0, but this may not ever happen either.

Centering

Centering your variables by subtracting the mean from all values can improve the interpretation of your results.

- Remember, a linear transformation does not change associations (correlations) between variables. In this case, it only changes the interpretation for some coefficients


```
stress.data = stress.data %>%  
  mutate(Anxiety.c = Anxiety - mean(Anxiety),  
         Support.c = Support - mean(Support))  
head(stress.data[,c("Anxiety", "Anxiety.c", "Support", "Suppo
```

```
##      Anxiety   Anxiety.c Support  Support.c  
## 1 10.18520   2.57086873   6.1602 -2.5697997  
## 2  5.58873  -2.02560127   8.9069  0.1769003  
## 3  6.58500  -1.02933127  10.5433  1.8133003  
## 4  8.95430   1.33996873  11.4605  2.7305003  
## 5  7.59910  -0.01523127   5.5516 -3.1783997  
## 6  8.15600   0.54166873   7.5117 -1.2182997
```

Why did we not center Y (Stress)?

```
summary(lm(Stress ~ Anxiety.c*Support.c, data = stress.data))
```

```
##
## Call:
## lm(formula = Stress ~ Anxiety.c * Support.c, data = stress.data)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -3.8163 -1.0783  0.0373  0.9200  3.6109
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    4.99580    0.14647   34.108 < 2e-16 ***
## Anxiety.c       0.25122    0.06489    3.872 0.000181 ***
## Support.c       0.34914    0.05238    6.666 9.82e-10 ***
## Anxiety.c:Support.c -0.04174    0.01309   -3.188 0.001850 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 1.462 on 114 degrees of freedom
## Multiple R-squared:  0.4084,    Adjusted R-squared:  0.3928
## F-statistic: 26.23 on 3 and 114 DF,  p-value: 5.645e-13
```

```
summary(imodel)
```

```
##
## Call:
## lm(formula = Stress ~ Anxiety * Support, data = stress.data)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -3.8163 -1.0783  0.0373  0.9200  3.6109
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   -2.73966    1.12101  -2.444  0.01606 *
## Anxiety         0.61561    0.13010   4.732 6.44e-06 ***
## Support        0.66697    0.09547   6.986 2.02e-10 ***
## Anxiety:Support -0.04174    0.01309  -3.188  0.00185 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 1.462 on 114 degrees of freedom
## Multiple R-squared:  0.4084,    Adjusted R-squared:  0.3928
## F-statistic: 26.23 on 3 and 114 DF,  p-value: 5.645e-13
```

What changed? What stayed the same?

Standardized regression equation

So far, we've only discussed the unstandardized regression equation. If you're interested in getting the standardized regression equation, you can follow the same procedure of standardizing your variables first and then entering them into your linear model.

This is OK

$$Y \sim z(X) + z(Z) + z(X)*z(Z)$$

$$Y \sim z(X)*z(Z)$$

This is not OK

$$Y \sim z(X) + z(Z) + z(X*Z)$$

```
stress.data = stress.data %>%  
  mutate(zStress = scale(Stress),  
         zAnxiety = scale(Anxiety),  
         zSupport = scale(Support))  
head(stress.data[,c("zStress", "zAnxiety", "zSupport")])
```

```
##      zStress      zAnxiety      zSupport  
## 1 -1.05630148  1.033649700 -0.78410538  
## 2  0.97450140 -0.814418147  0.05397639  
## 3  0.52978175 -0.413855421  0.55327992  
## 4  1.87547440  0.538751068  0.83313888  
## 5  0.04640497 -0.006123922 -0.96980334  
## 6 -0.02939564  0.217784639 -0.37173144
```

```
zmodel = lm(zStress ~ zAnxiety*zSupport, stress.data)
summary(zmodel)
```

```
##
## Call:
## lm(formula = zStress ~ zAnxiety * zSupport, data = stress.data)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -2.03400 -0.57471  0.01989  0.49037  1.92453
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   -0.09818    0.07807   -1.258  0.211093
## zAnxiety        0.33302    0.08602    3.872  0.000181 ***
## zSupport        0.60987    0.09149    6.666  9.82e-10 ***
## zAnxiety:zSupport -0.18134    0.05688   -3.188  0.001850 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.7792 on 114 degrees of freedom
## Multiple R-squared:  0.4084,    Adjusted R-squared:  0.3928
## F-statistic: 26.23 on 3 and 114 DF,  p-value: 5.645e-13
```

Which is correct?

```
zmodel = lm(zStress ~ zAnxiety + zSupport + zAxS, stress.data)
summary(zmodel)
```

```
##
## Call:
## lm(formula = zStress ~ zAnxiety + zSupport + zAxS, data = stress.data)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -2.03400 -0.57471  0.01989  0.49037  1.92453
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) -1.610e-15  7.173e-02   0.000  1.00000
## zAnxiety      8.161e-01  1.725e-01   4.732 6.44e-06 ***
## zSupport      1.165e+00  1.668e-01   6.986 2.02e-10 ***
## zAxS          -5.003e-01  1.569e-01  -3.188  0.00185 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.7792 on 114 degrees of freedom
## Multiple R-squared:  0.4084,    Adjusted R-squared:  0.3928
## F-statistic: 26.23 on 3 and 114 DF,  p-value: 5.645e-13
```

Which is correct?

```
summary(imodel)
```

```
##
## Call:
## lm(formula = Stress ~ Anxiety * Support, data = stress.data)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -3.8163 -1.0783  0.0373  0.9200  3.6109
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   -2.73966    1.12101  -2.444  0.01606 *
## Anxiety         0.61561    0.13010   4.732 6.44e-06 ***
## Support        0.66697    0.09547   6.986 2.02e-10 ***
## Anxiety:Support -0.04174    0.01309  -3.188  0.00185 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 1.462 on 114 degrees of freedom
## Multiple R-squared:  0.4084,    Adjusted R-squared:  0.3928
## F-statistic: 26.23 on 3 and 114 DF,  p-value: 5.645e-13
```

OG model


```
zmodel = lm(zStress ~ zAnxiety*zSupport, stress.data)
summary(zmodel)
```

```
##
## Call:
## lm(formula = zStress ~ zAnxiety * zSupport, data = stress.data)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -2.03400 -0.57471  0.01989  0.49037  1.92453
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   -0.09818    0.07807   -1.258  0.211093
## zAnxiety       0.33302    0.08602    3.872  0.000181 ***
## zSupport       0.60987    0.09149    6.666  9.82e-10 ***
## zAnxiety:zSupport -0.18134    0.05688   -3.188  0.001850 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.7792 on 114 degrees of freedom
## Multiple R-squared:  0.4084,    Adjusted R-squared:  0.3928
## F-statistic: 26.23 on 3 and 114 DF,  p-value: 5.645e-13
```

```
summary(lm(Stress ~ Anxiety.c*Support.c, data = stress.data))
```

```
##
## Call:
## lm(formula = Stress ~ Anxiety.c * Support.c, data = stress.data)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -3.8163 -1.0783  0.0373  0.9200  3.6109
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    4.99580    0.14647   34.108 < 2e-16 ***
## Anxiety.c       0.25122    0.06489    3.872 0.000181 ***
## Support.c       0.34914    0.05238    6.666 9.82e-10 ***
## Anxiety.c:Support.c -0.04174    0.01309   -3.188 0.001850 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 1.462 on 114 degrees of freedom
## Multiple R-squared:  0.4084,    Adjusted R-squared:  0.3928
## F-statistic: 26.23 on 3 and 114 DF,  p-value: 5.645e-13
```

Centered model. Changed? Same?

Pop Quiz

Go back to the first pop quiz. Assume you have now centered the appropriate variables, and now re-interpret your coefficients.

Next time...

Guest lecture by Josh Jackson on Bayesian Analyses!