

CONFORMAL SPHERICAL SPLINES FOR THE LAPLACE–BELTRAMI OPERATOR ON GENUS-ZERO SURFACES: CONSTRUCTION, ALGORITHM, AND EXPERIMENTS*

SHELVEAN KAPITA†

Abstract. Every smooth closed Riemannian surface of genus zero is isometric to the unit sphere carrying a conformal metric, the round metric multiplied by a positive factor. This paper solves the Poisson equation and the eigenvalue problem of the Laplace–Beltrami operator of such a metric with smooth spherical splines, taking the conformal factor as the only description of the geometry. Because the Dirichlet energy is conformally invariant in two dimensions, the stiffness matrix is the round-sphere matrix for every conformal metric, and the factor enters only the load vector, the mean-value constraint and the weighted mass matrix. A surface embedded in space with a conformal parametrization, a surface of revolution, and a surface given only as a triangle mesh are instances of the same problem that differ in how the factor is obtained; the first two are treated here, and the factor is checked against an area identity before any equation is discretized. The unknown is a spline on a fixed spherical triangulation in broken Bernstein–Bézier form, with C^r smoothness imposed algebraically through edge functionals and the conforming space realized by a null-space matrix computed once and reused for every metric. Each step of the computation is stated with the formulas it needs. Experiments on the round metric, a prolate spheroid, a dumbbell of revolution, an Evans–Fung red-blood-cell profile and a prescribed factor with no embedding show the rates $d+1$ in L^2 and d in energy for spline degree d , the rate $2d$ for eigenvalues with reproduction of the even-order eigenvalues of the round metric up to degree d to roundoff, and, against parametric surface finite elements on the same triangulations, smaller errors at comparable numbers of unknowns for degrees four and six.

Key words. Laplace–Beltrami operator, Poisson equation on surfaces, spherical splines, conformal parametrization, Bernstein–Bézier methods, genus-zero surfaces

MSC codes. 65N30, 65N25, 65D07, 41A15, 58J50

1. Introduction. Let λ be a smooth function on the unit sphere $\mathbb{S}^2$ and let $\bar{g} = e^{2\lambda}g_s$ be the conformal metric it defines, g_s being the round metric. We write M for the Riemannian surface $(\mathbb{S}^2, \bar{g})$, Δ_M for its Laplace–Beltrami operator and $dA_M = e^{2\lambda}dA$ for its area element. For a function f on M with zero mean, the Poisson equation

$$(1.1) \quad -\Delta_M u = f \quad \text{on } M, \quad \int_M u \, dA_M = 0,$$

and the eigenvalue problem $-\Delta_M u = \mu u$ are the two model problems for second-order elliptic computation on a surface, and they share their matrices. By the uniformization theorem [28] every smooth closed Riemannian surface of genus zero is isometric to such an M for some λ , so (1.1) covers the Laplace–Beltrami operator of every smooth closed genus-zero surface, and the isometry is what a conformal parametrization of an embedded surface provides. The question of this paper is practical: given λ , how does one turn either problem into a matrix problem for a smooth high-order spline on the sphere, and what has to be computed at each step? The answer has three parts, and the paper is organized around them.

First, the geometry. The whole of it is the conformal factor $w = e^{2\lambda}$, one positive function on the sphere. How w is obtained depends on the instance: for a surface

*Submitted to the editors DATE.

†Department of Mathematics, Texas A&M University, College Station, TX 77843, USA (kapita@tamu.edu).

in $\mathbb{R}^3$ with a conformal parametrization $\Phi: \mathbb{S}^2 \rightarrow M$, w is the area factor of Φ ; for a surface of revolution, w comes from a one-dimensional nonlinear equation, the Mercator matching, and is checked against a closed-form area; a prescribed λ needs no surface at all; and a surface given only as a triangle mesh requires the map to be computed, which is deferred to a separate paper. All four instances feed the same computation (Section 4).

Second, the approximation space. The unknown is a spherical spline of degree d and smoothness r on a fixed spherical triangulation [3, 34]. We write it in broken Bernstein–Bézier form [29]: every spherical triangle keeps its own $\binom{d+2}{2}$ coefficients, C^r smoothness across an edge is a set of linear functionals whose only geometric input is the spherical barycentric coordinates of the opposite vertex, the conforming space is the kernel of the stacked smoothness matrix, and a null-space matrix computed once turns the constrained problem into a symmetric positive definite system. Nothing in this part depends on λ (Section 3).

Third, the equation. Because the Dirichlet energy is conformally invariant in two dimensions, the weak form of (1.1) in the round metric reads

$$(1.2) \quad \int_{\mathbb{S}^2} \nabla u \cdot \nabla v \, dA = \int_{\mathbb{S}^2} w f v \, dA \quad \text{for all } v, \quad \int_{\mathbb{S}^2} w u \, dA = 0,$$

with the round gradient and area element. The stiffness matrix is therefore the round-sphere matrix for every conformal metric, assembled once, and λ enters the load vector and the mean constraint through the single factor w at the quadrature points (Section 5); the eigenvalue problem is the generalized eigenproblem of the same stiffness matrix and the weighted mass matrix. Section 6 lists the computation step by step with the formulas each step evaluates.

The contribution is the formulation in terms of the conformal metric and what it makes computable, rather than a new spline space: the conformal metric on the sphere as the object discretized, with embedded surfaces, surfaces of revolution, prescribed factors and meshed surfaces as instances that differ only in how w is supplied; the geometry carried by one scalar weight without approximation whenever λ is known exactly, so that the discretization itself introduces no geometric consistency error; the metric-independent stiffness matrix and the reusable smoothness and null-space matrices; the eigenvalue formulation with its parity reproduction for the round metric; and a computational comparison with parametric surface finite elements. For the round metric the linear system is that of Baramidze and Lai [10].

The experiments (Section 7) use five instances: the round metric, a prolate spheroid, a strongly necked dumbbell of revolution, the Evans–Fung red-blood-cell profile in physical units, and a prescribed factor with no embedding. On all of them, the spline solution of the Poisson equation converges at the rates $d+1$ in $L^2(M)$ and d in the $H^1(M)$ seminorm expected from the approximation theory of spherical splines [34, Ch. 14], and the eigenvalues converge at the rate $2d$ with reproduction of the even-order eigenvalues of the round metric to roundoff. The analysis behind the rates, and surfaces given only as triangle meshes, for which λ must itself be computed, are deferred to separate papers; here the emphasis is on what is computed and how.

Throughout, $\mathbb{S}^2$ is the unit sphere with round metric g_S , surface measure dA and tangential gradient ∇ ; $W^{k,q}$ are Sobolev spaces with norm $\|\cdot\|_{k,q,\Omega}$ and top-order seminorm $|\cdot|_{k,q,\Omega}$; eigenvalues are μ and λ is reserved for the logarithm of the conformal factor; references of the form Theorem 13.30 are to [34].

2. Related work. *Spherical splines.* The foundational papers are [2, 5, 3, 4], which introduced circular and spherical Bernstein–Bézier polynomials, established dimension and local bases for homogeneous spline spaces, and applied the construction to fitting scattered data on spheres and sphere-like surfaces. The approximation theory and the quasi-interpolation machinery are due to Neamtu and Schumaker [37], [34, Ch. 14]. Spherical splines have been used extensively for data fitting, by minimal energy and least squares [9, 13, 11] and with shape constraints on the Bernstein–Bézier coefficients [12]; such pointwise constraints transfer unchanged to genus-zero surfaces through the parametrization of this paper.

Spherical splines for PDEs, and the sphere-like setting of Baramidze and Lai. The closest prior work is that of Baramidze and Lai [10]: a second-order elliptic equation of Laplace–Beltrami type on the unit sphere is solved by the Galerkin method in $S_d^r(\Delta)$, with existence, uniqueness and convergence proved, the smoothness conditions enforced through the constrained least-squares and augmented Lagrangian machinery of [7], and numerical rates reported; [13] introduces nonhomogeneous spherical splines, restrictions of all trivariate polynomials of degree at most d , so that constants are reproduced for every degree. In the same constrained-coefficient framework, Lai and Lee [32] replace the Galerkin form by collocation, the strong form imposed at interior collocation points and the smoothness conditions imposed as constraints on the coefficients, with the overdetermined constrained system solved by least squares, convergence proved for domains of positive reach, and the approach extended to the three-dimensional Monge–Ampère equation [33]. That realization applies to the transplanted problem of this paper without change, since the strong form $-\Delta_{\mathbb{S}^2} \tilde{u} = w\tilde{f}$ involves only the spherical Laplacian of the Bernstein basis, algebraic in the coefficients by the formulas of Section 3, and the smoothness matrix is the same; we use the Galerkin form because it keeps the stiffness matrix symmetric positive semidefinite and free of the weight, which the eigenproblem needs. Surfaces other than the sphere enter that line of work through the sphere-like construction of [5, 3], a radial graph identified with the sphere by $x \mapsto x/|x|$, which requires M to be star-shaped and puts the surface into both the stiffness and the mass form through a general pulled-back metric. Under the conformal parametrization used here the Dirichlet form is invariant (Lemma 4.1), the stiffness matrix is the round-sphere matrix of [10], and the surface enters only through the scalar weight w ; the nonhomogeneous spaces of [13] would lift the parity restriction at the cost of Proposition 5.3.

Finite elements on surfaces. The dominant paradigm is the surface finite element method of Dziuk [23], surveyed in [24] and, for the parametric, trace and narrow-band variants with a full account of the geometric error, in Bonito, Demlow and Nochetto [17]: the surface is approximated by an interpolating polyhedron or an isoparametric lift, and a geometric consistency error enters alongside the Galerkin error. Higher-order parametric elements and their pointwise and a posteriori estimates are due to Demlow [21] and Bonito and Demlow [16], adaptivity on implicit surfaces to Demlow and Dziuk [22], and the a priori theory for the Laplace–Beltrami eigenvalue problem to Bonito, Demlow and Owen [18]. In all of these high order requires curved elements and lifted quadrature, and C^1 conformity on surfaces is delicate. Here, once a conformal parametrization is known, the geometry is represented through a smooth scalar weight with no further approximation, the discretization introduces no geometric consistency error, C^r smoothness comes from the edge functionals, and the stiffness matrix is independent of the surface. The price is the conformal factor itself, which for a surface given only as a triangle mesh must be computed from an approximately conformal map [26, 31], the subject of a companion paper.

Eigenvalue approximation and spectral geometry. The Galerkin approximation theory of symmetric eigenvalue problems is classical [8, 15]; Hersch's inequality [27] and its extension along the spectrum [30] give a posteriori bounds that every computed genus-zero spectrum must respect.

3. Spherical splines in broken Bernstein–Bézier form. This section sets out the spline machinery in the form in which it is used throughout the paper and in the code. Every spherical triangle keeps its own Bernstein–Bézier coefficients. Smoothness across an edge is a finite set of linear functionals of the coefficients on the two triangles that share the edge. The conforming spline space is the kernel of the matrix obtained by stacking these functionals, and no global basis is ever constructed. This is the Bernstein-constraint representation of [29], specialized to the sphere; the ingredients are those of [34, Ch. 13], and references of the form Theorem 13.30 are to that book. The section closes with the structural features of the spherical theory that shape everything that follows.

3.1. Spherical triangulations. A spherical triangle $T = \langle v_1, v_2, v_3 \rangle$ is the region of $\mathbb{S}^2$ bounded by the great-circle arcs joining $v_1, v_2, v_3 \in \mathbb{S}^2$, assumed to lie in an open hemisphere so that the vertices are linearly independent in $\mathbb{R}^3$. A spherical triangulation $\triangle$ is a finite set of spherical triangles covering $\mathbb{S}^2$, any two meeting in a vertex, an edge, or not at all, with N_V, N_E, N_T vertices, edges and triangles. Every edge belongs to exactly two triangles, and Euler's relation gives

$$(3.1) \quad N_E = 3N_V - 6, \quad N_T = 2N_V - 4$$

(Theorem 13.33). The mesh size $|\triangle|$ is the longest geodesic edge and $\theta_\triangle$ the smallest angle. All experiments use icosahedral refinements: level L has $N_T = 20 \cdot 4^L$ triangles, obtained by splitting every triangle of level $L-1$ at the geodesic midpoints of its edges.

3.2. Spherical barycentric coordinates and Bernstein polynomials. Let $V_T = [v_1 \ v_2 \ v_3] \in \mathbb{R}^{3 \times 3}$ be the matrix whose columns are the vertices of T . It is invertible because T lies in a hemisphere. The spherical barycentric coordinates of a point $x \in \mathbb{R}^3$ relative to T are

$$(3.2) \quad b(x) = (b_1, b_2, b_3) = V_T^{-1}x, \quad \text{so that} \quad x = b_1v_1 + b_2v_2 + b_3v_3.$$

They are linear, not affine, functions of x : nonnegative on T , with $b_1 + b_2 + b_3 > 1$ in the interior of T . The spherical Bernstein basis polynomials of degree d relative to T are

$$(3.3) \quad B_{ijk}^d(x) = \frac{d!}{i!j!k!} b_1(x)^i b_2(x)^j b_3(x)^k, \quad i + j + k = d,$$

restricted to $\mathbb{S}^2$. Each is the restriction of a homogeneous trivariate polynomial of degree d , and the $\binom{d+2}{2}$ functions (3.3) span the space $\mathcal{B}_d$ of spherical polynomials of degree d , which is exactly the restriction to $\mathbb{S}^2$ of all homogeneous polynomials of degree d in $\mathbb{R}^3$ (Section 13.1). Since $\mathcal{B}_d$ does not depend on T , every $p \in \mathcal{B}_d$ has a unique B-form relative to every spherical triangle,

$$(3.4) \quad p = \sum_{i+j+k=d} c_{ijk} B_{ijk}^d,$$

and we associate the B-coefficient c_{ijk} with the spherical domain point

$$(3.5) \quad \xi_{ijk} = \frac{iv_1 + jv_2 + kv_3}{|iv_1 + jv_2 + kv_3|}, \quad i + j + k = d.$$

We write $c_T \in \mathbb{R}^{\binom{d+2}{2}}$ for the vector of B-coefficients of p on T , ordered by domain points. Decomposing into spherical harmonics,

$$(3.6) \quad \mathcal{B}_d = \mathcal{Y}_d \oplus \mathcal{Y}_{d-2} \oplus \cdots \quad (\text{down to } \mathcal{Y}_0 \text{ or } \mathcal{Y}_1 \text{ by parity}),$$

where $\mathcal{Y}_\ell$ is the $(2\ell + 1)$ -dimensional space of spherical harmonics of exact degree ℓ (Theorem 13.22).

For a spherical triangulation Δ the broken spline space is

$$(3.7) \quad S_d^{-1}(\Delta) = \{s : \mathbb{S}^2 \rightarrow \mathbb{R} : s|_T \in \mathcal{B}_d \ \forall T \in \Delta\},$$

with no continuity across edges. A function in $S_d^{-1}(\Delta)$ is determined by the concatenation

$$(3.8) \quad c = (c_{T_1}, \dots, c_{T_{N_T}}) \in \mathbb{R}^N, \quad N = N_T \binom{d+2}{2},$$

of its B-coefficient vectors on the triangles of Δ . No coefficient is shared between triangles, and a domain point on a common edge carries one coefficient from each of the two triangles that contain it. The spline spaces of interest are

$$(3.9) \quad S_d^r(\Delta) = \{s \in C^r(\mathbb{S}^2) : s|_T \in \mathcal{B}_d \ \forall T \in \Delta\} \subset S_d^{-1}(\Delta), \quad r \geq 0.$$

Every planar Bernstein–Bézier algorithm whose input is a vector of barycentric coordinates applies to (3.3) verbatim, since the algorithms never use $b_1 + b_2 + b_3 = 1$: de Casteljau evaluation, directional differentiation, and subdivision [34, Ch. 13]. Since every global spherical polynomial restricts to each triangle as a member of $\mathcal{B}_d$ and is smooth,

$$(3.10) \quad \mathcal{B}_d \subset S_d^r(\Delta) \quad \text{for every } r \geq 0.$$

3.3. Derivatives of the homogeneous extension. Regard B_{ijk}^d as the homogeneous polynomial (3.3) on $\mathbb{R}^3$. For a direction $z \in \mathbb{R}^3$ let D_z be the directional derivative and $a = (a_1, a_2, a_3) = V_T^{-1}z$ the spherical barycentric coordinates of z . Then

$$(3.11) \quad D_z B_{ijk}^d = d(a_1 B_{i-1,j,k}^{d-1} + a_2 B_{i,j-1,k}^{d-1} + a_3 B_{i,j,k-1}^{d-1}),$$

with the convention that a Bernstein polynomial with a negative index is zero (Theorem 13.28). Taking $z = e_1, e_2, e_3$ gives the ambient gradient of the homogeneous extension H_p of $p \in \mathcal{B}_d$ in coefficient form, with $\alpha = (i, j, k)$ and e_m the m th unit multi-index,

$$(3.12) \quad \frac{\partial H_p}{\partial x_\ell} = d \sum_{|\alpha|=d} c_\alpha \sum_{m=1}^3 (V_T^{-1})_{m\ell} B_{\alpha-e_m}^{d-1}, \quad \ell = 1, 2, 3.$$

We write G_T for the resulting sparse matrix, the Bernstein gradient matrix of [29] with the rows of V_T^{-1} in the place of the planar barycentric gradients. Euler’s identity $x \cdot \nabla H_p(x) = d H_p(x)$ gives, for $|x| = 1$, the tangential gradient

$$(3.13) \quad \nabla_{\mathbb{S}^2} p(x) = (I - xx^\top) \nabla H_p(x) = \nabla H_p(x) - d p(x) x.$$

The first form is what the assembly evaluates at quadrature points; the second shows that the tangential gradient is algebraic in the coefficients, with ambient components in $\mathcal{B}_{d+1}$ (Theorem 13.15).

3.4. Restriction to an edge. Let $e = \langle v_1, v_2 \rangle$ be the edge of T opposite v_3 ; it is an arc of the great circle through v_1 and v_2 . On e the third coordinate vanishes, $b_3 = 0$, so by (3.3)

$$(3.14) \quad p|_e = \sum_{i+j=d} c_{ij0} B_{ij0}^d|_e, \quad B_{ij0}^d(x) = \binom{d}{i} b_1^i b_2^j \quad \text{for } x = b_1 v_1 + b_2 v_2 \in e.$$

The $B_{ij0}^d|_e$ are the linearly independent circular Bernstein polynomials on the arc [2], so two polynomials from the two triangles sharing e agree on e if and only if their coefficients at the $d+1$ shared domain points agree. In arc length the restriction is a trigonometric polynomial (Remark 13.4), which is why quadrature is unavoidable in assembly.

3.5. Smoothness functionals across an edge. Let $T = \langle v_1, v_2, v_3 \rangle$ and $\tilde{T} = \langle v_4, v_3, v_2 \rangle$ share the edge $e = \langle v_2, v_3 \rangle$, and let $s \in S_d^{-1}(\triangle)$ have B-coefficients c_{ijk} on T and $\tilde{c}_{ijk}$ on $\tilde{T}$, each indexed by the vertex order of its own triangle. Let $(\beta_1, \beta_2, \beta_3) = V_T^{-1} v_4$ be the spherical barycentric coordinates of the fourth vertex relative to T . By Theorem 13.30, s is C^r across e if and only if

$$(3.15) \quad \tilde{c}_{njk} = \sum_{\nu+\mu+\kappa=n} c_{\nu, k+\mu, j+\kappa} B_{\nu\mu\kappa}^n(v_4), \quad j+k = d-n, \quad n = 0, \dots, r,$$

where $B_{\nu\mu\kappa}^n(v_4) = \frac{n!}{\nu!\mu!\kappa!} \beta_1^\nu \beta_2^\mu \beta_3^\kappa$. The first two layers read

$$(3.16) \quad n = 0 : \quad \tilde{c}_{0jk} = c_{0kj}, \quad j+k = d,$$

$$(3.17) \quad n = 1 : \quad \tilde{c}_{1jk} = \beta_1 c_{1kj} + \beta_2 c_{0, k+1, j} + \beta_3 c_{0, k, j+1}, \quad j+k = d-1.$$

The layer $n = 0$ is the coefficient matching of Section 3.4. Each C^1 condition (3.17) ties one coefficient of $\tilde{T}$ in the first row parallel to e to three coefficients of T , the four circled points of Figure SM1 in Section SM1 of the supplement; the stencil is the planar one and only the numbers $\beta_1, \beta_2, \beta_3$ change. The layer n involves the coefficients in the rows $0, \dots, n$ parallel to e on both triangles, and the difference of the two sides of (3.15) is a linear functional of the broken coefficient vector c supported on the two triangles adjacent to e . We call these the smoothness functionals of order r across e and write

$$(3.18) \quad J_{0,e}^{(r)} \in \mathbb{R}^{m_e \times N}, \quad m_e = \sum_{n=0}^r (d+1-n),$$

for the matrix whose rows are these functionals, so that s is C^r across e if and only if $J_{0,e}^{(r)} c = 0$. A row of layer n has at most $\binom{n+2}{2} + 1$ nonzero entries. For $r = 0$ we write $C_{0,e} = J_{0,e}^{(0)}$. The subscript 0 marks the scalar slot of the constraint complex of [29]; only that slot is needed in this paper, since the Laplace–Beltrami problem is scalar and of second order.

The functionals (3.15) describe C^r continuity on the curved surface exactly: Theorem 13.30 is proved by extending $s|_T$ and $s|_{\tilde{T}}$ homogeneously to the tetrahedra with apex at the origin, which share the planar face $\langle 0, v_2, v_3 \rangle$, and applying the trivariate C^r conditions across that face. No element map is involved. In the terminology of [29, Sec. 7], where the transfer of C^r functionals to curved patches through element maps is left open, the sphere is the case in which the polynomials live on the surface itself, so the only geometric input is $V_T^{-1} v_4$.

Stacking the blocks $J_{0,e}^{(r)}$ over all N_E edges gives the smoothness matrix of the space,

$$(3.19) \quad J_0^{(r)} \in \mathbb{R}^{m \times N}, \quad m = N_E \sum_{n=0}^r (d+1-n), \quad S_d^r(\Delta) = \ker J_0^{(r)} \subset S_d^{-1}(\Delta).$$

The implementation stores the edge blocks, so the matrix has $O(N_E)$ rows and nonzeros for fixed d and r . Its rows are linearly dependent, since the conditions around a vertex are related, and this is allowed throughout; the number of dependencies is counted in Section SM1 of the supplement and confirmed on the meshes used.

3.6. Dimension from the rank. The dimension of the spline space is $N - \text{rank } J_0^{(r)}$ whether or not the rows are independent, with the rank delivered by the elimination that builds Z ; closed-form counts (Theorem 13.45) are recalled in Section SM1 of the supplement and checked against the computed ranks. For $d < 3r + 2$ the dimension can depend on the positions of the vertices [4], and the rank is then the only reliable count. The entire construction lives on the linear structure of $\mathbb{R}^3$: a general closed surface carries no ambient cone of polynomials, so any extension is a transfer from the sphere, by the radial projection of [5] for star-shaped surfaces or by the conformal parametrization of the next section for every smooth closed genus-zero surface.

4. The conformal metric and its instances.

4.1. The conformal metric on the sphere. Let $\lambda \in C^\infty(\mathbb{S}^2)$, $\bar{g} := e^{2\lambda} g_s$, and $M := (\mathbb{S}^2, \bar{g})$. We write $w := e^{2\lambda}$ for the conformal factor. The area element, gradient and Laplace–Beltrami operator of $\bar{g}$ are

$$(4.1) \quad dA_M = e^{2\lambda} dA, \quad \nabla_M u = e^{-2\lambda} \nabla u, \quad \Delta_M u = e^{-2\lambda} \Delta_{\mathbb{S}^2} u,$$

the last being the conformal transformation law in dimension two [28, Ch. 2]. A conformal diffeomorphism of $\mathbb{S}^2$ pulls $\bar{g}$ back to another conformal metric with the same geometry, so λ is determined by the geometry only up to the noncompact Möbius group [28], and an unnormalized representative can concentrate w arbitrarily. We fix this freedom by Hersch’s centering condition [27],

$$(4.2) \quad \int_{\mathbb{S}^2} x_i w dA = 0, \quad i = 1, 2, 3,$$

where x_1, x_2, x_3 are the ambient coordinate functions restricted to $\mathbb{S}^2$; a centered representative always exists [27, 36]. The remaining additive freedom in λ is a global scaling, which we normalize by

$$(4.3) \quad \text{Area}(M) = \int_{\mathbb{S}^2} w dA = 4\pi,$$

so that computed spectra are comparable with those of the round metric; the Poisson problem needs neither normalization, and the red blood cell is run in its own units without (4.3). We write $\Lambda_0 := \max_{\mathbb{S}^2} |\lambda|$ and $E := e^{\Lambda_0}$, so that $E^{-2} \leq w \leq E^2$; constants C are generic and depend at most on the arguments displayed.

Two forms carry the whole computation. On the round sphere define

$$(4.4) \quad a(u, v) := \int_{\mathbb{S}^2} \nabla u \cdot \nabla v dA, \quad b_w(u, v) := \int_{\mathbb{S}^2} w u v dA, \quad u, v \in W^{1,2}(\mathbb{S}^2),$$

and write $\|u\|_{L_w^2}^2 := b_w(u, u)$.

LEMMA 4.1 (Mass and energy of the conformal metric). *For all $u, v \in W^{1,2}(\mathbb{S}^2)$,*

$$(4.5) \quad \int_M u v dA_M = b_w(u, v),$$

$$(4.6) \quad \int_M \nabla_M u \cdot_{\bar{g}} \nabla_M v dA_M = a(u, v),$$

and $E^{-1} \|u\|_{L^2(\mathbb{S}^2)} \leq \|u\|_{L_w^2} \leq E \|u\|_{L^2(\mathbb{S}^2)}$. In particular $W^{1,2}(M) = W^{1,2}(\mathbb{S}^2)$ with equivalent norms.

Proof. (4.5) is the first identity of (4.1). For (4.6), $\bar{g}(\nabla_M u, \nabla_M v) = e^{-2\lambda} \nabla u \cdot \nabla v$ pointwise by the second identity; multiplying by $dA_M = e^{2\lambda} dA$ the factors cancel. This cancellation is the conformal invariance of the Dirichlet energy, special to dimension two. The norm comparison is $e^{-2\Lambda_0} \leq w \leq e^{2\Lambda_0}$. $\square$

The approximation power of the spherical spline spaces is Theorem 14.26: if Δ covers $\mathbb{S}^2$ with $|\text{star}(v)| < 1$ for every vertex and $|\Delta| \leq 1/6$, if $d \geq 3r + 2$, and if $0 \leq m \leq d$ with $m \equiv d \pmod{2}$, then for every $g \in W^{m+1,2}(\mathbb{S}^2)$ there is $s \in S_d^r(\Delta)$ with

$$(4.7) \quad |g - s|_{k,2,\mathbb{S}^2} \leq C(d, \theta_\Delta) |\Delta|^{m+1-k} |g|_{m+1,2,\mathbb{S}^2}, \quad 0 \leq k \leq m,$$

where θ_Δ is the smallest angle of Δ and $|\cdot|_{m+1,2,\mathbb{S}^2}$ is the seminorm of Section 14.5 of [34]. By Lemma 4.1 the same s approximates g in the norms of M : with $m = d$ and $k = 0, 1$,

$$(4.8) \quad \|g - s\|_{L^2(M)} + |\Delta| |g - s|_{1,2,M} \leq C(d, \theta_\Delta) E |\Delta|^{d+1} |g|_{d+1,2,\mathbb{S}^2},$$

since the energy seminorm of M equals that of the round sphere by (4.6) and the $L^2(M)$ norm is the weighted norm by (4.5). The spline spaces themselves need no change: $S_d^r(\Delta)$ is a space of functions on $\mathbb{S}^2$, hence on M , and C^r smoothness, the smoothness matrix $J_0^{(r)}$, the null-space matrix Z , the dimension $N - \text{rank } J_0^{(r)}$, determining sets and local bases are the same for every λ , because none of them involves the metric.

4.2. Instances: how the conformal factor is obtained. *An embedded surface with a conformal parametrization.* Let $\Sigma \subset \mathbb{R}^3$ be a smooth closed surface of genus zero with its induced metric, and let $\Phi: \mathbb{S}^2 \rightarrow \Sigma$ be a conformal diffeomorphism, which exists by uniformization [28]. Then $\Phi^* g_\Sigma = e^{2\lambda} g_s$ for one λ , so Φ is an isometry from M onto Σ , and every Laplace–Beltrami problem on Σ is the corresponding problem on M with data $f \circ \Phi$ and solution $u \circ \Phi^{-1}$. The factor is the area magnification of Φ : at $x \in \mathbb{S}^2$ with orthonormal tangent vectors τ_1, τ_2 ,

$$(4.9) \quad w(x) = |d\Phi_x(\tau_1) \times d\Phi_x(\tau_2)| = \sigma_1(x) \sigma_2(x),$$

where $\sigma_1 \geq \sigma_2 > 0$ are the singular values of $d\Phi_x$, and conformality is $\sigma_1 = \sigma_2$. In colatitude and longitude (Θ, ϕ) ,

$$(4.10) \quad w(\Theta, \phi) = \frac{|\partial_\Theta \Phi \times \partial_\phi \Phi|}{\sin \Theta},$$

which a code evaluates when Φ is given with its derivatives; with Φ alone, centered differences with step 10^{-5} reproduced the geometry diagnostics of the experiments below to about ten digits, and the ratio σ_1/σ_2 from the same differences measures the

deviation from conformality, assumed here negligible against the discretization error (an approximately conformal map perturbs the metric in both forms; its treatment is deferred). The values of w are needed only at quadrature points.

A surface of revolution. Let Σ be given by a generating curve $(r(\eta), z(\eta))$, $\eta \in [0, \pi]$, rotated about the z -axis, with $r > 0$ on $(0, \pi)$ and the curve meeting the axis regularly at both ends; the spheroid, the dumbbell and the red blood cell are of this form. The metric of Σ in (η, ϕ) is $(r'^2 + z'^2) d\eta^2 + r^2 d\phi^2$, and its Mercator variable is

$$(4.11) \quad t(\eta) = \ln \tan(\eta/2) + G(\eta), \quad G'(\eta) = \frac{\sqrt{r'^2 + z'^2}}{r} - \frac{1}{\sin \eta},$$

where G' extends smoothly to the poles. The round sphere has the same structure with $G \equiv 0$, and matching the two Mercator variables at the equator defines the conformal map $\Theta \mapsto \eta(\Theta)$ through the monotone scalar equation

$$(4.12) \quad \ln \tan(\eta/2) + \tilde{G}(\eta) = \ln \tan(\Theta/2), \quad \tilde{G} = G - G(\pi/2),$$

solved by a safeguarded Newton iteration, and $\Phi(\Theta, \phi) = (r(\eta) \cos \phi, r(\eta) \sin \phi, z(\eta))$. The factor is $w = r(\eta)^2 / \sin^2 \Theta$, which we evaluate in the numerically stable half-angle form

$$(4.13) \quad w = \left(\frac{r(\eta)}{\sin \eta} \right)^2 e^{-2\tilde{G}(\eta)} \left(\frac{1 + T^2}{1 + \tau^2} \right)^2, \quad \tau = \tan(\eta/2), \quad T = \tan(\Theta/2) = \tau e^{\tilde{G}(\eta)},$$

which is accurate at the poles. Star-shapedness is not used, so profiles with concavities are admissible. Equatorial symmetry makes the Hersch centering (4.2) automatic, and the exact area $2\pi \int_0^\pi r \sqrt{r'^2 + z'^2} d\eta$ is an independent identity for the computed factor, against which $\int_{\mathbb{S}^2} w dA$ under the assembly quadrature is compared as the first check of every run; in the experiments the agreement is at roundoff. For a radial graph $\{R(\psi) v(\psi, \phi)\}$ over the sphere the construction is the special case $r = R \sin \psi$, $z = R \cos \psi$, with $G' = (\sqrt{R^2 + R'^2} - R)/(R \sin \psi)$ and prefactor $R(\eta)^2$ in (4.13); the spheroid is the case with closed-form G .

A prescribed factor. Any smooth λ defines an instance; the computation needs only w at the quadrature points, and Section 7 includes one.

A surface given as a triangle mesh. The map Φ must then be computed, by harmonic energy minimization with Möbius normalization [26] or conformalized mean curvature flow [31], and is only approximately conformal; that instance is the subject of a companion paper.

5. The Poisson equation as a weighted problem on the round sphere.

The weak form of (1.1) is written in the round metric through the two identities of Lemma 4.1 and then in the broken coefficients of Section 3, with conformity imposed through the smoothness matrix. The section closes with the eigenproblem, which uses the same matrices.

5.1. The weak form in the round metric. Let $f \in L^2(M)$ with $\int_M f dA_M = 0$. The weak form of (1.1) is: find $u \in W^{1,2}(M)$ with $\int_M u dA_M = 0$ such that

$$(5.1) \quad \int_M \nabla_M u \cdot_{\bar{g}} \nabla_M v dA_M = \int_M f v dA_M \quad \text{for all } v \in W^{1,2}(M).$$

By (4.6) on the left and (4.5) on the right, and since $W^{1,2}(M) = W^{1,2}(\mathbb{S}^2)$, this is the problem on the round sphere: find $u \in W^{1,2}(\mathbb{S}^2)$ with $b_w(u, 1) = 0$ such that

$$(5.2) \quad a(u, v) = b_w(f, v) \quad \text{for all } v \in W^{1,2}(\mathbb{S}^2),$$

with the forms a and b_w of (4.4); the compatibility condition is $b_w(f, 1) = 0$. Three facts organize what follows. The metric enters only through w , on the right-hand side and in the constraint; a is the Dirichlet form of the round sphere. The problem is well posed: on $\{v : b_w(v, 1) = 0\}$ the form a is coercive, since $a(v, v) \geq \mu_1(M) b_w(v, v)$ there by Proposition 5.2 below with $\mu_1(M) > 0$, so Lax–Milgram gives a unique solution, which satisfies (5.2) for all v because both sides vanish at $v = 1$. The strong form is $-\Delta_{\mathbb{S}^2} u = wf$ by (4.1); for a manufactured u the product wf in the load vector is $-\Delta_{\mathbb{S}^2} u$ and contains no w , which the first experiment of Section 7 uses to separate the effect of the factor in the load vector from its effect in the constraint and the norms.

5.2. Standing assumptions. Let Δ be a spherical triangulation with mesh size $|\Delta|$ and smallest angle θ_Δ , and $S_d^r(\Delta) = \ker J_0^{(r)}$ the spline space of Section 3.5. The standing assumptions are (A1) Δ covers $\mathbb{S}^2$, $|\text{star}(v)| < 1$ for every vertex and $|\Delta| \leq 1/6$, the hypotheses under which the dimension and approximation theory of spherical splines are available (Theorems 13.43, 13.45, 14.25, 14.26); (A2) d is even and $d \geq 3r + 2$ with $r \geq 0$; (A3) $w = e^{2\lambda}$ with λ smooth, normalized by (4.2) and, except for surfaces run in physical units, by (4.3). The parity requirement in (A2) is not technical: $\mathcal{B}_d$ contains the constants if and only if d is even, and by Theorem 13.22

$$(5.3) \quad \mathcal{B}_d = \mathcal{Y}_0 \oplus \mathcal{Y}_2 \oplus \cdots \oplus \mathcal{Y}_d \quad (d \text{ even}).$$

Even d puts the constant function, which fixes the mean-value constraint and is the zero-eigenvalue eigenfunction, in the discrete space; for odd d it is not representable even on a single triangle (Theorem 13.23). The inequality $d \geq 3r + 2$ is needed only for (4.7).

5.3. The discrete problem in broken coefficients. Set $S_h := S_d^r(\Delta) \subset W^{1,2}(\mathbb{S}^2)$; $r = 0$ is already conforming, and higher r buys a smoother solution at fewer degrees of freedom. The discrete problem reads: find $u_h \in S_h$ with $b_w(u_h, 1) = 0$ such that

$$(5.4) \quad a(u_h, v_h) = b_w(f, v_h) \quad \text{for all } v_h \in S_h.$$

We write it in the broken coefficients of Section 3. On each triangle T the element stiffness matrix K_T , the element weighted mass matrix $M_{w,T}$, and the element load vector F_T , indexed by the multi-indices α, β of degree d , are

$$(5.5) \quad \begin{aligned} (K_T)_{\alpha\beta} &= \int_T \nabla_{\mathbb{S}^2} B_\alpha^d \cdot \nabla_{\mathbb{S}^2} B_\beta^d dA, & (M_{w,T})_{\alpha\beta} &= \int_T w B_\alpha^d B_\beta^d dA, \\ (F_T)_\alpha &= \int_T w f B_\alpha^d dA, \end{aligned}$$

with the tangential gradients given by (3.13). The broken stiffness matrix $K = \text{diag}_T K_T$ and the broken weighted mass matrix $M_w = \text{diag}_T M_{w,T}$ are block diagonal of size $N \times N$, one block per triangle, the broken load vector $F \in \mathbb{R}^N$ is the concatenation of the F_T , and for $c, y \in \mathbb{R}^N$ representing $s, v \in S_d^{-1}(\Delta)$ one has $y^\top K c = a(s, v)$,

413 $y^\top M_w c = b_w(s, v)$ and $y^\top F = b_w(f, v)$ with the forms extended triangle by triangle.
 414 Let $c_1 \in \mathbb{R}^N$ be the broken coefficient vector of the constant function 1. Since $1 \in \mathcal{B}_d$
 415 for even d (Theorem 13.23), its B-form on each triangle is unique, and its entries are
 416 not all equal because $b_1 + b_2 + b_3$ is not 1 on the sphere; c_1 is computed once per
 417 triangulation and degree by solving the square Bernstein evaluation system at the
 418 domain points of each triangle, which is exact up to roundoff. Writing $J := J_0^{(r)}$ for
 419 the smoothness matrix, (5.4) is the constrained linear problem

$$420 \quad (5.6) \quad c \in \ker J, \quad c_1^\top M_w c = 0, \quad y^\top K c = y^\top F \quad \text{for all } y \in \ker J.$$

421 The broken stiffness matrix K is the round-sphere Dirichlet matrix, assembled once;
 422 M enters only through w in F and $M_w c_1$. This is the constrained Galerkin problem of
 423 [29] and the constrained B-form setting of [7]: the broken matrices are never modified,
 424 and conformity is a side condition on the coefficient vector; for $w \equiv 1$ it is the
 425 discretization of [10]. It can be solved on the same space by a null-space matrix, by
 426 the augmented Lagrangian iteration of [7], or as a saddle-point system with Lagrange
 427 multipliers [29, Sec. 6]. The experiments use the null-space method [14, Sec. 6],
 428 checked against the saddle-point form in Section 7.1.

429 *The null-space method.* Let $Z \in \mathbb{R}^{N \times n_h}$ have full column rank and range $Z =$
 430 $\ker J$, so that $n_h = \dim S_h$ and every conforming coefficient vector is $c = Z\hat{c}$ for a
 431 unique $\hat{c} \in \mathbb{R}^{n_h}$. Substituting into (5.6) and appending the mean-value constraint
 432 with a scalar multiplier ν gives the reduced system

$$433 \quad (5.7) \quad \begin{bmatrix} Z^\top K Z & g \\ g^\top & 0 \end{bmatrix} \begin{bmatrix} \hat{c} \\ \nu \end{bmatrix} = \begin{bmatrix} Z^\top F \\ 0 \end{bmatrix}, \quad g := Z^\top M_w c_1, \quad c = Z\hat{c}.$$

434 The reduced stiffness matrix $Z^\top K Z$ is symmetric positive semidefinite with kernel
 435 spanned by the reduced coefficients $\hat{c}_1$ of the constant; since $g^\top \hat{c}_1 = \int_{\mathbb{S}^2} w \, dA > 0$,
 436 the bordered matrix is nonsingular. The multiplier ν vanishes whenever the discrete
 437 compatibility $c_1^\top F = 0$ holds to roundoff, and its size checks the quadrature of the
 438 right-hand side. In the language of [34, Ch. 5], the columns of Z extend the coefficients
 439 on a determining set to all N broken coefficients.

440 The construction of Z by sparse rank-revealing elimination of J in a locality order,
 441 the saddle-point realization of the same equations, and the reading of residuals in the
 442 broken representation (the raw residual $Kc - F$ is a Lagrange multiplier in range $J^\top$
 443 and vanishes only after projection by $Z^\top$) are given in Section SM2 of the supplement.

444 **5.4. What error to expect.** The rates observed in Section 7 follow from the
 445 approximation estimate (4.8) by standard arguments, which we state with their hy-
 446 potheses.

447 **PROPOSITION 5.1.** *Let $w \in C^\infty(\mathbb{S}^2)$ with $0 < w_{\min} \leq w \leq w_{\max}$, let (A1) and*
 448 *(A2) hold, and let $u \in W^{d+1,2}(\mathbb{S}^2)$ solve (1.1). Then the solution u_h of (5.4) satisfies*
 449

$$450 \quad (5.8) \quad |u - u_h|_{1,2,M} \leq C |\Delta|^d |u|_{d+1,2,\mathbb{S}^2}, \quad \|u - u_h\|_{L^2(M)} \leq C |\Delta|^{d+1} |u|_{d+1,2,\mathbb{S}^2},$$

451 *with C depending on d , θ_Δ , $w_{\min}$, $w_{\max}$ and $\|w\|_{C^1(\mathbb{S}^2)}$. For the eigenproblem (5.11),*
 452 *if μ is an eigenvalue of (5.9) with eigenspace $E \subset W^{d+1,2}(\mathbb{S}^2)$, then the discrete*
 453 *eigenvalues converging to μ satisfy $0 \leq \mu_h - \mu \leq C |\Delta|^{2d}$.*

454 *Proof.* The energy estimate is Céa's lemma [19, Ch. 2] on the w -mean-zero sub-
 455 space, where a is coercive by Proposition 5.2, with (4.8) and (4.6). For the L^2 estimate

let $e = u - u_h$ made w -mean-free and let z solve $-\Delta_{\mathbb{S}^2} z = we$, $b_w(z, 1) = 0$; elliptic regularity on the closed manifold $\mathbb{S}^2$ [38, Ch. 5] gives $\|z\|_{W^{2,2}} \leq C w_{\max} \|e\|_{L^2}$. Galerkin orthogonality gives $b_w(e, e) = a(z - z_h, e)$ for every $z_h \in S_h$, (4.7) with $m = 1$ bounds $\inf_{z_h} \|z - z_h\|_{1,2}$ by $C|\Delta| \|z\|_{W^{2,2}}$, and dividing by $\|e\|_{L_w^2} \geq w_{\min}^{1/2} \|e\|_{L^2}$ and using the energy estimate gives the L^2 rate in the weighted norm, which is the $L^2(M)$ norm by (4.5). The eigenvalue statement is the Babuška–Osborn estimate for conforming approximation of a symmetric pencil with compact solution operator [8, Sec. 7]: the eigenvalue error is bounded by the square of the best energy approximation of the eigenspace, which (4.8) bounds by $|\Delta|^d$. $\square$

The growth of the constants with the conformal factor and with the position along the spectrum is the subject of a separate paper.

5.5. The Laplace–Beltrami eigenproblem. The eigenvalue problem on M is: find $u \not\equiv 0$ and $\mu \in \mathbb{R}$ with

$$(5.9) \quad -\Delta_M u = \mu u \quad \text{on } M,$$

understood weakly in $W^{1,2}(M)$, and on the round sphere the generalized eigenvalue problem for the pencil (a, b_w) : find $u \not\equiv 0$ and $\mu \in \mathbb{R}$ with

$$(5.10) \quad a(u, v) = \mu b_w(u, v) \quad \text{for all } v \in W^{1,2}(\mathbb{S}^2).$$

PROPOSITION 5.2 (The spectrum as a weighted problem). *(u, μ) is an eigenpair of (5.9) if and only if it is an eigenpair of (5.10), and no constant in this correspondence depends on λ .*

Proof. Insert (4.5) and (4.6) into the weak form of (5.9). $\square$

The pencil is symmetric with b_w positive definite, so the spectrum is real, discrete, $0 = \mu_0 < \mu_1 \leq \mu_2 \leq \dots \rightarrow \infty$ [20, Ch. I], with b_w -orthogonal eigenfunctions and the constant eigenfunction at $\mu_0 = 0$. The discrete eigenvalue problem reads: find $u_h \in S_h \setminus \{0\}$ and $\mu_h \in \mathbb{R}$ with

$$(5.11) \quad a(u_h, v_h) = \mu_h b_w(u_h, v_h) \quad \text{for all } v_h \in S_h,$$

which in broken coefficients is the constrained pencil $y^\top K c = \mu_h y^\top M_w c$ for all $y \in \ker J$, and with the null-space matrix the reduced pencil

$$(5.12) \quad (Z^\top K Z) \hat{c} = \mu_h (Z^\top M_w Z) \hat{c}, \quad c = Z \hat{c},$$

in which $Z^\top M_w Z$ is positive definite. The saddle-point form is the pencil

$$(5.13) \quad \mathcal{A}x = \mu \mathcal{M}x, \quad \mathcal{A} = \begin{bmatrix} K & J_b^\top \\ J_b & 0 \end{bmatrix}, \quad \mathcal{M} = \begin{bmatrix} M_w & 0 \\ 0 & 0 \end{bmatrix};$$

its finite eigenvalues with nonzero primal component are exactly those of (5.11), and the multiplier directions contribute only infinite eigenvalues [29, Sec. 6], so shift-invert Arnoldi iteration [35] on $(\mathcal{A} - \sigma \mathcal{M})^{-1} \mathcal{M}$ finds the constrained spectrum without any null-space matrix. Three facts follow from the structure. First, since $S_h \subset W^{1,2}(\mathbb{S}^2)$, the Courant–Fischer principle [20, Ch. I] applied over subspaces of S_h gives the one-sided bounds

$$(5.14) \quad \mu_{i,h} \geq \mu_i, \quad i = 0, 1, \dots, n_h - 1.$$

Second, since d is even, the constants lie in S_h by (3.10), and $a(1, v_h) = 0 = \mu b_w(1, v_h)$ shows that $(1, 0)$ is an exact discrete eigenpair, $\mu_{0,h} = 0 = \mu_0$. Third, the same principle applied to a perturbed weight gives a spectrally uniform stability estimate: if $\hat{w}$ is the weight actually used and $\delta := \|\hat{w}/w - 1\|_{L^\infty(\mathbb{S}^2)} < 1$, then $(1 - \delta)b_w \leq b_{\hat{w}} \leq (1 + \delta)b_w$ as quadratic forms, and the monotonicity of the Courant–Fischer minimum in the mass form gives

$$(5.15) \quad \frac{\mu_i}{1 + \delta} \leq \hat{\mu}_i \leq \frac{\mu_i}{1 - \delta}, \quad i \geq 0,$$

for the eigenvalues $\hat{\mu}_i$ of $(a, b_{\hat{w}})$; the same argument covers quadrature errors in K and M_w . Errors in the weight and the quadrature therefore enter the spectrum as relative errors uniform along the spectrum. Convergence at the rate $|\Delta|^{2d}$ is Proposition 5.1, with no geometric consistency term, in contrast with the surface finite element eigenvalue theory of [18]. One structural statement is specific to the sphere and serves as an implementation test.

PROPOSITION 5.3 (Parity fingerprint). *Let $\lambda \equiv 0$, so that $w \equiv 1$ and (5.11) is the discrete eigenvalue problem of the round metric, and let d be even. Then for every even ℓ with $0 \leq \ell \leq d$, every spherical harmonic $Y \in \mathcal{Y}_\ell$ is an exact discrete eigenfunction with exact eigenvalue $\ell(\ell + 1)$:*

$$a(Y, v_h) = \ell(\ell + 1) b_1(Y, v_h) \quad \text{for all } v_h \in S_h,$$

so the discrete spectrum contains $\ell(\ell + 1)$ with multiplicity at least $2\ell + 1$, and for $|\Delta|$ small enough the discrete eigenvalue cluster converging to $\ell(\ell + 1)$ has total multiplicity exactly $2\ell + 1$. The eigenvalues of odd order ℓ converge from above at the rate $|\Delta|^{2d}$.

Proof. By (5.3) and (3.10), $\mathcal{Y}_\ell \subset \mathcal{B}_d \subset S_h$ for even $\ell \leq d$, and $-\Delta_{\mathbb{S}^2} Y = \ell(\ell + 1)Y$ for $Y \in \mathcal{Y}_\ell$ [6, Ch. 2], so $(Y, \ell(\ell + 1))$ satisfies (5.11) verbatim; the $2\ell + 1$ harmonics of degree ℓ are independent in S_h , which gives the multiplicity bound. By [8, Sec. 7], for $|\Delta|$ small enough exactly $2\ell + 1$ discrete eigenvalues, counted with multiplicity, lie in a neighborhood of $\ell(\ell + 1)$ that excludes the other eigenvalues of the sphere, and these are the cluster. $\square$

At $d = 6$ the discrete spectrum of the round metric therefore contains 0, 6, 20 and 42 with multiplicities 1, 5, 9 and 13 up to quadrature error, while the clusters at 2, 12 and 30 converge at twelfth order; a quadrature defect perturbs the even clusters, a constraint-rank defect changes the multiplicities, and an assembly error destroys the rate. For every λ , Hersch’s inequality $\mu_1(M) \text{Area}(M) \leq 8\pi$ [27] and its extension $\mu_k(M) \text{Area}(M) \leq 8\pi k$ [30] bound the eigenvalues from above; since the discrete eigenvalues lie above the exact ones by (5.14), a normalized discrete eigenvalue exceeding $2k$ signals underresolution or a failure of the normalization (4.2).

6. The algorithm. Algorithm 6.1 lists the computation in execution order. The conformal factor is obtained and checked before any equation is discretized, and the smoothness, null-space and stiffness matrices are built once per triangulation and degree and reused for every λ .

The formulas each step evaluates, the Newton iteration and half-angle form for the conformal factor of a surface of revolution, the radial projection of the reference quadrature rule, the quadrature formulas for the load and constraint vectors, the solver diagnostics, and the cost of a new surface or a new right-hand side are given in Section SM3 of the supplement. J , Z and $Z^\top K Z$ are computed once per (Δ, d, r) ; a new λ costs one pass over the quadrature points and one factorization of (5.7); a new

Algorithm 6.1 Poisson equation for a conformal metric on the sphere by spherical splines

Require: the conformal factor $w = e^{2\lambda}$, as a formula, or as a conformal parametrization Φ or a generating curve from which it is computed; right-hand side f ; spherical triangulation Δ ; degree d (even) and smoothness r

Ensure: broken coefficient vector c of the spline solution $u_h \approx u$

- 1: **Geometry.** Build the routine $x \mapsto w(x)$: for a surface of revolution solve (4.12) for $\eta(\Theta)$ and evaluate (4.13); for a parametrized surface evaluate (4.10); for a prescribed λ evaluate $e^{2\lambda}$. Check $\int_{\mathbb{S}^2} w dA$ against the exact area when one is known and the Hersch moments (4.2) against zero.
 - 2: **Spline space (once per Δ, d, r).** Assemble the edge blocks (3.15) into J ; eliminate to reduced row echelon form in the locality order and form Z by (SM2.1); compare $N - \text{rank } J$ with the closed-form counts of Section SM1.
 - 3: **Stiffness (once per Δ, d).** Assemble the broken $K = \text{diag}_T K_T$ from (5.5) with the tangential gradient (3.13); form and store $Z^\top K Z$.
 - 4: **Weighted data.** At the quadrature points of each triangle evaluate w, f (for an embedded surface, $f \circ \Phi$) and the Bernstein basis; assemble the load vector F and the constraint vector $M_w c_1$ of (5.5), where c_1 interpolates the constant. Check $c_1^\top F$ against zero.
 - 5: **Solve.** Solve the bordered system (5.7) by a sparse direct solver; set $c = Z\hat{c}$. Check the multiplier ν against zero and the projected residual $Z^\top(Kc - F)$ against zero.
 - 6: **Output.** Evaluate u_h from c by de Casteljau on any triangle; for an embedded surface the value at $p \in \Sigma$ is $u_h(\Phi^{-1}(p))$. Errors are measured in $L^2(M)$ and $H^1(M)$ through (4.5) and (4.6).
-

right-hand side costs one pass and one triangular solve; the eigenproblem costs one assembly of M_w and one Lanczos run.

7. Numerical experiments. The smoothness matrix J , the null-space matrix Z and the stiffness matrix K are reused unchanged across instances; the metric is supplied through the conformal factor of step 1 of Algorithm 6.1. Local matrices use a conical Gauss product rule of order 12 per direction on each spherical triangle. The bordered system (5.7) is solved by a sparse direct solver, and the reduced pencil (5.12) by shift-inverted Lanczos at $\sigma = -1$. Icosahedral level ℓ has $20 \cdot 4^\ell$ triangles and geodesic mesh sizes 0.628, 0.326, 0.165, 0.083 at levels 1 to 4; the degrees are $(d, r) = (2, 0), (4, 0)$ and $(6, 1)$, the last the smallest even degree admitted by $d \geq 3r + 2$ for C^1 splines.

The instances are the round metric, $\lambda \equiv 0$; three surfaces of revolution, the prolate spheroid $x^2 + y^2 + (z/c)^2 = 1$ with $c = 3/2$, the dumbbell $R(\psi) = 1 - \frac{3}{4}\sin^2 \psi$ with two bulbs of radius one joined by a neck of radius $1/4$, and the Evans–Fung red blood cell [25], a biconcave discocyte run in physical units (micrometers), whose factors are computed from the generating curves by (4.11), with the spheroid’s Mercator integral in closed form (7.1); and a prescribed factor with no embedding, $\lambda = 0.6(x_1^2 - x_2^2) - 0.4x_3^2$, whose symmetry satisfies (4.2). The three surfaces of revolution pass the area identity to relative error below 3×10^{-15} ; $\text{osc } \lambda$ is 0, 0.535, 2.23, 0.373 and 1.2 respectively, and the dumbbell is the hard case, where $e^{2 \text{osc } \lambda} \approx 86$ enters the constants of (4.8) (Table SM2 in the supplement).

7.1. Algebraic checks. The mesh, quadrature, smoothness matrix, null-space matrix and solvers are checked independently of the metric: Euler relations, quadrature area, rank and dependency count of J , reproduction of the constant and of xy , agreement of the null-space and saddle-point realizations to 6×10^{-14} , one-sidedness (5.14) to 3×10^{-13} , and product-to-moment assembly [1] against direct quadrature to 10^{-15} (Section SM4 of the supplement).

7.2. The Poisson equation. Two tests separate the two places where the metric enters. In Test 1 the exact solution is $u = e^{x_1}x_2 + x_1x_3^3$, not a spherical polynomial of any degree, with $f = -\Delta_M u = -w^{-1}\Delta_{\mathbb{S}^2}u$ computed symbolically; the product wf in the load vector is then $-\Delta_{\mathbb{S}^2}u$, so the discrete solutions for the different instances differ only through the constraint vector $M_w c_1$ and the weighted norm in which the error is measured, and Test 1 isolates the spline approximation from the metric. In Test 2 the right-hand side is prescribed on the embedded surface, $f(p) = p_1$ made mean-free, so the load vector sees the metric at every quadrature point; its reference is a dense spherical-harmonic Galerkin solve of (5.2). Errors are measured through (4.5) and (4.6) with the assembly quadrature. Table 1 reports Test 1 and Table 2 the degree-4 and 6 rows of Test 2, whose full record is in Section SM5 of the supplement.

TABLE 1

Poisson problem, Test 1: exact solution $u = \tilde{u} \circ \Phi^{-1}$ with $\tilde{u} = e^{x_1}x_2 + x_1x_3^3$ on the sphere, right-hand side $f = -\Delta_M u$, icosahedral levels 1 to 4. The $H^1(M)$ seminorm error is the same on all four surfaces, because the load vector $\int_{\mathbb{S}^2} w \tilde{f} B_\alpha dA = -\int_{\mathbb{S}^2} (\Delta_{\mathbb{S}^2} \tilde{u}) B_\alpha dA$ does not contain w ; the $L^2(M)$ error sees w through the constant fixed by the weighted mean and through the weighted norm, and is listed per surface. Rates are against the geodesic mesh size; the $L^2(M)$ rates on the spheroid, dumbbell and red blood cell differ from those of the sphere by at most 0.16 at level 2 and by less than 0.03 at levels 3 and 4. The red blood cell is in micrometers.

(d, r)	lev.	dim S_h	$H^1(M)$, all surfaces		$L^2(M)$, sphere		$L^2(M)$, other surfaces		
			error	rate	error	rate	spheroid	dumbbell	red blood cell
(2, 0)	2	642	$4.59 \cdot 10^{-2}$	2.04	$1.75 \cdot 10^{-3}$	3.04	$2.09 \cdot 10^{-3}$	$1.33 \cdot 10^{-3}$	$5.56 \cdot 10^{-3}$
	3	2562	$1.16 \cdot 10^{-2}$	2.02	$2.24 \cdot 10^{-4}$	3.02	$2.69 \cdot 10^{-4}$	$1.72 \cdot 10^{-4}$	$7.11 \cdot 10^{-4}$
	4	10242	$2.92 \cdot 10^{-3}$	2.01	$2.82 \cdot 10^{-5}$	3.02	$3.39 \cdot 10^{-5}$	$2.18 \cdot 10^{-5}$	$8.96 \cdot 10^{-5}$
(4, 0)	2	2562	$1.70 \cdot 10^{-4}$	4.19	$3.70 \cdot 10^{-6}$	5.20	$4.46 \cdot 10^{-6}$	$2.77 \cdot 10^{-6}$	$1.17 \cdot 10^{-5}$
	3	10242	$1.07 \cdot 10^{-5}$	4.06	$1.17 \cdot 10^{-7}$	5.07	$1.42 \cdot 10^{-7}$	$8.84 \cdot 10^{-8}$	$3.71 \cdot 10^{-7}$
	4	40962	$6.70 \cdot 10^{-7}$	4.03	$3.69 \cdot 10^{-9}$	5.04	$4.45 \cdot 10^{-9}$	$2.78 \cdot 10^{-9}$	$1.16 \cdot 10^{-8}$
(6, 1)	2	3206	$1.40 \cdot 10^{-6}$	6.04	$2.57 \cdot 10^{-8}$	6.92	$3.25 \cdot 10^{-8}$	$2.38 \cdot 10^{-8}$	$7.98 \cdot 10^{-8}$
	3	12806	$2.49 \cdot 10^{-8}$	5.92	$2.48 \cdot 10^{-10}$	6.82	$3.14 \cdot 10^{-10}$	$2.30 \cdot 10^{-10}$	$7.69 \cdot 10^{-10}$
	4	51206	$4.22 \cdot 10^{-10}$	5.94	$2.25 \cdot 10^{-12}$	6.84	$2.82 \cdot 10^{-12}$	$1.98 \cdot 10^{-12}$	$7.03 \cdot 10^{-12}$

In Test 1 the $H^1(M)$ rates over the last refinement are 2.0, 4.0 and 5.9 for $d = 2, 4, 6$, the same to the digits shown across the instances, and the $L^2(M)$ rates are 3.0, 5.0 and 6.8, with the $d = 6$ errors reaching 2×10^{-12} in $L^2(M)$ at 51206 unknowns; the conformal factor enters the $L^2(M)$ errors as a multiplicative constant within the bound E of Lemma 4.1 and nothing else, and the prescribed factor behaves like the embedded surfaces (Section SM5 of the supplement). In Test 2, where the weight is in the load vector, the rates on the sphere, spheroid and red blood cell are again d and $d + 1$ to two digits at $d = 2, 4$ and 5.6 to 6.1 in energy at $d = 6$; on the dumbbell the finest (6, 1) row and one (2, 0) row fall short by an amount within the accuracy of the spherical-harmonic reference, as discussed in the supplement. The multiplier ν is below 10^{-13} and the projected residual below 10^{-12} in every row.

Comparison with surface finite elements. Test 1 was repeated on the spheroid and the dumbbell with parametric surface finite elements [23, 21]: Dziuk's polyhedral P_1

TABLE 2

Poisson problem, Test 2: right-hand side $f(p) = p_1$ prescribed on the physical surface, so that w enters the load vector at every quadrature point; degrees 4 and 6 (degree 2 and the reference details are in Section SM5 of the supplement). Errors in $L^2(M)$ and the $H^1(M)$ seminorm with rates in parentheses; the red blood cell is in micrometers. The dumbbell reference is converged only to about 10^{-7} , which limits its last rows.

geometry	(d, r)	level	$\dim S_h$	$L^2(M)$ (rate)	$H^1(M)$ (rate)
sphere	(4, 0)	2	2562	1.64×10^{-6} (5.21)	7.50×10^{-5} (4.19)
	(4, 0)	3	10242	5.20×10^{-8} (5.05)	4.72×10^{-6} (4.05)
	(6, 1)	2	3206	9.75×10^{-9} (6.96)	5.63×10^{-7} (6.07)
spheroid	(4, 0)	2	2562	2.44×10^{-6} (5.04)	9.69×10^{-5} (4.10)
	(4, 0)	3	10242	8.39×10^{-8} (4.94)	6.35×10^{-6} (3.99)
	(6, 1)	2	3206	3.50×10^{-8} (6.32)	1.36×10^{-6} (5.60)
dumbbell	(4, 0)	2	2562	1.45×10^{-5} (4.30)	3.75×10^{-4} (3.44)
	(4, 0)	3	10242	4.05×10^{-7} (5.23)	2.32×10^{-5} (4.07)
	(6, 1)	2	3206	8.46×10^{-7} (4.95)	2.79×10^{-5} (4.23)
red blood cell	(4, 0)	2	2562	2.27×10^{-4} (5.14)	3.29×10^{-3} (4.13)
	(4, 0)	3	10242	7.57×10^{-6} (4.98)	2.15×10^{-4} (3.99)
	(6, 1)	2	3206	4.16×10^{-6} (6.63)	6.47×10^{-5} (5.70)

method and isoparametric P_2 and P_3 elements on the same icosahedral triangulations pushed to M by Φ , every Lagrange node on the exact surface, so that the geometric error is the smallest the parametric method admits at that degree; errors are measured on the discrete surface with the lifted exact solution. Table 3 reports errors and the wall-clock time of one Python implementation for both methods at the finest level run. At equal degrees of freedom the C^0 spline of degree 2 and the P_2 element are within a factor of two; the spline of degree 4 has an $L^2(M)$ error an order of magnitude below that of P_3 at 10^4 unknowns; and the C^1 spline of degree 6 reaches 3×10^{-10} where P_3 reaches 10^{-6} with twice the unknowns. The finite element errors on the dumbbell are two to three times those on the spheroid at the same level, consistent with the larger geometric approximation error on the more strongly curved surface, whereas the spline energy errors agree on the two instances to the digits shown, as expected when the stiffness matrix is independent of the metric and the map carries the geometry. In time, the per-instance work of the spline at a given number of unknowns is comparable to P_1 and two to three times P_2 , the difference being the order-12 product quadrature of the weighted load vector; the metric-independent setup is amortized over instances. The ratios at the finest configurations are in Table 3.

7.3. The eigenproblem on the round sphere: the parity fingerprint.

Table 4 reports the discrete spectra on the round sphere ($w \equiv 1$) against the exact eigenvalues $\ell(\ell+1)$ with multiplicities $2\ell+1$, split into the even- ℓ and odd- ℓ clusters of Proposition 5.3. The prediction is realized exactly: the even clusters are reproduced to at worst 2.5×10^{-13} , independent of the mesh, at all three degrees, while the odd clusters converge from above with measured rates 4.02, 8.35, and 11.95 against the expected $2d = 4, 8, 12$ of Section 5.5. At $d = 6$ the exact values 0, 6, 20, 42 appear with multiplicities 1, 5, 9, 13. At level 1 the vertex stars have diameter about 1.26, violating $|\text{star}(v)| < 1$ in (A1), and the fingerprint and rates hold regardless.

TABLE 3

Conformal spherical splines against parametric surface finite elements (Dziuk P_1 , isoparametric P_2 , P_3 , nodes on the exact surface) for Poisson Test 1 on the spheroid and the dumbbell: finest level run for each degree, errors in $L^2(M)$ and the $H^1(M)$ seminorm with, in parentheses, the convergence rate over the last refinement, and wall-clock seconds. For the splines, “setup is the geometry-independent work (smoothness matrix, null-space matrix, round-sphere stiffness), done once per triangulation and degree and reused for every surface; “surface is the per-surface work (weight at the quadrature points, load and constraint vectors, bordered solve). For the FEM, “setup is the isoparametric mesh and “surface is assembly plus solve. Same machine, single thread, quadrature of order 12 (splines) and 6 (FEM).

geometry	method	dof	$L^2(M)$	$H^1(M)$	setup (s)	surface (s)
spheroid	spline (2, 0)	10242	$3.39 \cdot 10^{-5}$ (3.00)	$2.92 \cdot 10^{-3}$ (2.00)	2.3	12.4
	spline (4, 0)	40962	$4.45 \cdot 10^{-9}$ (5.01)	$6.70 \cdot 10^{-7}$ (4.01)	3.1	14.4
	spline (6, 1)	12806	$3.14 \cdot 10^{-10}$ (6.79)	$2.49 \cdot 10^{-8}$ (5.90)	1.5	5.5
	FEM P_1	10242	$1.44 \cdot 10^{-3}$ (2.00)	$9.20 \cdot 10^{-2}$ (1.00)	0.8	12.9
	FEM P_2	40962	$4.77 \cdot 10^{-6}$ (3.00)	$9.29 \cdot 10^{-4}$ (2.00)	1.1	15.1
	FEM P_3	23042	$1.05 \cdot 10^{-6}$ (4.02)	$1.53 \cdot 10^{-4}$ (3.00)	0.4	4.4
dumbbell	spline (2, 0)	10242	$2.17 \cdot 10^{-5}$ (3.00)	$2.92 \cdot 10^{-3}$ (2.00)	2.2	24.2
	spline (4, 0)	40962	$2.78 \cdot 10^{-9}$ (5.01)	$6.70 \cdot 10^{-7}$ (4.01)	3.2	25.6
	spline (6, 1)	12806	$2.30 \cdot 10^{-10}$ (6.80)	$2.49 \cdot 10^{-8}$ (5.90)	1.4	8.4
	FEM P_1	10242	$1.00 \cdot 10^{-3}$ (1.99)	$1.43 \cdot 10^{-1}$ (1.00)	0.8	25.4
	FEM P_2	40962	$3.26 \cdot 10^{-6}$ (3.06)	$2.55 \cdot 10^{-3}$ (1.99)	1.2	27.7
	FEM P_3	23042	$2.14 \cdot 10^{-6}$ (4.04)	$6.42 \cdot 10^{-4}$ (2.96)	0.5	7.4

TABLE 4

Round sphere. Maximum eigenvalue errors over the even- ℓ and odd- ℓ clusters (first 16 eigenvalues for $d = 2, 4$; first 50 for $d = 6$), and the measured convergence rate of the odd clusters against the theoretical $2d$. The even clusters are exact up to quadrature and roundoff at every mesh, as Proposition 5.3 predicts.

d	r	level	$\dim S_d^r$	even- ℓ err	odd- ℓ err	rate ($2d$)
2	0	1	162	1.7×10^{-14}	4.94×10^{-2}	
2	0	2	642	3.4×10^{-14}	3.39×10^{-3}	4.09
2	0	3	2562	2.0×10^{-14}	2.18×10^{-4}	4.02
4	0	1	642	1.7×10^{-14}	6.83×10^{-6}	
4	0	2	2562	1.8×10^{-14}	2.88×10^{-8}	8.35
6	1	1	806	1.5×10^{-12}	4.29×10^{-6}	
6	1	2	3206	2.5×10^{-13}	1.71×10^{-9}	11.95

7.4. The eigenproblem on the curved surfaces. For the spheroid the Mercator matching (4.12) reads

$$(7.1) \quad \ln \tan(\theta/2) + G_0(\theta) = \ln \tan(\Theta/2), \quad G_0(\theta) = \int_{\pi/2}^{\theta} \frac{\sqrt{\cos^2 t + c^2 \sin^2 t} - 1}{\sin t} dt,$$

the isometric latitude of the Mercator projection of an ellipsoid, and $\int_{\mathbb{S}^2} w dA$ agrees with the closed-form area $2\pi + 2\pi c \arcsin(e)/e$, $e = \sqrt{1 - 1/c^2}$, to relative error 8.4×10^{-16} . None of the three spectra has an elementary closed form, so the reference is a dense Galerkin solve of the same pencil in the real spherical-harmonic basis, in which the zonal weight couples only equal azimuthal orders; its self-convergence is given in Table 5. The normalized first eigenvalues $\mu_1 \text{Area}/(4\pi)$ are 1.5514, 0.780 and 1.784, all below the Hersch bound 2; on the red blood cell the equatorial pair

626 $\mu_{1,2} = 0.16715 \mu\text{m}^{-2}$ lies below the axial mode $\mu_3 = 0.23963$, so in-plane variation is
 627 cheaper than top-to-bottom variation on the flat discocyte.

628 Table 5 gives the rates. On the spheroid ($\text{osc } \lambda = 0.535$) they are 3.98, 7.99 and
 629 10.62 for $d = 2, 4, 6$, the last at an error of 2×10^{-8} where the quadrature of the weight
 630 is not yet negligible. On the red blood cell ($\text{osc } \lambda = 0.373$) they are 3.98, 8.02 and
 631 10.65 with terminal errors 1.3×10^{-10} and 3.4×10^{-11} at $d = 4, 6$. On the dumbbell
 632 ($\text{osc } \lambda = 2.23$) the coarse errors are orders of magnitude above the spheroid values at
 633 identical dimensions and the rates climb toward $2d$ with refinement (3.94 at $d = 2$;
 634 6.31 then 7.00 at $d = 4$; 9.03 at $d = 6$ from one pair of levels), consistent with the
 635 expected order and a pre-asymptotic range that widens with $\text{osc } \lambda$. One-sidedness
 636 holds to 6×10^{-14} throughout. Figure 1 shows computed eigenfunctions of the three
 637 embedded instances.




FIG. 1. *Computed eigenfunctions at $d = 6$, $r = 1$, level 2, drawn on the embedded surfaces. From left: spheroid, seventh eigenfunction, $\mu = 5.38$; dumbbell, $\mu = 57.62$, with azimuthal and axial nodal lines crossing both bulbs; red blood cell, a mode localized in the dimples, $\mu = 1.28 \mu\text{m}^{-2}$; red blood cell, an azimuthal rim mode, $\mu = 2.66 \mu\text{m}^{-2}$. Sign and scale are arbitrary.*

TABLE 5

Eigenvalues on the curved surfaces (levels 2 and 3; level 1 in the supplement's record). Maximum eigenvalue error of the spline pencil against the spherical-harmonic reference (first 16 eigenvalues; first 30 at $d = 6$ on the spheroid) and measured rates against $2d$. Reference self-convergence: 9.4×10^{-13} (spheroid), 8.4×10^{-8} (dumbbell), 2.2×10^{-13} (red blood cell, in μm^{-2}).

geometry	d	r	level	$\dim S_d^r$	max error	rate ($2d$)
spheroid	2	0	2	642	4.47×10^{-3}	4.27
	2	0	3	2562	2.94×10^{-4}	3.98
	4	0	2	2562	3.63×10^{-7}	7.99
	6	1	2	3206	1.95×10^{-8}	10.62
dumbbell	2	0	2	642	1.28×10^0	2.63
	2	0	3	2562	8.68×10^{-2}	3.94
	4	0	2	2562	4.64×10^{-3}	6.31
	4	0	3	10242	3.88×10^{-5}	7.00
	6	1	2	3206	3.62×10^{-5}	9.03
red blood cell	2	0	2	642	5.84×10^{-4}	4.26
	2	0	3	2562	3.85×10^{-5}	3.98
	4	0	2	2562	3.10×10^{-8}	7.70
	4	0	3	10242	1.29×10^{-10}	8.02
	6	1	2	3206	3.35×10^{-11}	10.65

638 **8. Discussion and conclusions.** For a conformal metric on the sphere the
 639 area element is the conformal factor times the round one and the Dirichlet energy is
 640 the round energy, so the stiffness matrix is the round-sphere matrix for every such

metric and the geometry is one scalar at the quadrature points of the load vector, the constraint vector and the weighted mass matrix. By uniformization every smooth closed genus-zero surface is such a metric, and an embedded surface with a conformal parametrization, a surface of revolution, a prescribed factor and a meshed surface are instances that differ only in how the factor is obtained. The factor is checked against an exact area where one exists before any equation is discretized, the C^r spline spaces serve every metric as they stand, and errors in the metric are measured in the round one with no change of constants in energy. The experiments show the rates $d + 1$ in $L^2(M)$ and d in energy for the Poisson problem and $2d$ for the eigenvalues on all instances, reproduction of the even-order eigenvalues of the round metric to roundoff, and, on the dumbbell with $\text{osc } \lambda = 2.23$, larger constants and a longer pre-asymptotic range with the limiting orders unchanged.

Relative to surface finite elements [23, 24, 17], the method replaces a sequence of physical meshes by a fixed spherical mesh plus one scalar function. It is restricted to genus zero, needs the conformal factor as input, and a metric with strong conformal concentration maps large regions into small spherical patches that a uniform mesh under-resolves; in return, when the factor is known exactly, the geometry is low-dimensional, computed to the accuracy of the area check, and, in the cases run here, at degrees four and six the errors are smaller than those of isoparametric cubic elements at comparable numbers of unknowns. Adaptive refinement driven by the conformal factor is the next step. Two matters are deferred to separate papers: the constants in (5.8) and in the eigenvalue estimate as functions of the conformal factor and the position along the spectrum, and surfaces given only as triangle meshes, for which the conformal factor must itself be computed.

Acknowledgments. Claude Fable 5.1 (Anthropic) assisted with drafting, editing, and code; the mathematics, numerical results, final text, and source verification are the author's responsibility.

REFERENCES

- [1] M. AINSWORTH, G. ANDRIAMARO, AND O. DAVYDOV, *Bernstein-Bézier finite elements of arbitrary order and optimal assembly procedures*, SIAM J. Sci. Comput., 33 (2011), pp. 3087–3109.
- [2] P. ALFELD, M. NEAMTU, AND L. L. SCHUMAKER, *Circular Bernstein-Bézier polynomials*, in Mathematical Methods for Curves and Surfaces, M. Dæhlen, T. Lyche, and L. L. Schumaker, eds., Vanderbilt University Press, Nashville, 1995, pp. 11–20.
- [3] P. ALFELD, M. NEAMTU, AND L. L. SCHUMAKER, *Bernstein-Bézier polynomials on spheres and sphere-like surfaces*, Comput. Aided Geom. Design, 13 (1996), pp. 333–349.
- [4] P. ALFELD, M. NEAMTU, AND L. L. SCHUMAKER, *Dimension and local bases of homogeneous spline spaces*, SIAM J. Math. Anal., 27 (1996), pp. 1482–1501.
- [5] P. ALFELD, M. NEAMTU, AND L. L. SCHUMAKER, *Fitting scattered data on sphere-like surfaces using spherical splines*, J. Comput. Appl. Math., 73 (1996), pp. 5–43.
- [6] K. ATKINSON AND W. HAN, *Spherical Harmonics and Approximations on the Unit Sphere: An Introduction*, vol. 2044 of Lecture Notes in Mathematics, Springer, Berlin, 2012.
- [7] G. AWANOU, M.-J. LAI, AND P. WENSTON, *The multivariate spline method for scattered data fitting and numerical solution of partial differential equations*, in Wavelets and Splines: Athens 2005, G. Chen and M.-J. Lai, eds., Nashboro Press, Brentwood, TN, 2006, pp. 24–75.
- [8] I. BABUŠKA AND J. OSBORN, *Eigenvalue problems*, in Handbook of Numerical Analysis, Vol. II, P. G. Ciarlet and J. L. Lions, eds., North-Holland, Amsterdam, 1991, pp. 641–787.
- [9] V. BARAMIDZE AND M.-J. LAI, *Error bounds for minimal energy interpolatory spherical splines*, in Approximation Theory XI: Gatlinburg 2004, C. K. Chui, M. Neamtu, and L. L. Schumaker, eds., Nashboro Press, Brentwood, TN, 2005, pp. 25–50.
- [10] V. BARAMIDZE AND M.-J. LAI, *Spherical spline solution to a PDE on the sphere*, in Wavelets

- and Splines: Athens 2005, G. Chen and M.-J. Lai, eds., Nashboro Press, Brentwood, TN, 2006, pp. 75–92.
- [11] V. BARAMIDZE AND M.-J. LAI, *Convergence of discrete and penalized least squares spherical splines*, J. Approx. Theory, 163 (2011), pp. 1091–1106.
- [12] V. BARAMIDZE AND M. J. LAI, *Nonnegative data interpolation by spherical splines*, J. Comput. Appl. Math., 342 (2018), pp. 463–477.
- [13] V. BARAMIDZE, M.-J. LAI, AND C. K. SHUM, *Spherical splines for data interpolation and fitting*, SIAM J. Sci. Comput., 28 (2006), pp. 241–259.
- [14] M. BENZI, G. H. GOLUB, AND J. LIESEN, *Numerical solution of saddle point problems*, Acta Numer., 14 (2005), pp. 1–137.
- [15] D. BOFFI, *Finite element approximation of eigenvalue problems*, Acta Numer., 19 (2010), pp. 1–120.
- [16] A. BONITO AND A. DEMLOW, *A posteriori error estimates for the Laplace–Beltrami operator on parametric c^2 surfaces*, SIAM J. Numer. Anal., 57 (2019), pp. 973–996.
- [17] A. BONITO, A. DEMLOW, AND R. H. NOCHETTO, *Finite element methods for the Laplace–Beltrami operator*, in Geometric Partial Differential Equations, Part I, vol. 21 of Handbook of Numerical Analysis, Elsevier, Amsterdam, 2020, pp. 1–103.
- [18] A. BONITO, A. DEMLOW, AND J. OWEN, *A priori error estimates for finite element approximations to eigenvalues and eigenfunctions of the Laplace–Beltrami operator*, SIAM J. Numer. Anal., 56 (2018), pp. 2963–2988.
- [19] S. C. BRENNER AND L. R. SCOTT, *The Mathematical Theory of Finite Element Methods*, Springer, New York, 3rd ed., 2008.
- [20] I. CHAVEL, *Eigenvalues in Riemannian Geometry*, Academic Press, Orlando, 1984.
- [21] A. DEMLOW, *Higher-order finite element methods and pointwise error estimates for elliptic problems on surfaces*, SIAM J. Numer. Anal., 47 (2009), pp. 805–827.
- [22] A. DEMLOW AND G. DZIUK, *An adaptive finite element method for the Laplace–Beltrami operator on implicitly defined surfaces*, SIAM J. Numer. Anal., 45 (2007), pp. 421–442.
- [23] G. DZIUK, *Finite elements for the Beltrami operator on arbitrary surfaces*, in Partial Differential Equations and Calculus of Variations, vol. 1357 of Lecture Notes in Mathematics, Springer, Berlin, 1988, pp. 142–155.
- [24] G. DZIUK AND C. M. ELLIOTT, *Finite element methods for surface PDEs*, Acta Numer., 22 (2013), pp. 289–396.
- [25] E. EVANS AND Y.-C. FUNG, *Improved measurements of the erythrocyte geometry*, Microvasc. Res., 4 (1972), pp. 335–347.
- [26] X. GU AND S.-T. YAU, *Global conformal surface parameterization*, in Proceedings of the Eurographics/ACM SIGGRAPH Symposium on Geometry Processing, 2003, pp. 127–137.
- [27] J. HERSCH, *Quatre propriétés isopérimétriques de membranes sphériques homogènes*, C. R. Acad. Sci. Paris Sér. A, 270 (1970), pp. 1645–1648.
- [28] J. JOST, *Compact Riemann Surfaces: An Introduction to Contemporary Mathematics*, Universitext, Springer, Berlin, 3rd ed., 2006.
- [29] S. KAPITA, *Bernstein constraint complexes for multivariate splines on triangulated surfaces*. Preprint, 2026.
- [30] M. KARPUKHIN, N. NADIRASHVILI, A. PENSKOI, AND I. POLTEROVICH, *An isoperimetric inequality for Laplace eigenvalues on the sphere*, J. Differential Geom., 118 (2021), pp. 313–333.
- [31] M. KAZHDAN, J. SOLOMON, AND M. BEN-CHEN, *Can mean-curvature flow be modified to be non-singular?*, Comput. Graph. Forum, 31 (2012), pp. 1745–1754.
- [32] M.-J. LAI AND J. LEE, *A multivariate spline based collocation method for numerical solution of partial differential equations*, SIAM J. Numer. Anal., 60 (2022), pp. 2405–2434.
- [33] M.-J. LAI AND J. LEE, *Trivariate spline collocation methods for numerical solution to 3D Monge–Ampère equation*, J. Sci. Comput., 95 (2023), p. 56.
- [34] M.-J. LAI AND L. L. SCHUMAKER, *Spline Functions on Triangulations*, Cambridge University Press, Cambridge, 2007.
- [35] R. B. LEHOUCQ, D. C. SORENSEN, AND C. YANG, *ARPACK Users’ Guide: Solution of Large-Scale Eigenvalue Problems with Implicitly Restarted Arnoldi Methods*, SIAM, Philadelphia, 1998.
- [36] P. LI AND S.-T. YAU, *A new conformal invariant and its applications to the Willmore conjecture and the first eigenvalue of compact surfaces*, Invent. Math., 69 (1982), pp. 269–291.
- [37] M. NEAMTU AND L. L. SCHUMAKER, *On the approximation order of splines on spherical triangulations*, Adv. Comput. Math., 21 (2004), pp. 3–20.
- [38] M. E. TAYLOR, *Partial Differential Equations I: Basic Theory*, Springer, New York, 2nd ed., 2011.