

Figure 1: Schedule using EDF.

## 1. Solution

- (a) The utilization factor is given by

$$U = \sum_{i=1}^3 \frac{C_i}{T_i} = \frac{2}{3} + \frac{\tau}{10},$$

and for the tasks to be schedulable using EDF it needs to hold that  $U \leq 1$ , which gives the condition

$$\tau \leq \frac{10}{3}. \quad (1)$$

So the maximum value for  $\tau$  is 3. The schedule length is given by  $\text{lcm}(3, 6, 10) = 30$ .

- (b) The schedule is shown in Figure 1. Task  $J_2$  is assumed to have priority when tasks have the same time until deadline. The worst-case response times can be read from the schedule and are given by  $R_1 = 1$ ,  $R_2 = 5$ , and  $R_3 = 9$ .
- (c) The utilization factor with  $\tau = 3$  is given by  $U = \frac{2}{3} + \frac{3}{10} \approx 0.967$ . Since  $U$  is higher than  $3(2^{1/3} - 1) = 0.78$  we cannot determine if the tasks are schedulable by RM. We therefore compute the worst-case response times to determine schedulability. Let  $R_i$  denote the worst-case response time for task  $J_i$ . The fixed priority is assigned such that  $J_1$  has the highest priority,  $J_2$  has the middle priority, and  $J_3$  has the lowest priority.

For task  $J_1$ :

$$R_1 = C_1 = 1 \leq D_1 = 3$$

For task  $J_2$ :

$$R_2^0 = C_2 = 2$$

$$R_2^1 = C_2 + \lceil \frac{R_2^0}{T_1} \rceil C_1 = 2 + \lceil \frac{2}{3} \rceil = 3$$

$$R_2^2 = C_2 + \lceil \frac{R_2^1}{T_1} \rceil C_1 = 2 + \lceil \frac{3}{3} \rceil = 3 \leq D_2 = 6$$

For task  $J_3$ :

$$R_3^0 = C_3 = 3$$

$$R_3^1 = C_3 + \lceil \frac{R_3^0}{T_1} \rceil C_1 + \lceil \frac{R_3^0}{T_2} \rceil C_2 = 3 + \lceil \frac{3}{3} \rceil + 2\lceil \frac{3}{6} \rceil = 6$$

$$R_3^2 = C_3 + \lceil \frac{R_3^1}{T_1} \rceil C_1 + \lceil \frac{R_3^1}{T_2} \rceil C_2 = 3 + \lceil \frac{6}{3} \rceil + 2\lceil \frac{6}{6} \rceil = 7$$

$$R_3^3 = C_3 + \lceil \frac{R_3^2}{T_1} \rceil C_1 + \lceil \frac{R_3^2}{T_2} \rceil C_2 = 3 + \lceil \frac{7}{3} \rceil + 2\lceil \frac{7}{6} \rceil = 10$$

$$R_3^4 = C_3 + \lceil \frac{R_3^3}{T_1} \rceil C_1 + \lceil \frac{R_3^3}{T_2} \rceil C_2 = 3 + \lceil \frac{10}{3} \rceil + 2\lceil \frac{10}{6} \rceil = 11 > D_3 = 10$$

$$R_3^5 = C_3 + \lceil \frac{R_3^4}{T_1} \rceil C_1 + \lceil \frac{R_3^4}{T_2} \rceil C_2 = 3 + \lceil \frac{11}{3} \rceil + 2\lceil \frac{11}{6} \rceil = 11 > D_3 = 10$$

Thus the three tasks are not schedulable by RM, since task  $J_3$  does not meet its deadline.

## 2. Solution

- (a) (1) The switching system can be modeled as a Hybrid Automaton

$$H = (Q, X, \text{Init}, f, E, D, G, R),$$

where

- $Q = \{q_1, q_2\}$ ,
- $X = \mathbb{R}^2$ ,
- $\text{Init} \in Q \times X$ ,
- $f(q_1, x) = \begin{bmatrix} -x_2 - x_1(1 - x_1^2 - x_2^2) \\ x_1 - x_2(1 - x_1^2 - x_2^2) \end{bmatrix}$ ,  $f(q_2, x) = \begin{bmatrix} x_2 \\ -(1 + x_1^2)x_1 - x_2 + M \cos(\omega t) \end{bmatrix}$ ,
- $E = \{(q_1, q_2), (q_2, q_1)\}$ ,
- $D(q_1) = \{x \in \mathbb{R}^2 : \|x\| < \alpha\}$ ,  $D(q_2) = \{x \in \mathbb{R}^2 : \|x\| \geq \alpha\}$ ,
- $G((q_1, q_2)) = \{x \in \mathbb{R}^2 : \|x\| = \alpha\}$ ,  $G((q_2, q_1)) = \{x \in \mathbb{R}^2 : \|x\| < \alpha\}$ ,
- $R((q_1, q_2), x) = R((q_2, q_1), x) = x$

- (b) (i) By differentiating  $V_1$  over the traj. of the first subsystem, we obtain

$$\begin{aligned} \dot{V}_1 &= -x_1x_2 - x_1^2(1 - x_1^2 - x_2^2) + x_1x_2 - x_2^2(1 - x_1^2 - x_2^2) \\ &= -(x_1^2 + x_2^2)(1 - x_1^2 - x_2^2) = -\|x\|(1 - \|x\|), \end{aligned}$$

which is negative for  $\|x\| < 1$ .

- (ii) By considering  $x(0)$  such that  $\|x(0)\| > 1$ ,  $\dot{V}_1$  is positive and hence  $V_1$ , and consequently  $\|x\|$ , blow to infinity. Hence, it is not globally AS.

- (c) (i)  $V_2$  becomes

$$V_2 = \frac{1}{2}x_1^2 + x_2^2 + x_1x_2 + x_1^2 + \frac{x_1^4}{2},$$

and after differentiation over the traj. of the second subsystem, we obtain

$$\begin{aligned} \dot{V}_2 &= x_1x_2 - 2x_1x_2(1 + x_1^2) - 2x_2^2 + 2Mx_2 \cos(\omega t) + x_2^2 - x_1^2(1 + x_1^2) - x_1x_2 + \\ &\quad Mx_1 \cos(\omega t) + 2x_1x_2 + 2x_1^3x_2 \\ &= -x_1^2 - x_2^2 + (x_1 + 2x_2)M \cos(\omega t) - x_1^4 = -\|x\|^2 - x_1^4 + (x_1 + 2x_2) \cos(\omega t) \\ &\leq -\|x\|^2 - x_1^4 + \sqrt{5}\|x\|M \cos(\omega t) \\ &= -\|x\|(\|x\| - M\sqrt{5}) - x_1^4, \end{aligned}$$

which is negative when  $\|x\| > M\sqrt{5}$ .

- (ii) For a sufficiently large value of  $M$ , it can be shown that  $\dot{x}_1 > 0$  and  $\dot{x}_2 > 0$  arbitrarily close to zero, hence the origin is not asymptotically stable.

- (d) The condition is

$$M\sqrt{5} < \alpha < 1.$$

### 3. Solution:

- (a) The system is not controllable, because

$$W_c = \begin{bmatrix} 1 & 3 \\ 2 & 6 \end{bmatrix}$$

and  $\det(W_c) = 0$ .

- (b) With  $L = [l_1 \quad l_2]$  the characteristic polynomial of the closed-loop system is

$$\begin{aligned} \det(zI - A + BL) &= \det \left( \begin{bmatrix} z - 1 + l_1 & l_2 - 1 \\ 2(l_1 - 1) & z - 2(1 - l_2) \end{bmatrix} \right) \\ &= (z - 1 + l_1)(z - 2(1 - l_2)) - 2(1 - l_1)(1 - l_2) \\ &= z(z - 3 + l_1 + 2l_2). \end{aligned}$$

One pole is located at  $z_1 = 0$ , while the other is located at  $z_2 = 3 - l_1 - 2l_2$ . The characteristic polynomial of the open-loop system is  $\det(zI - A) = z(z - 3)$  and we observe that one pole cannot be moved by state-feedback. This is due to the uncontrollability of the system.

- (c) If  $\text{rank}([\lambda I - \Phi \quad \Gamma]) < n$  for a  $\lambda \in \mathbb{C}$ , then there exists a  $w$ , such that  $w^T[\lambda I - \Phi \quad \Gamma] = 0$ . Therefore,  $w^T\Phi = \lambda w^T$  and  $w^T\Gamma = 0$ . Note that in this case  $\lambda$  is an eigenvalue of  $\Phi$  and  $w$  is the corresponding left eigenvector. Multiplying  $W_c$  with  $w^T$  from the left gives us

$$\begin{aligned} w^T W_c &= [w^T\Gamma \quad w^T\Phi\Gamma \quad w^T\Phi^2\Gamma \quad \dots \quad w^T\Phi^{n-1}\Gamma] \\ &= [w^T\Gamma \quad \lambda w^T\Gamma \quad \lambda^2 w^T\Gamma \quad \dots \quad \lambda^{n-1} w^T\Gamma] = 0 \end{aligned}$$

which shows us that  $\text{rank}(W_c) < n$ . Hence, the system is uncontrollable.

- (d) We only need to test  $z$ 's that are eigenvalues of  $\Phi$  because only then  $\Phi$  will lose rank. Otherwise,  $zI - \Phi$  has full rank and therefore  $[zI - \Phi \quad \Gamma]$  has full rank.
- (e) As shown in (b) the poles of the system are at  $z_1 = 0$  and  $z_2 = 3$ . Applying the PBH test to the given system show us that

$$\text{rank}([z_1 I - \Phi \quad \Gamma]) = \text{rank} \left( \begin{bmatrix} -1 & -1 & 1 \\ -2 & -2 & 2 \end{bmatrix} \right) = 1$$

and

$$\text{rank}([z_2 I - \Phi \quad \Gamma]) = \text{rank} \left( \begin{bmatrix} 2 & -1 & 1 \\ -2 & 1 & 2 \end{bmatrix} \right) = 2.$$

Hence, the system is uncontrollable. The PBH test does not only show when a system is controllable but also which poles can be moved by a feedback controller. This implies that we can still stabilize an uncontrollable system if all unstable poles are controllable.

#### 4. Solution

(a) Consider the relation:

$$\sim = \{(s_1, s'_1), (s_2, s'_2), (s_3, s'_3), (s_4, s'_4), (s_5, s'_5), (s_2, s'_4), (s_6, s'_6)\} \subseteq S \times S',$$

The relation  $\sim$  is a simulation relation from  $\mathcal{T}_1$  to  $\mathcal{T}_2$  since the following hold:

- (a) For the state  $s_1 \in S$ , there exists the state  $s'_1 \in S'$  such that  $(s_1, s'_1) \in \sim$ .
- (b) We have that  $(s_6, s'_6) \in \sim$  with  $s_6 \in S_F$  and  $s'_6 \in S'_F$ .
- (c)
  - $(s_1, s'_1) \in \sim, (s_1, a, s_2) \in \rightarrow$ , and at the same time  $(s'_1, a, s'_2) \in \rightarrow'$ , and  $(s_2, s'_2) \in \sim$ .
  - $(s_1, s'_1) \in \sim, (s_1, a, s_3) \in \rightarrow$ , and at the same time  $(s'_1, a, s'_3) \in \rightarrow'$ , and  $(s_3, s'_3) \in \sim$ .
  - $(s_2, s'_2) \in \sim, (s_2, a, s_5) \in \rightarrow$ , and at the same time  $(s'_2, a, s'_5) \in \rightarrow'$ , and  $(s_5, s'_5) \in \sim$ .
  - $(s_2, s'_4) \in \sim, (s_2, a, s_5) \in \rightarrow$ , and at the same time  $(s'_4, a, s'_5) \in \rightarrow'$ , and  $(s_5, s'_5) \in \sim$ .
  - $(s_3, s'_3) \in \sim, (s_3, b, s_5) \in \rightarrow$ , and at the same time  $(s'_3, b, s'_5) \in \rightarrow'$ , and  $(s_5, s'_5) \in \sim$ .
  - $(s_4, s'_5) \in \sim, (s_4, a, s_6) \in \rightarrow$ , and at the same time  $(s'_5, a, s'_6) \in \rightarrow'$ , and  $(s_6, s'_6) \in \sim$ .
  - $(s_5, s'_5) \in \sim, (s_5, a, s_6) \in \rightarrow$ , and at the same time  $(s'_5, a, s'_6) \in \rightarrow'$ , and  $(s_6, s'_6) \in \sim$ .

It can be shown analogously, that the three conditions for the relation  $\sim^{-1}$  hold. Therefore,  $\sim$  is a *bisimulation relation* and the Transition Systems  $\mathcal{T}_1, \mathcal{T}_2$  are *bisimilar*.

(b) We start with the partition  $S_{/\sim} = \{\{s_1, s_2, s_3, s_4, s_5\}, \{s_6\}\}$ .

**Step 1:** Set  $P = \{s_1, s_2, s_3, s_4, s_5\}$  and  $P' = \{s_6\}$ . We have that  $\emptyset \neq P \cap \text{Pre}_a(P') = \{s_1, s_2, s_3, s_4, s_5\} \cap \{s_4, s_5\} = \{s_4, s_5\} \neq P$ . Then, set:

$$\begin{aligned} P_1 &= P \cap \text{Pre}_a(P') = \{s_4, s_5\}, \\ P_2 &= P \setminus \text{Pre}_a(P') = \{s_1, s_2, s_3\}, \\ S_{/\sim} &= \{\{s_6\}, \{s_4, s_5\}, \{s_1, s_2, s_3\}\}. \end{aligned}$$

**Step 2:** Set  $P = P' = \{s_1, s_2, s_3\}$ . We have that  $\emptyset \neq P \cap \text{Pre}_a(P) = \{s_1, s_2, s_3\} \cap \{s_1\} = \{s_1\} \neq P$ . Then, set:

$$\begin{aligned} P_1 &= P \cap \text{Pre}_a(P') = \{s_1\}, \\ P_2 &= P \setminus \text{Pre}_a(P') = \{s_1, s_2, s_3\} \setminus \{s_1\} = \{s_2, s_3\}, \\ S_{/\sim} &= \{\{s_1\}, \{s_6\}, \{s_2, s_3\}, \{s_4, s_5\}, \}. \end{aligned}$$

**Step 3:** Set  $P = \{s_2, s_3\}$ ,  $P' = \{s_4, s_5\}$ . We have that

$$\emptyset \neq P \cap \text{Pre}_b(P') = \{s_2, s_3\} \cap \{s_3\} = \{s_3\} \neq P.$$

Then, set:

$$\begin{aligned} P_1 &= P \cap \text{Pre}_b(P') = \{s_3\}, \\ P_2 &= P \setminus \text{Pre}_b(P') = \{s_2, s_3\} \setminus \{s_3\} = \{s_2\}, \\ S_{/\sim} &= \{\{s_1\}, \{s_2\}, \{s_3\}, \{s_4, s_5\}, \{s_6\}\}. \end{aligned}$$

We next check if the algorithm is finished, i.e., that we cannot refine more. The only element which is not a singleton is  $\{s_4, s_5\}$ . Thus, it suffices to show that for all  $P' \in S_{/\sim}$  and  $\sigma \in \{a, b\}$  it holds that  $\{s_4, s_5\} \cap \text{Pre}_\sigma(P') = \emptyset$  or  $\{s_4, s_5\}$ , which holds, since the only  $P'$  for which  $\{s_4, s_5\} \cap \text{Pre}_\sigma(P') \neq \emptyset$  is  $P' = \{s_6\}$  and also  $\{s_4, s_5\} \cap \text{Pre}_\sigma(P') = \{s_4, s_5\}$ .




Figure 2: The transition system  $\mathcal{T}'_3$ .

- (c) (i) No, it is not a bisimulation relation because it is not a relation from  $\mathcal{T}_3$  to  $\mathcal{T}_4$ . Indeed, since  $(s_2, s''_2) \in \sim$  and  $s_2 \xrightarrow{b} s_2$ , there must be a  $r \in S'$  such that  $s''_2 \xrightarrow{b} r$  and  $(s_2, r) \in \sim$ , which is not the case.
- (ii) The transition system  $\mathcal{T}'_3$  is depicted in Fig. 2. The relation

$$\sim = \{(s_1, s'_1), (s_1, s''_1), (s_2, s'_2), (s_2, s''_2), (s_3, s'_3)\} \subseteq S \times S',$$

is a simulation relation from  $\mathcal{T}_3$  to  $\mathcal{T}'_3$  since the following hold:

- i. For the state  $s_1 \in S$ , there exists a state  $s'_1 \in S'$  such that  $(s_1, s'_1) \in \sim$ .
- ii. We have that  $(s_3, s'_3) \in \sim$  with  $s_3 \in S_F$  and  $s'_3 \in S'_F$ .
- iii.
  - $(s_1, s'_1) \in \sim$ ,  $(s_1, a, s_2) \in \rightarrow$ , and at the same time  $(s'_1, a, s'_2) \in \rightarrow'$ , and  $(s_2, s'_2) \in \sim$ .
  - $(s_1, s'_1) \in \sim$ ,  $(s_1, b, s_1) \in \rightarrow$ , and at the same time  $(s'_1, b, s'_1) \in \rightarrow'$ , and  $(s_1, s'_1) \in \sim$ .
  - $(s_1, s''_1) \in \sim$ ,  $(s_1, a, s_2) \in \rightarrow$ , and at the same time  $(s''_1, a, s''_2) \in \rightarrow'$ , and  $(s_2, s''_2) \in \sim$ .
  - $(s_1, s''_1) \in \sim$ ,  $(s_1, b, s_1) \in \rightarrow$ , and at the same time  $(s''_1, b, s'_1) \in \rightarrow'$ , and  $(s_1, s'_1) \in \sim$ .
  - $(s_2, s'_2) \in \sim$ ,  $(s_2, a, s_3) \in \rightarrow$ , and at the same time  $(s'_2, a, s'_3) \in \rightarrow'$ , and  $(s_3, s'_3) \in \sim$ .
  - $(s_2, s'_2) \in \sim$ ,  $(s_2, b, s_2) \in \rightarrow$ , and at the same time  $(s'_2, b, s''_2) \in \rightarrow'$ , and  $(s_2, s''_2) \in \sim$ .
  - $(s_2, s''_2) \in \sim$ ,  $(s_2, a, s_3) \in \rightarrow$ , and at the same time  $(s''_2, a, s'_3) \in \rightarrow'$ , and  $(s_3, s'_3) \in \sim$ .
  - $(s_2, s''_2) \in \sim$ ,  $(s_2, b, s_2) \in \rightarrow$ , and at the same time  $(s''_2, b, s'_2) \in \rightarrow'$ , and  $(s_2, s'_2) \in \sim$ .

It can be shown analogously, that the three conditions for the relation  $\sim^{-1}$  hold. Therefore,  $\sim$  is a *bisimulation relation* and the Transition Systems  $\mathcal{T}_3, \mathcal{T}'_3$  are *bisimilar*.

- (d) Such a Transition system is  $\mathcal{T}'_5$  in Fig. 3. We can check the simulation by relating  $s_1, s_2, s_3 \sim s'_1, s''_1$  and  $s_4 \sim s'_2$ .




Figure 3: The transition system  $\mathcal{T}'_5$ .

## 5. Solution

(a) We have

$$\begin{aligned}\dot{x}_1 &= \dot{p}_2 - \dot{p}_1 = u_2 - u_1 \\ \dot{x}_2 &= \dot{p}_3 - \dot{p}_2 = u_3 - u_2.\end{aligned}\tag{2}$$

Therefore,

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ u_3 \end{bmatrix}.\tag{3}$$

(b) We have

$$\begin{aligned}u_1 &= p_2 - p_1 = x_1, \\ u_2 &= p_3 - p_2 = x_2, \\ u_3 &= p_1 - p_3 = -x_1 - x_2.\end{aligned}\tag{4}$$

Therefore

$$\begin{bmatrix} u_1 \\ u_2 \\ u_3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ -1 & -1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}.\tag{5}$$

(c) Using the results in (a) and (b), we have

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} -1 & 1 \\ -1 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}.\tag{6}$$

(d) Using the suggested Lyapunov function, we find

$$\dot{V}(x) = -x_1^2 - 2x_2^2,\tag{7}$$

therefore,  $x(t)$  asymptotically converges to zero, which implies that the robots asymptotically meet at the same place.

(e) With the sampled measurements, we have  $\dot{x} = BK(x + e)$ , or equivalently

$$\begin{aligned}\dot{x}_1 &= -x_1 + x_2 - e_1 + e_2, \\ \dot{x}_2 &= -x_1 - 2x_2 - e_1 - 2e_2.\end{aligned}\tag{8}$$

(f) Using the same Lyapunov function, we have

$$\dot{V}(t) = -x_1^2 - x_1e_1 + x_1e_2 - 2x_2^2 - e_1x_2 - 2e_2x_2,\tag{9}$$



which can be written as

$$\dot{V}(t) = -x(t)^T Q x(t) + x(t)^T R e(t), \quad (10)$$

where

$$Q = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}, \quad R = \begin{bmatrix} -1 & 1 \\ -1 & -2 \end{bmatrix}. \quad (11)$$

Therefore,  $\dot{V}(t)$  is guaranteed if

$$-\lambda_Q \|x(t)\|^2 + \|x(t)\| \lambda_R \|e(t)\| \leq 0, \quad (12)$$

where  $\lambda_Q$  is the smallest eigenvalue of  $Q$  and  $\lambda_R$  is the 2-norm of  $R$ . Solving the inequality for  $\|e(t)\|$  gives

$$\|e(t)\| \leq \frac{\lambda_Q}{\lambda_R} \|x(t)\|. \quad (13)$$

As for numerical values, we have immediately  $\lambda_Q = 1$ , while a quick calculation gives  $\lambda_R = \sqrt{(7 + \sqrt{13})}/2$ . Therefore, we conclude

$$\alpha = \sqrt{\frac{2}{7 + \sqrt{13}}}. \quad (14)$$