

MEM6804 Modeling and Simulation for Logistics & Supply Chain

物流与供应链建模与仿真

Theory Analysis

Lecture 2: Elements of Probability and Statistics

SHEN Haihui 沈海辉

Sino-US Global Logistics Institute
Shanghai Jiao Tong University

🏠 shenhaihui.github.io/teaching/mem6804f
✉ shenhaihui@sjtu.edu.cn

Spring 2021 (full-time)



上海交通大学
SHANGHAI JIAO TONG UNIVERSITY

董浩云航运与物流研究院
CY TUNG Institute of Maritime and Logistics
中美物流研究院 (工程系统管理研究院)
Sino-US Global Logistics Institute (Institute of Industrial & System Engineering)



- 1 Probability Space
- 2 Random Variables & Distributions
- 3 Expectations
- 4 Common Distributions
- 5 Useful Inequalities
- 6 Convergence
- 7 Properties of a Random Sample

- 1 Probability Space
- 2 Random Variables & Distributions
- 3 Expectations
- 4 Common Distributions
- 5 Useful Inequalities
- 6 Convergence
- 7 Properties of a Random Sample

Probability Space

A **probability space** is a triplet $(\Omega, \mathcal{F}, \mathbb{P})$:

- Ω , sample space: A set of *all* possible outcomes.



Probability Space

A **probability space** is a triplet $(\Omega, \mathcal{F}, \mathbb{P})$:

- Ω , sample space: A set of *all* possible outcomes.
 - A set of *some* outcomes, as a subset of Ω , is called an **event**.

A **probability space** is a triplet $(\Omega, \mathcal{F}, \mathbb{P})$:

- Ω , sample space: A set of *all* possible outcomes.
 - A set of *some* outcomes, as a subset of Ω , is called an **event**.
- \mathcal{F} , σ -algebra (or σ -field): A set of events, i.e., a set of some subsets of Ω , such that:

A **probability space** is a triplet $(\Omega, \mathcal{F}, \mathbb{P})$:

- Ω , sample space: A set of *all* possible outcomes.
 - A set of *some* outcomes, as a subset of Ω , is called an **event**.
- \mathcal{F} , σ -algebra (or σ -field): A set of events, i.e., a set of some subsets of Ω , such that:
 - ① $\Omega \in \mathcal{F}$;

A **probability space** is a triplet $(\Omega, \mathcal{F}, \mathbb{P})$:

- Ω , sample space: A set of *all* possible outcomes.
 - A set of *some* outcomes, as a subset of Ω , is called an **event**.
- \mathcal{F} , σ -algebra (or σ -field): A set of events, i.e., a set of some subsets of Ω , such that:
 - ① $\Omega \in \mathcal{F}$;
 - ② Closed under complementation: If $A \in \mathcal{F}$, then $A^c \in \mathcal{F}$;

A **probability space** is a triplet $(\Omega, \mathcal{F}, \mathbb{P})$:

- Ω , sample space: A set of *all* possible outcomes.
 - A set of *some* outcomes, as a subset of Ω , is called an **event**.
- \mathcal{F} , σ -algebra (or σ -field): A set of events, i.e., a set of some subsets of Ω , such that:
 - ① $\Omega \in \mathcal{F}$;
 - ② Closed under complementation: If $A \in \mathcal{F}$, then $A^c \in \mathcal{F}$;
 - ③ Closed under countable unions:[†] If $A_i \in \mathcal{F}$, $i = 1, 2, \dots$, is a **countable** sequence of sets, then $\bigcup_{i=1}^{\infty} A_i \in \mathcal{F}$.

[†] It implies that \mathcal{F} is also closed under countable intersections.



A **probability space** is a triplet $(\Omega, \mathcal{F}, \mathbb{P})$:

- Ω , sample space: A set of *all* possible outcomes.
 - A set of *some* outcomes, as a subset of Ω , is called an **event**.
- \mathcal{F} , σ -algebra (or σ -field): A set of events, i.e., a set of some subsets of Ω , such that:
 - ① $\Omega \in \mathcal{F}$;
 - ② Closed under complementation: If $A \in \mathcal{F}$, then $A^c \in \mathcal{F}$;
 - ③ Closed under countable unions:[†] If $A_i \in \mathcal{F}$, $i = 1, 2, \dots$, is a **countable** sequence of sets, then $\bigcup_{i=1}^{\infty} A_i \in \mathcal{F}$.
- $\mathbb{P} : \mathcal{F} \rightarrow [0, 1]$, probability function (or probability measure): A function that assigns probabilities to events, such that:

[†] It implies that \mathcal{F} is also closed under countable intersections.



A **probability space** is a triplet $(\Omega, \mathcal{F}, \mathbb{P})$:

- Ω , sample space: A set of *all* possible outcomes.
 - A set of *some* outcomes, as a subset of Ω , is called an **event**.
- \mathcal{F} , σ -algebra (or σ -field): A set of events, i.e., a set of some subsets of Ω , such that:
 - ① $\Omega \in \mathcal{F}$;
 - ② Closed under complementation: If $A \in \mathcal{F}$, then $A^c \in \mathcal{F}$;
 - ③ Closed under countable unions:[†] If $A_i \in \mathcal{F}$, $i = 1, 2, \dots$, is a **countable** sequence of sets, then $\bigcup_{i=1}^{\infty} A_i \in \mathcal{F}$.
- $\mathbb{P} : \mathcal{F} \rightarrow [0, 1]$, probability function (or probability measure): A function that assigns probabilities to events, such that:
 - ① $\mathbb{P}(A) \in [0, 1]$ for any $A \in \mathcal{F}$;

[†] It implies that \mathcal{F} is also closed under countable intersections.



A **probability space** is a triplet $(\Omega, \mathcal{F}, \mathbb{P})$:

- Ω , sample space: A set of *all* possible outcomes.
 - A set of *some* outcomes, as a subset of Ω , is called an **event**.
- \mathcal{F} , σ -algebra (or σ -field): A set of events, i.e., a set of some subsets of Ω , such that:
 - ① $\Omega \in \mathcal{F}$;
 - ② Closed under complementation: If $A \in \mathcal{F}$, then $A^c \in \mathcal{F}$;
 - ③ Closed under countable unions:[†] If $A_i \in \mathcal{F}$, $i = 1, 2, \dots$, is a **countable** sequence of sets, then $\bigcup_{i=1}^{\infty} A_i \in \mathcal{F}$.
- $\mathbb{P} : \mathcal{F} \rightarrow [0, 1]$, probability function (or probability measure): A function that assigns probabilities to events, such that:
 - ① $\mathbb{P}(A) \in [0, 1]$ for any $A \in \mathcal{F}$;
 - ② $\mathbb{P}(\Omega) = 1$;

[†] It implies that \mathcal{F} is also closed under countable intersections.



A **probability space** is a triplet $(\Omega, \mathcal{F}, \mathbb{P})$:

- Ω , sample space: A set of *all* possible outcomes.
 - A set of *some* outcomes, as a subset of Ω , is called an **event**.
- \mathcal{F} , σ -algebra (or σ -field): A set of events, i.e., a set of some subsets of Ω , such that:
 - ① $\Omega \in \mathcal{F}$;
 - ② Closed under complementation: If $A \in \mathcal{F}$, then $A^c \in \mathcal{F}$;
 - ③ Closed under countable unions:[†] If $A_i \in \mathcal{F}$, $i = 1, 2, \dots$, is a **countable** sequence of sets, then $\bigcup_{i=1}^{\infty} A_i \in \mathcal{F}$.
- $\mathbb{P} : \mathcal{F} \rightarrow [0, 1]$, probability function (or probability measure): A function that assigns probabilities to events, such that:
 - ① $\mathbb{P}(A) \in [0, 1]$ for any $A \in \mathcal{F}$;
 - ② $\mathbb{P}(\Omega) = 1$;
 - ③ Countably additive: If $A_i \in \mathcal{F}$, $i = 1, 2, \dots$, is a **countable** sequence of **disjoint** sets, then $\mathbb{P}(\bigcup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} \mathbb{P}(A_i)$.

[†] It implies that \mathcal{F} is also closed under countable intersections.



- Example 1: Flip a fair coin.
 - $\Omega = \{\text{H (head)}, \text{T (tail)}\};$
 - $\mathcal{F} = \{\emptyset, \{\text{H}\}, \{\text{T}\}, \Omega\};$
 - $\mathbb{P}(\emptyset) = 0, \mathbb{P}(\{\text{H}\}) = 1/2, \mathbb{P}(\{\text{T}\}) = 1/2, \text{ and } \mathbb{P}(\Omega) = 1.$

- Example 1: Flip a fair coin.
 - $\Omega = \{H \text{ (head)}, T \text{ (tail)}\}$;
 - $\mathcal{F} = \{\emptyset, \{H\}, \{T\}, \Omega\}$;
 - $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\{H\}) = 1/2$, $\mathbb{P}(\{T\}) = 1/2$, and $\mathbb{P}(\Omega) = 1$.
- Example 2: Draw a ball out of 3 balls (red, green, blue).
 - $\Omega = \{R \text{ (red)}, G \text{ (green)}, B \text{ (blue)}\}$;
 - $\mathcal{F} = \{\emptyset, \{R\}, \{G\}, \{B\}, \{R,G\}, \{R,B\}, \{G,B\}, \Omega\}$;
 - $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\{R\}) = \mathbb{P}(\{G\}) = \mathbb{P}(\{B\}) = 1/3$,
 $\mathbb{P}(\{R,G\}) = \mathbb{P}(\{R,B\}) = \mathbb{P}(\{G,B\}) = 2/3$, and $\mathbb{P}(\Omega) = 1$;



- Example 1: Flip a fair coin.
 - $\Omega = \{H \text{ (head)}, T \text{ (tail)}\}$;
 - $\mathcal{F} = \{\emptyset, \{H\}, \{T\}, \Omega\}$;
 - $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\{H\}) = 1/2$, $\mathbb{P}(\{T\}) = 1/2$, and $\mathbb{P}(\Omega) = 1$.
- Example 2: Draw a ball out of 3 balls (red, green, blue).
 - $\Omega = \{R \text{ (red)}, G \text{ (green)}, B \text{ (blue)}\}$;
 - $\mathcal{F} = \{\emptyset, \{R\}, \{G\}, \{B\}, \{R,G\}, \{R,B\}, \{G,B\}, \Omega\}$;
 - $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\{R\}) = \mathbb{P}(\{G\}) = \mathbb{P}(\{B\}) = 1/3$,
 $\mathbb{P}(\{R,G\}) = \mathbb{P}(\{R,B\}) = \mathbb{P}(\{G,B\}) = 2/3$, and $\mathbb{P}(\Omega) = 1$;
 - $\mathcal{F}_1 = \{\emptyset, \{R\}, \{G,B\}, \Omega\}$, $\mathcal{F}_2 = \{\emptyset, \{G\}, \{R,B\}, \Omega\}$...

- Example 1: Flip a fair coin.
 - $\Omega = \{H \text{ (head)}, T \text{ (tail)}\}$;
 - $\mathcal{F} = \{\emptyset, \{H\}, \{T\}, \Omega\}$;
 - $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\{H\}) = 1/2$, $\mathbb{P}(\{T\}) = 1/2$, and $\mathbb{P}(\Omega) = 1$.
- Example 2: Draw a ball out of 3 balls (red, green, blue).
 - $\Omega = \{R \text{ (red)}, G \text{ (green)}, B \text{ (blue)}\}$;
 - $\mathcal{F} = \{\emptyset, \{R\}, \{G\}, \{B\}, \{R,G\}, \{R,B\}, \{G,B\}, \Omega\}$;
 - $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\{R\}) = \mathbb{P}(\{G\}) = \mathbb{P}(\{B\}) = 1/3$,
 $\mathbb{P}(\{R,G\}) = \mathbb{P}(\{R,B\}) = \mathbb{P}(\{G,B\}) = 2/3$, and $\mathbb{P}(\Omega) = 1$;
 - $\mathcal{F}_1 = \{\emptyset, \{R\}, \{G,B\}, \Omega\}$, $\mathcal{F}_2 = \{\emptyset, \{G\}, \{R,B\}, \Omega\}$...
- Example 3: Randomly “draw” a number in $[0, 1]$
 - $\Omega = [0, 1]$;
 - $\mathcal{F}_1 = \{\emptyset, [0, a), [a, 1], \Omega\}$, $\mathcal{F}_2 = \{\emptyset, (0, a), \{0\} \cup [a, 1], \Omega\}$...



- Example 1: Flip a fair coin.
 - $\Omega = \{H \text{ (head)}, T \text{ (tail)}\};$
 - $\mathcal{F} = \{\emptyset, \{H\}, \{T\}, \Omega\};$
 - $\mathbb{P}(\emptyset) = 0, \mathbb{P}(\{H\}) = 1/2, \mathbb{P}(\{T\}) = 1/2, \text{ and } \mathbb{P}(\Omega) = 1.$
- Example 2: Draw a ball out of 3 balls (red, green, blue).
 - $\Omega = \{R \text{ (red)}, G \text{ (green)}, B \text{ (blue)}\};$
 - $\mathcal{F} = \{\emptyset, \{R\}, \{G\}, \{B\}, \{R,G\}, \{R,B\}, \{G,B\}, \Omega\};$
 - $\mathbb{P}(\emptyset) = 0, \mathbb{P}(\{R\}) = \mathbb{P}(\{G\}) = \mathbb{P}(\{B\}) = 1/3,$
 $\mathbb{P}(\{R,G\}) = \mathbb{P}(\{R,B\}) = \mathbb{P}(\{G,B\}) = 2/3, \text{ and } \mathbb{P}(\Omega) = 1;$
 - $\mathcal{F}_1 = \{\emptyset, \{R\}, \{G,B\}, \Omega\}, \mathcal{F}_2 = \{\emptyset, \{G\}, \{R,B\}, \Omega\}...$
- Example 3: Randomly “draw” a number in $[0, 1]$
 - $\Omega = [0, 1];$
 - $\mathcal{F}_1 = \{\emptyset, [0, a), [a, 1], \Omega\}, \mathcal{F}_2 = \{\emptyset, (0, a), \{0\} \cup [a, 1], \Omega\}...$
 - A more practical and interesting \mathcal{F} is the one that contains all intervals (no matter open or closed) on $[0, 1]$.



- **Independence** of Events: Two events A and B in \mathcal{F} are called statistically independent events when

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \mathbb{P}(B).$$

- **Independence** of Events: Two events A and B in \mathcal{F} are called statistically independent events when

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \mathbb{P}(B).$$

- **Conditional Probability**: If A and B are events in \mathcal{F} and $\mathbb{P}(B) > 0$, then the conditional probability of A given B , denoted as $\mathbb{P}(A|B)$, is

$$\mathbb{P}(A|B) := \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}.$$



- **Independence** of Events: Two events A and B in \mathcal{F} are called statistically independent events when

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \mathbb{P}(B).$$

- **Conditional Probability**: If A and B are events in \mathcal{F} and $\mathbb{P}(B) > 0$, then the conditional probability of A given B , denoted as $\mathbb{P}(A|B)$, is

$$\mathbb{P}(A|B) := \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}.$$

- Bayes' Rule:

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(B|A) \mathbb{P}(A)}{\mathbb{P}(B)}.$$



- **Independence** of Events: Two events A and B in \mathcal{F} are called statistically independent events when

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \mathbb{P}(B).$$

- **Conditional Probability**: If A and B are events in \mathcal{F} and $\mathbb{P}(B) > 0$, then the conditional probability of A given B , denoted as $\mathbb{P}(A|B)$, is

$$\mathbb{P}(A|B) := \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}.$$

- Bayes' Rule:

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(B|A) \mathbb{P}(A)}{\mathbb{P}(B)}.$$

- Events A and B are independent $\iff \mathbb{P}(A|B) = \mathbb{P}(A)$.



- For more than two events:
 - **Mutual independence** (or collective independence) intuitively means that each event is independent of any combination of other events;
 - **Pairwise independence** means any two events in the collection are independent of each other.

- For more than two events:
 - **Mutual independence** (or collective independence) intuitively means that each event is independent of any combination of other events;
 - **Pairwise independence** means any two events in the collection are independent of each other.
- Sets A_1, \dots, A_n are (mutually) independent if for any $I \subset \{1, \dots, n\}$ we have $\mathbb{P}(\cap_{i \in I} A_i) = \prod_{i \in I} \mathbb{P}(A_i)$.



- For more than two events:
 - **Mutual independence** (or collective independence) intuitively means that each event is independent of any combination of other events;
 - **Pairwise independence** means any two events in the collection are independent of each other.
- Sets A_1, \dots, A_n are (mutually) independent if for any $I \subset \{1, \dots, n\}$ we have $\mathbb{P}(\cap_{i \in I} A_i) = \prod_{i \in I} \mathbb{P}(A_i)$.
- **Warning:** Only having $\mathbb{P}(\cap_{i=1}^n A_i) = \prod_{i=1}^n \mathbb{P}(A_i)$ is not sufficient!

- For more than two events:
 - **Mutual independence** (or collective independence) intuitively means that each event is independent of any combination of other events;
 - **Pairwise independence** means any two events in the collection are independent of each other.
- Sets A_1, \dots, A_n are (mutually) independent if for any $I \subset \{1, \dots, n\}$ we have $\mathbb{P}(\cap_{i \in I} A_i) = \prod_{i \in I} \mathbb{P}(A_i)$.
- **Warning:** Only having $\mathbb{P}(\cap_{i=1}^n A_i) = \prod_{i=1}^n \mathbb{P}(A_i)$ is not sufficient!
- Sets A_1, \dots, A_n are pairwise independent if for any $i \neq j$ we have $\mathbb{P}(A_i \cap A_j) = \mathbb{P}(A_i) \mathbb{P}(A_j)$.

- For more than two events:
 - **Mutual independence** (or collective independence) intuitively means that each event is independent of any combination of other events;
 - **Pairwise independence** means any two events in the collection are independent of each other.
- Sets A_1, \dots, A_n are (mutually) independent if for any $I \subset \{1, \dots, n\}$ we have $\mathbb{P}(\cap_{i \in I} A_i) = \prod_{i \in I} \mathbb{P}(A_i)$.
- **Warning:** Only having $\mathbb{P}(\cap_{i=1}^n A_i) = \prod_{i=1}^n \mathbb{P}(A_i)$ is not sufficient!
- Sets A_1, \dots, A_n are pairwise independent if for any $i \neq j$ we have $\mathbb{P}(A_i \cap A_j) = \mathbb{P}(A_i) \mathbb{P}(A_j)$.
- Clearly, mutual independence implies pairwise independence, but not vice versa!

Consider a sequence of sets $\{A_n : n \geq 1\}$.

Consider a sequence of sets $\{A_n : n \geq 1\}$.

(The First) Borel-Cantelli Lemma

If $\sum_{n=1}^{\infty} \mathbb{P}(A_n) < \infty$, then $\mathbb{P}(A_n \text{ i.o.}) = 0$, where “i.o.” denotes “infinitely often”.



Consider a sequence of sets $\{A_n : n \geq 1\}$.

(The First) Borel-Cantelli Lemma

If $\sum_{n=1}^{\infty} \mathbb{P}(A_n) < \infty$, then $\mathbb{P}(A_n \text{ i.o.}) = 0$, where “i.o.” denotes “infinitely often”.

The Secon Borel-Cantelli Lemma

If $\sum_{n=1}^{\infty} \mathbb{P}(A_n) = \infty$ and $\{A_n\}$ are independent,[†] then $\mathbb{P}(A_n \text{ i.o.}) = 1$.

[†]The assumption of independence can be weakened to pairwise independence, with more difficult proof.



Consider a sequence of sets $\{A_n : n \geq 1\}$.

(The First) Borel-Cantelli Lemma

If $\sum_{n=1}^{\infty} \mathbb{P}(A_n) < \infty$, then $\mathbb{P}(A_n \text{ i.o.}) = 0$, where “i.o.” denotes “infinitely often”.

The Secon Borel-Cantelli Lemma

If $\sum_{n=1}^{\infty} \mathbb{P}(A_n) = \infty$ and $\{A_n\}$ are independent,[†] then $\mathbb{P}(A_n \text{ i.o.}) = 1$.

- Remark: For event A , if $\mathbb{P}(A) = 1$, then we say A happens **almost surely** (a.s.).

[†]The assumption of independence can be weakened to pairwise independence, with more difficult proof.



- 1 Probability Space
- 2 Random Variables & Distributions**
- 3 Expectations
- 4 Common Distributions
- 5 Useful Inequalities
- 6 Convergence
- 7 Properties of a Random Sample

- A **random variable** (RV) is a function from a sample space Ω into the set of real numbers \mathbb{R} .

- A **random variable** (RV) is a function from a sample space Ω into the set of real numbers \mathbb{R} .
- Formally, given the probability space $(\Omega, \mathcal{F}, \mathbb{P})$, a RV X is a function $X : \Omega \rightarrow \mathbb{R}$, such that for any $a \in \mathbb{R}$,

$$\{\omega \in \Omega : X(\omega) \leq a\} \in \mathcal{F}.$$

- A **random variable** (RV) is a function from a sample space Ω into the set of real numbers \mathbb{R} .
- Formally, given the probability space $(\Omega, \mathcal{F}, \mathbb{P})$, a RV X is a function $X : \Omega \rightarrow \mathbb{R}$, such that for any $a \in \mathbb{R}$,

$$\{\omega \in \Omega : X(\omega) \leq a\} \in \mathcal{F}.$$

- For a particular element $\omega \in \Omega$, $X(\omega)$ is called a *realization* of X .

- A **random variable** (RV) is a function from a sample space Ω into the set of real numbers \mathbb{R} .
- Formally, given the probability space $(\Omega, \mathcal{F}, \mathbb{P})$, a RV X is a function $X : \Omega \rightarrow \mathbb{R}$, such that for any $a \in \mathbb{R}$,

$$\{\omega \in \Omega : X(\omega) \leq a\} \in \mathcal{F}.$$

- For a particular element $\omega \in \Omega$, $X(\omega)$ is called a *realization* of X .
 - Usually, we will simply denote $X(\omega)$ as x when ω is not explicitly shown.

- A **random variable** (RV) is a function from a sample space Ω into the set of real numbers \mathbb{R} .
- Formally, given the probability space $(\Omega, \mathcal{F}, \mathbb{P})$, a RV X is a function $X : \Omega \rightarrow \mathbb{R}$, such that for any $a \in \mathbb{R}$,

$$\{\omega \in \Omega : X(\omega) \leq a\} \in \mathcal{F}.$$

- For a particular element $\omega \in \Omega$, $X(\omega)$ is called a *realization* of X .
 - Usually, we will simply denote $X(\omega)$ as x when ω is not explicitly shown.
 - A popular convention is to denote the RVs by upper-case letters (e.g., X and Y) and their realizations by lower-case letters (e.g., x and y).

- Example 1': Let $X(H) = 0$, $X(T) = 1$.



- Example 1': Let $X(H) = 0$, $X(T) = 1$.
- Example 2':
 - Under $(\Omega, \mathcal{F}, \mathbb{P})$, let $X(R) = 0$, $X(G) = 1$, and $X(B) = 2$.



- Example 1': Let $X(H) = 0$, $X(T) = 1$.
- Example 2':
 - Under $(\Omega, \mathcal{F}, \mathbb{P})$, let $X(R) = 0$, $X(G) = 1$, and $X(B) = 2$.
 - Under $(\Omega, \mathcal{F}_1, \mathbb{P})$, let $X(R) = 0$, $X(G) = 1$, and $X(B) = 1$.



- Example 1': Let $X(H) = 0$, $X(T) = 1$.
- Example 2':
 - Under $(\Omega, \mathcal{F}, \mathbb{P})$, let $X(R) = 0$, $X(G) = 1$, and $X(B) = 2$.
 - Under $(\Omega, \mathcal{F}_1, \mathbb{P})$, let $X(R) = 0$, $X(G) = 1$, and $X(B) = 1$.
- Example 3':
 - Under $(\Omega, \mathcal{F}_1, \mathbb{P})$, let $X(\omega) := \begin{cases} 0, & \text{if } \omega \in [0, a), \\ 1, & \text{if } \omega \in [a, 1]. \end{cases}$



- Example 1': Let $X(H) = 0$, $X(T) = 1$.
- Example 2':
 - Under $(\Omega, \mathcal{F}, \mathbb{P})$, let $X(R) = 0$, $X(G) = 1$, and $X(B) = 2$.
 - Under $(\Omega, \mathcal{F}_1, \mathbb{P})$, let $X(R) = 0$, $X(G) = 1$, and $X(B) = 1$.
- Example 3':
 - Under $(\Omega, \mathcal{F}_1, \mathbb{P})$, let $X(\omega) := \begin{cases} 0, & \text{if } \omega \in [0, a), \\ 1, & \text{if } \omega \in [a, 1]. \end{cases}$
 - Under $(\Omega, \mathcal{F}, \mathbb{P})$, let $X(\omega) = \omega$ for $\omega \in [0, 1]$.



- The **cumulative distribution function** (CDF) of a RV X , denoted by $F : \mathbb{R} \rightarrow [0, 1]$, is defined by

$$F(x) := \mathbb{P}(X \leq x) = \mathbb{P}(\{\omega \in \Omega : X(\omega) \leq x\}), \quad \forall x \in \mathbb{R},$$



- The **cumulative distribution function** (CDF) of a RV X , denoted by $F : \mathbb{R} \rightarrow [0, 1]$, is defined by

$$F(x) := \mathbb{P}(X \leq x) = \mathbb{P}(\{\omega \in \Omega : X(\omega) \leq x\}), \quad \forall x \in \mathbb{R},$$

and the following is satisfied:

- The **cumulative distribution function** (CDF) of a RV X , denoted by $F : \mathbb{R} \rightarrow [0, 1]$, is defined by

$$F(x) := \mathbb{P}(X \leq x) = \mathbb{P}(\{\omega \in \Omega : X(\omega) \leq x\}), \quad \forall x \in \mathbb{R},$$

and the following is satisfied:

- $\lim_{x \rightarrow -\infty} F(x) = 0$ and $\lim_{x \rightarrow +\infty} F(x) = 1$;



- The **cumulative distribution function** (CDF) of a RV X , denoted by $F : \mathbb{R} \rightarrow [0, 1]$, is defined by

$$F(x) := \mathbb{P}(X \leq x) = \mathbb{P}(\{\omega \in \Omega : X(\omega) \leq x\}), \quad \forall x \in \mathbb{R},$$

and the following is satisfied:

- $\lim_{x \rightarrow -\infty} F(x) = 0$ and $\lim_{x \rightarrow +\infty} F(x) = 1$;
- $F(x)$ is nondecreasing in x ;



- The **cumulative distribution function** (CDF) of a RV X , denoted by $F : \mathbb{R} \rightarrow [0, 1]$, is defined by

$$F(x) := \mathbb{P}(X \leq x) = \mathbb{P}(\{\omega \in \Omega : X(\omega) \leq x\}), \quad \forall x \in \mathbb{R},$$

and the following is satisfied:

- $\lim_{x \rightarrow -\infty} F(x) = 0$ and $\lim_{x \rightarrow +\infty} F(x) = 1$;
- $F(x)$ is nondecreasing in x ;
- $F(x)$ is right-continuous, that is, for any $x_0 \in \mathbb{R}$,

$$\lim_{x \downarrow x_0} F(x) = F(x_0).$$



- A RV X is said to be **discrete** if the set of its possible values is countable.



- A RV X is said to be **discrete** if the set of its possible values is countable.
- The **probability mass function** (pmf) of a discrete RV X is given by

$$p(x) := \mathbb{P}(X = x) = \mathbb{P}(\{\omega \in \Omega : X(\omega) = x\}), \quad \forall x \in \mathbb{R},$$



- A RV X is said to be **discrete** if the set of its possible values is countable.
- The **probability mass function** (pmf) of a discrete RV X is given by

$$p(x) := \mathbb{P}(X = x) = \mathbb{P}(\{\omega \in \Omega : X(\omega) = x\}), \quad \forall x \in \mathbb{R},$$

and the following is satisfied:

- $p(x) \geq 0$ for all $x \in \mathbb{R}$;
- $\sum_{x \in \mathbb{R}} p(x) = 1$.



- A RV X is said to be **discrete** if the set of its possible values is countable.
- The **probability mass function** (pmf) of a discrete RV X is given by

$$p(x) := \mathbb{P}(X = x) = \mathbb{P}(\{\omega \in \Omega : X(\omega) = x\}), \quad \forall x \in \mathbb{R},$$

and the following is satisfied:

- $p(x) \geq 0$ for all $x \in \mathbb{R}$;
 - $\sum_{x \in \mathbb{R}} p(x) = 1$.
- It is easy to see that $F(x) = \sum_{y \in (-\infty, x]} p(y)$.



- A RV X is said to be **continuous** if there exists a **probability density function** (pdf) $f(x)$ such that

$$F(x) = \mathbb{P}(X \leq x) = \int_{-\infty}^x f(t)dt, \quad \forall x \in \mathbb{R},$$



- A RV X is said to be **continuous** if there exists a **probability density function** (pdf) $f(x)$ such that

$$F(x) = \mathbb{P}(X \leq x) = \int_{-\infty}^x f(t)dt, \quad \forall x \in \mathbb{R},$$

and the following is satisfied:

- $f(x) \geq 0$ for all $x \in \mathbb{R}$;
- $\int_{-\infty}^{+\infty} f(t)dt = 1$.



- A RV X is said to be **continuous** if there exists a **probability density function** (pdf) $f(x)$ such that

$$F(x) = \mathbb{P}(X \leq x) = \int_{-\infty}^x f(t)dt, \quad \forall x \in \mathbb{R},$$

and the following is satisfied:

- $f(x) \geq 0$ for all $x \in \mathbb{R}$;
 - $\int_{-\infty}^{+\infty} f(t)dt = 1$.
- Observe that $\frac{d}{dx}F(x) = f(x)$.



- The **joint** CDF of RVs X and Y , denoted by $F : \mathbb{R} \times \mathbb{R} \rightarrow [0, 1]$, is defined by

$$\begin{aligned} F(x, y) &:= \mathbb{P}(X \leq x, Y \leq y) \\ &= \mathbb{P}(\{\omega : X(\omega) \leq x\} \cap \{\omega : Y(\omega) \leq y\}), \quad \forall x, y \in \mathbb{R}. \end{aligned}$$

- The **joint** CDF of RVs X and Y , denoted by $F : \mathbb{R} \times \mathbb{R} \rightarrow [0, 1]$, is defined by

$$\begin{aligned} F(x, y) &:= \mathbb{P}(X \leq x, Y \leq y) \\ &= \mathbb{P}(\{\omega : X(\omega) \leq x\} \cap \{\omega : Y(\omega) \leq y\}), \quad \forall x, y \in \mathbb{R}. \end{aligned}$$

- For discrete RVs X and Y , the **joint** pmf is given by

$$\begin{aligned} p(x, y) &:= \mathbb{P}(X = x, Y = y) \\ &= \mathbb{P}(\{\omega : X(\omega) = x\} \cap \{\omega : Y(\omega) = y\}), \quad \forall x, y \in \mathbb{R}. \end{aligned}$$

- The **joint** CDF of RVs X and Y , denoted by $F : \mathbb{R} \times \mathbb{R} \rightarrow [0, 1]$, is defined by

$$\begin{aligned} F(x, y) &:= \mathbb{P}(X \leq x, Y \leq y) \\ &= \mathbb{P}(\{\omega : X(\omega) \leq x\} \cap \{\omega : Y(\omega) \leq y\}), \quad \forall x, y \in \mathbb{R}. \end{aligned}$$

- For discrete RVs X and Y , the **joint** pmf is given by

$$\begin{aligned} p(x, y) &:= \mathbb{P}(X = x, Y = y) \\ &= \mathbb{P}(\{\omega : X(\omega) = x\} \cap \{\omega : Y(\omega) = y\}), \quad \forall x, y \in \mathbb{R}. \end{aligned}$$

- For continuous RVs X and Y , the **joint** pdf is $f(x, y)$ such that

$$F(x, y) = \int_{-\infty}^y \int_{-\infty}^x f(t, u) dt du, \quad \forall x, y \in \mathbb{R}.$$

- The **joint** CDF of RVs X and Y , denoted by $F : \mathbb{R} \times \mathbb{R} \rightarrow [0, 1]$, is defined by

$$\begin{aligned} F(x, y) &:= \mathbb{P}(X \leq x, Y \leq y) \\ &= \mathbb{P}(\{\omega : X(\omega) \leq x\} \cap \{\omega : Y(\omega) \leq y\}), \quad \forall x, y \in \mathbb{R}. \end{aligned}$$

- For discrete RVs X and Y , the **joint** pmf is given by

$$\begin{aligned} p(x, y) &:= \mathbb{P}(X = x, Y = y) \\ &= \mathbb{P}(\{\omega : X(\omega) = x\} \cap \{\omega : Y(\omega) = y\}), \quad \forall x, y \in \mathbb{R}. \end{aligned}$$

- For continuous RVs X and Y , the **joint** pdf is $f(x, y)$ such that

$$F(x, y) = \int_{-\infty}^y \int_{-\infty}^x f(t, u) dt du, \quad \forall x, y \in \mathbb{R}.$$

- Observe that $\frac{\partial^2 F(x, y)}{\partial x \partial y} = f(x, y)$.



- Given the random vector $(X, Y)^T$, the distribution of X or Y is called the **marginal distribution**.
 - The marginal CDF of X is $F_X(x) = F(x, +\infty)$.

- Given the random vector $(X, Y)^T$, the distribution of X or Y is called the **marginal distribution**.
 - The marginal CDF of X is $F_X(x) = F(x, +\infty)$.
- If $(X, Y)^T$ is discrete, the marginal pmf of X is

$$p_X(x) = \sum_{y \in \mathbb{R}} p(x, y).$$

- Given the random vector $(X, Y)^T$, the distribution of X or Y is called the **marginal distribution**.

- The marginal CDF of X is $F_X(x) = F(x, +\infty)$.

- If $(X, Y)^T$ is discrete, the marginal pmf of X is

$$p_X(x) = \sum_{y \in \mathbb{R}} p(x, y).$$

- If $(X, Y)^T$ is continuous, the marginal pdf of X is

$$f_X(x) = \int_{-\infty}^{+\infty} f(x, y) dy.$$

- Given the random vector $(X, Y)^T$, the distribution of X or Y is called the **marginal distribution**.

- The marginal CDF of X is $F_X(x) = F(x, +\infty)$.

- If $(X, Y)^T$ is discrete, the marginal pmf of X is

$$p_X(x) = \sum_{y \in \mathbb{R}} p(x, y).$$

- If $(X, Y)^T$ is continuous, the marginal pdf of X is

$$f_X(x) = \int_{-\infty}^{+\infty} f(x, y) dy.$$

- For Y , its marginal CDF, and pmf or pdf, can be determined similarly.

- If $(X, Y)^T$ is discrete, for any y such that $\mathbb{P}(Y = y) = p_Y(y) > 0$, the **conditional** pmf of X given that $Y = y$ is defined as

$$p(x|y) := \mathbb{P}(X = x|Y = y) = \frac{p(x, y)}{p_Y(y)}.$$

- If $(X, Y)^T$ is discrete, for any y such that $\mathbb{P}(Y = y) = p_Y(y) > 0$, the **conditional** pmf of X given that $Y = y$ is defined as

$$p(x|y) := \mathbb{P}(X = x|Y = y) = \frac{p(x, y)}{p_Y(y)}.$$

- If $(X, Y)^T$ is continuous, for any y such that $f_Y(y) > 0$, the **conditional** pdf of X given that $Y = y$ is defined as

$$f(x|y) := \frac{f(x, y)}{f_Y(y)}.$$

Intuitively, $f(x|y)$ can be understood as follows (although it is not the most rigorous approach):

Intuitively, $f(x|y)$ can be understood as follows (although it is not the most rigorous approach):

- 1 Note that

$$F(x|Y = y) = \lim_{\Delta \rightarrow 0} F(x|Y \text{ between } y \text{ and } y + \Delta)$$



Intuitively, $f(x|y)$ can be understood as follows (although it is not the most rigorous approach):

- 1 Note that

$$\begin{aligned} F(x|Y = y) &= \lim_{\Delta \rightarrow 0} F(x|Y \text{ between } y \text{ and } y + \Delta) \\ &= \lim_{\Delta \rightarrow 0} \frac{\mathbb{P}(X \leq x, Y \text{ between } y \text{ and } y + \Delta)}{\mathbb{P}(Y \text{ between } y \text{ and } y + \Delta)} \end{aligned}$$



Intuitively, $f(x|y)$ can be understood as follows (although it is not the most rigorous approach):

- 1 Note that

$$\begin{aligned} F(x|Y = y) &= \lim_{\Delta \rightarrow 0} F(x|Y \text{ between } y \text{ and } y + \Delta) \\ &= \lim_{\Delta \rightarrow 0} \frac{\mathbb{P}(X \leq x, Y \text{ between } y \text{ and } y + \Delta)}{\mathbb{P}(Y \text{ between } y \text{ and } y + \Delta)} \\ &= \frac{\lim_{\Delta \rightarrow 0} [F(x, y + \Delta) - F(x, y)] / \Delta}{\lim_{\Delta \rightarrow 0} [F_Y(y + \Delta) - F_Y(y)] / \Delta} \end{aligned}$$



Intuitively, $f(x|y)$ can be understood as follows (although it is not the most rigorous approach):

① Note that

$$\begin{aligned} F(x|Y = y) &= \lim_{\Delta \rightarrow 0} F(x|Y \text{ between } y \text{ and } y + \Delta) \\ &= \lim_{\Delta \rightarrow 0} \frac{\mathbb{P}(X \leq x, Y \text{ between } y \text{ and } y + \Delta)}{\mathbb{P}(Y \text{ between } y \text{ and } y + \Delta)} \\ &= \frac{\lim_{\Delta \rightarrow 0} [F(x, y + \Delta) - F(x, y)] / \Delta}{\lim_{\Delta \rightarrow 0} [F_Y(y + \Delta) - F_Y(y)] / \Delta} \\ &= \frac{\frac{\partial}{\partial y} F(x, y)}{\frac{d}{dy} F_Y(y)} \end{aligned}$$



Intuitively, $f(x|y)$ can be understood as follows (although it is not the most rigorous approach):

① Note that

$$\begin{aligned} F(x|Y=y) &= \lim_{\Delta \rightarrow 0} F(x|Y \text{ between } y \text{ and } y + \Delta) \\ &= \lim_{\Delta \rightarrow 0} \frac{\mathbb{P}(X \leq x, Y \text{ between } y \text{ and } y + \Delta)}{\mathbb{P}(Y \text{ between } y \text{ and } y + \Delta)} \\ &= \frac{\lim_{\Delta \rightarrow 0} [F(x, y + \Delta) - F(x, y)] / \Delta}{\lim_{\Delta \rightarrow 0} [F_Y(y + \Delta) - F_Y(y)] / \Delta} \\ &= \frac{\frac{\partial}{\partial y} F(x, y)}{\frac{d}{dy} F_Y(y)} = \frac{\frac{\partial}{\partial y} \int_{-\infty}^y \int_{-\infty}^x f(t, u) dt du}{f_Y(y)} \end{aligned}$$



Intuitively, $f(x|y)$ can be understood as follows (although it is not the most rigorous approach):

① Note that

$$\begin{aligned} F(x|Y=y) &= \lim_{\Delta \rightarrow 0} F(x|Y \text{ between } y \text{ and } y + \Delta) \\ &= \lim_{\Delta \rightarrow 0} \frac{\mathbb{P}(X \leq x, Y \text{ between } y \text{ and } y + \Delta)}{\mathbb{P}(Y \text{ between } y \text{ and } y + \Delta)} \\ &= \frac{\lim_{\Delta \rightarrow 0} [F(x, y + \Delta) - F(x, y)] / \Delta}{\lim_{\Delta \rightarrow 0} [F_Y(y + \Delta) - F_Y(y)] / \Delta} \\ &= \frac{\frac{\partial}{\partial y} F(x, y)}{\frac{d}{dy} F_Y(y)} = \frac{\frac{\partial}{\partial y} \int_{-\infty}^y \int_{-\infty}^x f(t, u) dt du}{f_Y(y)} \\ &= \frac{\int_{-\infty}^x f(t, y) dt}{f_Y(y)}. \end{aligned}$$



Intuitively, $f(x|y)$ can be understood as follows (although it is not the most rigorous approach):

① Note that

$$\begin{aligned} F(x|Y=y) &= \lim_{\Delta \rightarrow 0} F(x|Y \text{ between } y \text{ and } y + \Delta) \\ &= \lim_{\Delta \rightarrow 0} \frac{\mathbb{P}(X \leq x, Y \text{ between } y \text{ and } y + \Delta)}{\mathbb{P}(Y \text{ between } y \text{ and } y + \Delta)} \\ &= \frac{\lim_{\Delta \rightarrow 0} [F(x, y + \Delta) - F(x, y)] / \Delta}{\lim_{\Delta \rightarrow 0} [F_Y(y + \Delta) - F_Y(y)] / \Delta} \\ &= \frac{\frac{\partial}{\partial y} F(x, y)}{\frac{d}{dy} F_Y(y)} = \frac{\frac{\partial}{\partial y} \int_{-\infty}^y \int_{-\infty}^x f(t, u) dt du}{f_Y(y)} \\ &= \frac{\int_{-\infty}^x f(t, y) dt}{f_Y(y)}. \end{aligned}$$

② Then, $f(x|y) = \frac{\partial}{\partial x} F(x|Y=y) = \frac{\frac{\partial}{\partial x} \int_{-\infty}^x f(t, y) dt}{f_Y(y)} = \frac{f(x, y)}{f_Y(y)}$.

- Two RVs X and Y are said to be statistically **independent**, which can be denoted as $X \perp Y$, when, for any $x, y \in \mathbb{R}$,

$$F(x, y) = F_X(x)F_Y(y),$$



- Two RVs X and Y are said to be statistically **independent**, which can be denoted as $X \perp Y$, when, for any $x, y \in \mathbb{R}$,

$$F(x, y) = F_X(x)F_Y(y), \text{ or,}$$

$$p(x, y) = p_X(x)p_Y(y),$$

- Two RVs X and Y are said to be statistically **independent**, which can be denoted as $X \perp Y$, when, for any $x, y \in \mathbb{R}$,

$$F(x, y) = F_X(x)F_Y(y), \text{ or,}$$

$$p(x, y) = p_X(x)p_Y(y), \text{ or,}$$

$$f(x, y) = f_X(x)f_Y(y).$$



- Two RVs X and Y are said to be statistically **independent**, which can be denoted as $X \perp Y$, when, for any $x, y \in \mathbb{R}$,

$$F(x, y) = F_X(x)F_Y(y), \text{ or,}$$

$$p(x, y) = p_X(x)p_Y(y), \text{ or,}$$

$$f(x, y) = f_X(x)f_Y(y).$$

- X and Y are independent \iff
 - $p(x|y) \equiv p_X(x)$ or $f(x|y) \equiv f_X(x)$ regardless of the value y ;

- Two RVs X and Y are said to be statistically **independent**, which can be denoted as $X \perp Y$, when, for any $x, y \in \mathbb{R}$,

$$F(x, y) = F_X(x)F_Y(y), \text{ or,}$$

$$p(x, y) = p_X(x)p_Y(y), \text{ or,}$$

$$f(x, y) = f_X(x)f_Y(y).$$

- X and Y are independent \iff
 - $p(x|y) \equiv p_X(x)$ or $f(x|y) \equiv f_X(x)$ regardless of the value y ;
 - $\mathbb{P}(X \in A, Y \in B) = \mathbb{P}(X \in A)\mathbb{P}(Y \in B)$ for any $A, B \subset \mathbb{R}$.

- For more than two RVs X_1, \dots, X_n , the joint CDF, joint pmf or pdf, and the marginal pmf or pdf, are defined analogically.

- For more than two RVs X_1, \dots, X_n , the joint CDF, joint pmf or pdf, and the marginal pmf or pdf, are defined analogically.
- RVs X_1, \dots, X_n are (mutually) independent if

$$F(x_1, \dots, x_n) \equiv F_{X_1}(x_1) \times \cdots \times F_{X_n}(x_n), \text{ or,}$$

$$p(x_1, \dots, x_n) \equiv p_{X_1}(x_1) \times \cdots \times p_{X_n}(x_n), \text{ or,}$$

$$f(x_1, \dots, x_n) \equiv f_{X_1}(x_1) \times \cdots \times f_{X_n}(x_n).$$

- For more than two RVs X_1, \dots, X_n , the joint CDF, joint pmf or pdf, and the marginal pmf or pdf, are defined analogically.
- RVs X_1, \dots, X_n are (mutually) independent if

$$F(x_1, \dots, x_n) \equiv F_{X_1}(x_1) \times \cdots \times F_{X_n}(x_n), \text{ or,}$$

$$p(x_1, \dots, x_n) \equiv p_{X_1}(x_1) \times \cdots \times p_{X_n}(x_n), \text{ or,}$$

$$f(x_1, \dots, x_n) \equiv f_{X_1}(x_1) \times \cdots \times f_{X_n}(x_n).$$

- RVs X_1, \dots, X_n are pairwise independent if for any $i \neq j$, $X_i \perp X_j$.

- 1 Probability Space
- 2 Random Variables & Distributions
- 3 Expectations**
- 4 Common Distributions
- 5 Useful Inequalities
- 6 Convergence
- 7 Properties of a Random Sample

- The **expectation**, or **expected value**, or **mean**, of a RV X is defined as

$$\mathbb{E}[X] := \int_{\Omega} X(\omega) d\mathbb{P}(\omega),$$

provided that $\int_{\Omega} |X(\omega)| d\mathbb{P}(\omega) < \infty$ or $X \geq 0$ a.s., where the integral is the Lebesgue integral, rather than the Riemann integral.

- The **expectation**, or **expected value**, or **mean**, of a RV X is defined as

$$\mathbb{E}[X] := \int_{\Omega} X(\omega) d\mathbb{P}(\omega),$$

provided that $\int_{\Omega} |X(\omega)| d\mathbb{P}(\omega) < \infty$ or $X \geq 0$ a.s., where the integral is the Lebesgue integral, rather than the Riemann integral.

- For function $h : \mathbb{R} \rightarrow \mathbb{R}$, $\mathbb{E}[h(X)] = \int_{\Omega} h(X(\omega)) d\mathbb{P}(\omega)$.



- The **expectation**, or **expected value**, or **mean**, of a RV X is defined as

$$\mathbb{E}[X] := \int_{\Omega} X(\omega) d\mathbb{P}(\omega),$$

provided that $\int_{\Omega} |X(\omega)| d\mathbb{P}(\omega) < \infty$ or $X \geq 0$ a.s., where the integral is the Lebesgue integral, rather than the Riemann integral.

- For function $h : \mathbb{R} \rightarrow \mathbb{R}$, $\mathbb{E}[h(X)] = \int_{\Omega} h(X(\omega)) d\mathbb{P}(\omega)$.
- If X is a discrete RV:
 - $\mathbb{E}[X] = \sum_{x \in \mathbb{R}} xp(x)$;
 - $\mathbb{E}[h(X)] = \sum_{x \in \mathbb{R}} h(x)p(x)$.



- The **expectation**, or **expected value**, or **mean**, of a RV X is defined as

$$\mathbb{E}[X] := \int_{\Omega} X(\omega) d\mathbb{P}(\omega),$$

provided that $\int_{\Omega} |X(\omega)| d\mathbb{P}(\omega) < \infty$ or $X \geq 0$ a.s., where the integral is the Lebesgue integral, rather than the Riemann integral.

- For function $h : \mathbb{R} \rightarrow \mathbb{R}$, $\mathbb{E}[h(X)] = \int_{\Omega} h(X(\omega)) d\mathbb{P}(\omega)$.
- If X is a discrete RV:
 - $\mathbb{E}[X] = \sum_{x \in \mathbb{R}} xp(x)$;
 - $\mathbb{E}[h(X)] = \sum_{x \in \mathbb{R}} h(x)p(x)$.
- If X is a continuous RV:
 - $\mathbb{E}[X] = \int_{-\infty}^{+\infty} xf(x)dx$;
 - $\mathbb{E}[h(X)] = \int_{-\infty}^{+\infty} h(x)f(x)dx$.

- For integer n , $\mathbb{E}[X^n]$ is called the n th **moment** of X , and $\mathbb{E}[(X - \mathbb{E}[X])^n]$ is called the n th **central moment** of X .

Expectations

- For integer n , $\mathbb{E}[X^n]$ is called the n th **moment** of X , and $\mathbb{E}[(X - \mathbb{E}[X])^n]$ is called the n th **central moment** of X .
- Some special moments:
 - Mean (1st moment): $\mu := \mathbb{E}[X]$.

- For integer n , $\mathbb{E}[X^n]$ is called the n th **moment** of X , and $\mathbb{E}[(X - \mathbb{E}[X])^n]$ is called the n th **central moment** of X .
- Some special moments:
 - Mean (1st moment): $\mu := \mathbb{E}[X]$.
 - **Variance** (2nd central moment):
$$\sigma^2 := \text{Var}(X) := \mathbb{E}[(X - \mathbb{E}[X])^2] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2.$$

- For integer n , $\mathbb{E}[X^n]$ is called the n th **moment** of X , and $\mathbb{E}[(X - \mathbb{E}[X])^n]$ is called the n th **central moment** of X .
- Some special moments:
 - Mean (1st moment): $\mu := \mathbb{E}[X]$.
 - **Variance** (2nd central moment):
$$\sigma^2 := \text{Var}(X) := \mathbb{E}[(X - \mathbb{E}[X])^2] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2.$$
- **Linear** association:
 - **Covariance**:
$$\text{Cov}(X, Y) := \mathbb{E}[(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])] = \mathbb{E}[XY] - \mathbb{E}[X] \mathbb{E}[Y].$$

- For integer n , $\mathbb{E}[X^n]$ is called the n th **moment** of X , and $\mathbb{E}[(X - \mathbb{E}[X])^n]$ is called the n th **central moment** of X .
- Some special moments:
 - Mean (1st moment): $\mu := \mathbb{E}[X]$.
 - **Variance** (2nd central moment):
$$\sigma^2 := \text{Var}(X) := \mathbb{E}[(X - \mathbb{E}[X])^2] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2.$$
- **Linear** association:
 - **Covariance**:
$$\text{Cov}(X, Y) := \mathbb{E}[(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])] = \mathbb{E}[XY] - \mathbb{E}[X] \mathbb{E}[Y].$$
 - **Correlation**: $\rho(X, Y) := \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}.$

- For integer n , $\mathbb{E}[X^n]$ is called the n th **moment** of X , and $\mathbb{E}[(X - \mathbb{E}[X])^n]$ is called the n th **central moment** of X .
- Some special moments:
 - Mean (1st moment): $\mu := \mathbb{E}[X]$.
 - **Variance** (2nd central moment):
$$\sigma^2 := \text{Var}(X) := \mathbb{E}[(X - \mathbb{E}[X])^2] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2.$$
- **Linear** association:
 - **Covariance**:
$$\text{Cov}(X, Y) := \mathbb{E}[(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])] = \mathbb{E}[XY] - \mathbb{E}[X] \mathbb{E}[Y].$$
 - **Correlation**: $\rho(X, Y) := \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}.$
- In general, $X \perp Y \xLeftrightarrow{\neq} \rho(X, Y) = 0 \iff \text{Cov}(X, Y) = 0.$



- For integer n , $\mathbb{E}[X^n]$ is called the n th **moment** of X , and $\mathbb{E}[(X - \mathbb{E}[X])^n]$ is called the n th **central moment** of X .
- Some special moments:
 - Mean (1st moment): $\mu := \mathbb{E}[X]$.
 - **Variance** (2nd central moment):
 $\sigma^2 := \text{Var}(X) := \mathbb{E}[(X - \mathbb{E}[X])^2] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$.
- **Linear** association:
 - **Covariance**:
 $\text{Cov}(X, Y) := \mathbb{E}[(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])] = \mathbb{E}[XY] - \mathbb{E}[X] \mathbb{E}[Y]$.
 - **Correlation**: $\rho(X, Y) := \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}$.
- In general, $X \perp Y \xLeftrightarrow{\neq} \rho(X, Y) = 0 \iff \text{Cov}(X, Y) = 0$.
- If $(X, Y)^\top$ follows a bivariate normal distribution,[†] then $X \perp Y \iff \rho(X, Y) = 0$.

[†] **CAUTION:** It means MORE than that X and Y both follow a normal distribution! More details latter.

- The conditional expectation of X given $Y = y$ is

$$\mathbb{E}[X|y] := \begin{cases} \sum_{x \in \mathbb{R}} xp(x|y), & \text{if } X \text{ is discrete,} \\ \int_{-\infty}^{+\infty} xf(x|y)dx, & \text{if } X \text{ is continuous.} \end{cases}$$



- The conditional expectation of X given $Y = y$ is

$$\mathbb{E}[X|y] := \begin{cases} \sum_{x \in \mathbb{R}} xp(x|y), & \text{if } X \text{ is discrete,} \\ \int_{-\infty}^{+\infty} xf(x|y)dx, & \text{if } X \text{ is continuous.} \end{cases}$$

- The conditional variance of X given $Y = y$ is

$$\text{Var}(X|y) := \mathbb{E}[(X - \mathbb{E}[X])^2|y] = \mathbb{E}[X^2|y] - (\mathbb{E}[X|y])^2.$$

- The conditional expectation of X given $Y = y$ is

$$\mathbb{E}[X|y] := \begin{cases} \sum_{x \in \mathbb{R}} xp(x|y), & \text{if } X \text{ is discrete,} \\ \int_{-\infty}^{+\infty} xf(x|y)dx, & \text{if } X \text{ is continuous.} \end{cases}$$

- The conditional variance of X given $Y = y$ is

$$\text{Var}(X|y) := \mathbb{E}[(X - \mathbb{E}[X])^2|y] = \mathbb{E}[X^2|y] - (\mathbb{E}[X|y])^2.$$

- If $X \not\perp Y$, then $\mathbb{E}[X|y]$ and $\text{Var}(X|y)$ are functions of y .

- The conditional expectation of X given $Y = y$ is

$$\mathbb{E}[X|y] := \begin{cases} \sum_{x \in \mathbb{R}} xp(x|y), & \text{if } X \text{ is discrete,} \\ \int_{-\infty}^{+\infty} xf(x|y)dx, & \text{if } X \text{ is continuous.} \end{cases}$$

- The conditional variance of X given $Y = y$ is

$$\text{Var}(X|y) := \mathbb{E}[(X - \mathbb{E}[X])^2|y] = \mathbb{E}[X^2|y] - (\mathbb{E}[X|y])^2.$$

- If $X \not\perp Y$, then $\mathbb{E}[X|y]$ and $\text{Var}(X|y)$ are functions of y .
- If $X \not\perp Y$, then $\mathbb{E}[X|Y]$ and $\text{Var}(X|Y)$ are also RVs, whose value depends on the value of Y .

- The conditional expectation of X given $Y = y$ is

$$\mathbb{E}[X|y] := \begin{cases} \sum_{x \in \mathbb{R}} xp(x|y), & \text{if } X \text{ is discrete,} \\ \int_{-\infty}^{+\infty} xf(x|y)dx, & \text{if } X \text{ is continuous.} \end{cases}$$

- The conditional variance of X given $Y = y$ is

$$\text{Var}(X|y) := \mathbb{E}[(X - \mathbb{E}[X])^2|y] = \mathbb{E}[X^2|y] - (\mathbb{E}[X|y])^2.$$

- If $X \not\perp Y$, then $\mathbb{E}[X|y]$ and $\text{Var}(X|y)$ are functions of y .
- If $X \not\perp Y$, then $\mathbb{E}[X|Y]$ and $\text{Var}(X|Y)$ are also RVs, whose value depends on the value of Y .
- If $X \perp Y$, then $\mathbb{E}[X|y] = \mathbb{E}[X|Y] = \mathbb{E}[X]$, and $\text{Var}(X|y) = \text{Var}(X|Y) = \text{Var}(X)$.



- $\mathbb{E}[aX + bY] = a \mathbb{E}[X] + b \mathbb{E}[Y].$

- $\mathbb{E}[aX + bY] = a \mathbb{E}[X] + b \mathbb{E}[Y]$.
- $\text{Var}(aX + bY) = a^2 \text{Var}(X) + 2ab \text{Cov}(X, Y) + b^2 \text{Var}(Y)$.



- $\mathbb{E}[aX + bY] = a \mathbb{E}[X] + b \mathbb{E}[Y]$.
- $\text{Var}(aX + bY) = a^2 \text{Var}(X) + 2ab \text{Cov}(X, Y) + b^2 \text{Var}(Y)$.
- $\text{Cov}(aX + bY, cW + dV) = ac \text{Cov}(X, W) + ad \text{Cov}(X, V) + bc \text{Cov}(Y, W) + bd \text{Cov}(Y, V)$.



- $\mathbb{E}[aX + bY] = a \mathbb{E}[X] + b \mathbb{E}[Y]$.
- $\text{Var}(aX + bY) = a^2 \text{Var}(X) + 2ab \text{Cov}(X, Y) + b^2 \text{Var}(Y)$.
- $\text{Cov}(aX + bY, cW + dV) = ac \text{Cov}(X, W) + ad \text{Cov}(X, V) + bc \text{Cov}(Y, W) + bd \text{Cov}(Y, V)$.
- $\mathbb{E}[\mathbb{E}[X|Y]] = \mathbb{E}[X]$.



- $\mathbb{E}[aX + bY] = a \mathbb{E}[X] + b \mathbb{E}[Y].$
- $\text{Var}(aX + bY) = a^2 \text{Var}(X) + 2ab \text{Cov}(X, Y) + b^2 \text{Var}(Y).$
- $\text{Cov}(aX + bY, cW + dV) = ac \text{Cov}(X, W) +$
 $ad \text{Cov}(X, V) + bc \text{Cov}(Y, W) + bd \text{Cov}(Y, V).$
- $\mathbb{E}[\mathbb{E}[X|Y]] = \mathbb{E}[X].$
- $\text{Var}(X) = \mathbb{E}[\text{Var}(X|Y)] + \text{Var}(\mathbb{E}[X|Y]).$



- $\mathbb{E}[aX + bY] = a \mathbb{E}[X] + b \mathbb{E}[Y].$
- $\text{Var}(aX + bY) = a^2 \text{Var}(X) + 2ab \text{Cov}(X, Y) + b^2 \text{Var}(Y).$
- $\text{Cov}(aX + bY, cW + dV) = ac \text{Cov}(X, W) + ad \text{Cov}(X, V) + bc \text{Cov}(Y, W) + bd \text{Cov}(Y, V).$
- $\mathbb{E}[\mathbb{E}[X|Y]] = \mathbb{E}[X].$
- $\text{Var}(X) = \mathbb{E}[\text{Var}(X|Y)] + \text{Var}(\mathbb{E}[X|Y]).$
- If $X \perp Y$, then $\mathbb{E}[XY] = \mathbb{E}[X] \mathbb{E}[Y].$

- 1 Probability Space
- 2 Random Variables & Distributions
- 3 Expectations
- 4 Common Distributions**
- 5 Useful Inequalities
- 6 Convergence
- 7 Properties of a Random Sample

- $X \sim \text{Bernoulli}(p)$ or $\text{Ber}(p)$, if

$$X = \begin{cases} 1, & \text{with probability } p, \\ 0, & \text{with probability } 1 - p, \end{cases} \quad p \in [0, 1].$$



- $X \sim \text{Bernoulli}(p)$ or $\text{Ber}(p)$, if

$$X = \begin{cases} 1, & \text{with probability } p, \\ 0, & \text{with probability } 1 - p, \end{cases} \quad p \in [0, 1].$$

- $\mathbb{E}[X] = p$, $\text{Var}(X) = p(1 - p)$.



- $X \sim \text{Bernoulli}(p)$ or $\text{Ber}(p)$, if

$$X = \begin{cases} 1, & \text{with probability } p, \\ 0, & \text{with probability } 1 - p, \end{cases} \quad p \in [0, 1].$$

- $\mathbb{E}[X] = p$, $\text{Var}(X) = p(1 - p)$.
- The value $X = 1$ is often termed a “success” and p is referred to as the success probability.



- $X \sim \text{Bernoulli}(p)$ or $\text{Ber}(p)$, if

$$X = \begin{cases} 1, & \text{with probability } p, \\ 0, & \text{with probability } 1 - p, \end{cases} \quad p \in [0, 1].$$

- $\mathbb{E}[X] = p$, $\text{Var}(X) = p(1 - p)$.
- The value $X = 1$ is often termed a “success” and p is referred to as the success probability.
- $Y \sim \text{binomial}(n, p)$ or $\text{B}(n, p)$: The number of successes among n (mutually) **independent and identically distributed** (iid) $\text{Ber}(p)$ trials.

- $X \sim \text{Bernoulli}(p)$ or $\text{Ber}(p)$, if

$$X = \begin{cases} 1, & \text{with probability } p, \\ 0, & \text{with probability } 1 - p, \end{cases} \quad p \in [0, 1].$$

- $\mathbb{E}[X] = p$, $\text{Var}(X) = p(1 - p)$.
- The value $X = 1$ is often termed a “success” and p is referred to as the success probability.
- $Y \sim \text{binomial}(n, p)$ or $\text{B}(n, p)$: The number of successes among n (mutually) **independent and identically distributed** (iid) $\text{Ber}(p)$ trials.
 - $Y = \sum_{i=1}^n X_i$, where $X_i \sim \text{Ber}(p)$ are iid.



- $X \sim \text{Bernoulli}(p)$ or $\text{Ber}(p)$, if

$$X = \begin{cases} 1, & \text{with probability } p, \\ 0, & \text{with probability } 1 - p, \end{cases} \quad p \in [0, 1].$$

- $\mathbb{E}[X] = p$, $\text{Var}(X) = p(1 - p)$.
- The value $X = 1$ is often termed a “success” and p is referred to as the success probability.
- $Y \sim \text{binomial}(n, p)$ or $\text{B}(n, p)$: The number of successes among n (mutually) **independent and identically distributed** (iid) $\text{Ber}(p)$ trials.
 - $Y = \sum_{i=1}^n X_i$, where $X_i \sim \text{Ber}(p)$ are iid.
 - $p(y) = \mathbb{P}(Y = y) = \binom{n}{y} p^y (1 - p)^{n-y}$, $y = 0, 1, \dots, n$.



- $X \sim \text{Bernoulli}(p)$ or $\text{Ber}(p)$, if

$$X = \begin{cases} 1, & \text{with probability } p, \\ 0, & \text{with probability } 1 - p, \end{cases} \quad p \in [0, 1].$$

- $\mathbb{E}[X] = p$, $\text{Var}(X) = p(1 - p)$.
- The value $X = 1$ is often termed a “success” and p is referred to as the success probability.
- $Y \sim \text{binomial}(n, p)$ or $\text{B}(n, p)$: The number of successes among n (mutually) **independent and identically distributed** (iid) $\text{Ber}(p)$ trials.
 - $Y = \sum_{i=1}^n X_i$, where $X_i \sim \text{Ber}(p)$ are iid.
 - $p(y) = \mathbb{P}(Y = y) = \binom{n}{y} p^y (1 - p)^{n-y}$, $y = 0, 1, \dots, n$.
 - $\mathbb{E}[Y] = np$, $\text{Var}(Y) = np(1 - p)$.



- $X \sim \text{Bernoulli}(p)$ or $\text{Ber}(p)$, if

$$X = \begin{cases} 1, & \text{with probability } p, \\ 0, & \text{with probability } 1 - p, \end{cases} \quad p \in [0, 1].$$

- $\mathbb{E}[X] = p$, $\text{Var}(X) = p(1 - p)$.
- The value $X = 1$ is often termed a “success” and p is referred to as the success probability.
- $Y \sim \text{binomial}(n, p)$ or $\text{B}(n, p)$: The number of successes among n (mutually) **independent and identically distributed** (iid) $\text{Ber}(p)$ trials.
 - $Y = \sum_{i=1}^n X_i$, where $X_i \sim \text{Ber}(p)$ are iid.
 - $p(y) = \mathbb{P}(Y = y) = \binom{n}{y} p^y (1 - p)^{n-y}$, $y = 0, 1, \dots, n$.
 - $\mathbb{E}[Y] = np$, $\text{Var}(Y) = np(1 - p)$.
- If $Y_1 \sim \text{B}(n_1, p)$ and $Y_2 \sim \text{B}(n_2, p)$ are independent, then $Y_1 + Y_2 \sim \text{B}(n_1 + n_2, p)$.



- $Y \sim \text{negative binomial}(r, p)$ or $\text{NB}(r, p)$: The number of iid $\text{Ber}(p)$ trials to obtain r successes.



- $Y \sim \text{negative binomial}(r, p)$ or $\text{NB}(r, p)$: The number of iid $\text{Ber}(p)$ trials to obtain r successes.
 - $p(y) = \mathbb{P}(Y = y) = \binom{y-1}{r-1} p^r (1-p)^{y-r}, \quad y = r, r+1, \dots$



- $Y \sim \text{negative binomial}(r, p)$ or $\text{NB}(r, p)$: The number of iid $\text{Ber}(p)$ trials to obtain r successes.
 - $p(y) = \mathbb{P}(Y = y) = \binom{y-1}{r-1} p^r (1-p)^{y-r}, \quad y = r, r+1, \dots$
 - $\mathbb{E}[Y] = r + r(1-p)/p, \text{Var}(Y) = r(1-p)/p^2$.



- $Y \sim \text{negative binomial}(r, p)$ or $\text{NB}(r, p)$: The number of iid $\text{Ber}(p)$ trials to obtain r successes.
 - $p(y) = \mathbb{P}(Y = y) = \binom{y-1}{r-1} p^r (1-p)^{y-r}, \quad y = r, r+1, \dots$
 - $\mathbb{E}[Y] = r + r(1-p)/p, \text{Var}(Y) = r(1-p)/p^2$.
 - When $r = 1$, it becomes the geometric distribution.



- $Y \sim \text{negative binomial}(r, p)$ or $\text{NB}(r, p)$: The number of iid $\text{Ber}(p)$ trials to obtain r successes.
 - $p(y) = \mathbb{P}(Y = y) = \binom{y-1}{r-1} p^r (1-p)^{y-r}, \quad y = r, r+1, \dots$
 - $\mathbb{E}[Y] = r + r(1-p)/p, \text{Var}(Y) = r(1-p)/p^2$.
 - When $r = 1$, it becomes the geometric distribution.
- $Y \sim \text{geometric}(p)$ or $\text{Geo}(p)$: The number of iid $\text{Ber}(p)$ trials to obtain the first success.



- $Y \sim \text{negative binomial}(r, p)$ or $\text{NB}(r, p)$: The number of iid $\text{Ber}(p)$ trials to obtain r successes.
 - $p(y) = \mathbb{P}(Y = y) = \binom{y-1}{r-1} p^r (1-p)^{y-r}, \quad y = r, r+1, \dots$
 - $\mathbb{E}[Y] = r + r(1-p)/p, \text{Var}(Y) = r(1-p)/p^2$.
 - When $r = 1$, it becomes the geometric distribution.
- $Y \sim \text{geometric}(p)$ or $\text{Geo}(p)$: The number of iid $\text{Ber}(p)$ trials to obtain the first success.
 - $p(y) = \mathbb{P}(Y = y) = p(1-p)^{y-1}, \quad y = 1, 2, \dots$
 - $\mathbb{E}[Y] = 1/p, \text{Var}(Y) = (1-p)/p^2$.



- $Y \sim \text{negative binomial}(r, p)$ or $\text{NB}(r, p)$: The number of iid $\text{Ber}(p)$ trials to obtain r successes.
 - $p(y) = \mathbb{P}(Y = y) = \binom{y-1}{r-1} p^r (1-p)^{y-r}, \quad y = r, r+1, \dots$
 - $\mathbb{E}[Y] = r + r(1-p)/p, \text{Var}(Y) = r(1-p)/p^2$.
 - When $r = 1$, it becomes the geometric distribution.
- $Y \sim \text{geometric}(p)$ or $\text{Geo}(p)$: The number of iid $\text{Ber}(p)$ trials to obtain the first success.
 - $p(y) = \mathbb{P}(Y = y) = p(1-p)^{y-1}, \quad y = 1, 2, \dots$
 - $\mathbb{E}[Y] = 1/p, \text{Var}(Y) = (1-p)/p^2$.
 - **Memoryless Property**: For integers $s > t$,

$$\begin{aligned}\mathbb{P}(Y > s | Y > t) &= \frac{\mathbb{P}(Y > s, Y > t)}{\mathbb{P}(Y > t)} = \frac{\mathbb{P}(Y > s)}{\mathbb{P}(Y > t)} = \frac{(1-p)^s}{(1-p)^t} = (1-p)^{s-t} \\ &= \mathbb{P}(Y > s-t).\end{aligned}$$



- $Y \sim \text{negative binomial}(r, p)$ or $\text{NB}(r, p)$: The number of iid $\text{Ber}(p)$ trials to obtain r successes.
 - $p(y) = \mathbb{P}(Y = y) = \binom{y-1}{r-1} p^r (1-p)^{y-r}, \quad y = r, r+1, \dots$
 - $\mathbb{E}[Y] = r + r(1-p)/p, \text{Var}(Y) = r(1-p)/p^2$.
 - When $r = 1$, it becomes the geometric distribution.
- $Y \sim \text{geometric}(p)$ or $\text{Geo}(p)$: The number of iid $\text{Ber}(p)$ trials to obtain the first success.
 - $p(y) = \mathbb{P}(Y = y) = p(1-p)^{y-1}, \quad y = 1, 2, \dots$
 - $\mathbb{E}[Y] = 1/p, \text{Var}(Y) = (1-p)/p^2$.
 - **Memoryless Property**: For integers $s > t$,

$$\begin{aligned}\mathbb{P}(Y > s | Y > t) &= \frac{\mathbb{P}(Y > s, Y > t)}{\mathbb{P}(Y > t)} = \frac{\mathbb{P}(Y > s)}{\mathbb{P}(Y > t)} = \frac{(1-p)^s}{(1-p)^t} = (1-p)^{s-t} \\ &= \mathbb{P}(Y > s-t).\end{aligned}$$

- If $Y_1 \sim \text{NB}(r_1, p)$ and $Y_2 \sim \text{NB}(r_2, p)$ are independent, then $Y_1 + Y_2 \sim \text{NB}(r_1 + r_2, p)$.

- Poisson distribution is often used to model the number of occurrence in a given time interval.

- Poisson distribution is often used to model the number of occurrence in a given time interval.
- One of the basic assumptions is that, *for very small time intervals, the probability of an occurrence is proportional to the length of the time interval.*[†]

[†] See more detailed discussion in Lec 3.

- Poisson distribution is often used to model the number of occurrence in a given time interval.
- One of the basic assumptions is that, *for very small time intervals, the probability of an occurrence is proportional to the length of the time interval.*[†]
- $X \sim \text{Poisson}(\lambda)$ or $\text{Pois}(\lambda)$, with $\lambda > 0$, if

$$p(x) = \mathbb{P}(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}, \quad x = 0, 1, \dots$$

[†] See more detailed discussion in Lec 3.

- Poisson distribution is often used to model the number of occurrence in a given time interval.
- One of the basic assumptions is that, *for very small time intervals, the probability of an occurrence is proportional to the length of the time interval.*[†]
- $X \sim \text{Poisson}(\lambda)$ or $\text{Pois}(\lambda)$, with $\lambda > 0$, if

$$p(x) = \mathbb{P}(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}, \quad x = 0, 1, \dots$$

- It can be verified that $\sum_{x=0}^{\infty} p(x) = 1$.

[†] See more detailed discussion in Lec 3.

- Poisson distribution is often used to model the number of occurrence in a given time interval.
- One of the basic assumptions is that, *for very small time intervals, the probability of an occurrence is proportional to the length of the time interval.*[†]
- $X \sim \text{Poisson}(\lambda)$ or $\text{Pois}(\lambda)$, with $\lambda > 0$, if

$$p(x) = \mathbb{P}(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}, \quad x = 0, 1, \dots$$

- It can be verified that $\sum_{x=0}^{\infty} p(x) = 1$.
- $\mathbb{E}[X] = \lambda$, $\text{Var}(X) = \lambda$.

[†] See more detailed discussion in Lec 3.

- Poisson distribution is often used to model the number of occurrence in a given time interval.
- One of the basic assumptions is that, *for very small time intervals, the probability of an occurrence is proportional to the length of the time interval.*[†]
- $X \sim \text{Poisson}(\lambda)$ or $\text{Pois}(\lambda)$, with $\lambda > 0$, if

$$p(x) = \mathbb{P}(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}, \quad x = 0, 1, \dots$$

- It can be verified that $\sum_{x=0}^{\infty} p(x) = 1$.
- $\mathbb{E}[X] = \lambda$, $\text{Var}(X) = \lambda$.
- If $X_1 \sim \text{Pois}(\lambda_1)$ and $X_2 \sim \text{Pois}(\lambda_2)$ are independent,
 - $X_1 + X_2 \sim \text{Pois}(\lambda_1 + \lambda_2)$;
 - Given $X_1 + X_2 = n$, $X_1 \sim \text{B}(n, \lambda_1/(\lambda_1 + \lambda_2))$.

[†] See more detailed discussion in Lec 3.



- $X \sim \text{Uniform}(a, b)$ or $\text{Unif}(a, b)$ with $a < b$, if its pdf is given by

$$f(x) = \begin{cases} \frac{1}{b-a}, & \text{if } x \in [a, b], \\ 0, & \text{otherwise.} \end{cases}$$

- $X \sim \text{Uniform}(a, b)$ or $\text{Unif}(a, b)$ with $a < b$, if its pdf is given by

$$f(x) = \begin{cases} \frac{1}{b-a}, & \text{if } x \in [a, b], \\ 0, & \text{otherwise.} \end{cases}$$

- $\mathbb{E}[X] = \frac{b+a}{2}$, $\text{Var}(X) = \frac{(b-a)^2}{12}$.

- $X \sim \text{Uniform}(a, b)$ or $\text{Unif}(a, b)$ with $a < b$, if its pdf is given by

$$f(x) = \begin{cases} \frac{1}{b-a}, & \text{if } x \in [a, b], \\ 0, & \text{otherwise.} \end{cases}$$

- $\mathbb{E}[X] = \frac{b+a}{2}$, $\text{Var}(X) = \frac{(b-a)^2}{12}$.
- $X \sim \text{exponential}(\lambda)$ or $\text{Exp}(\lambda)$, with $\lambda > 0$, if its pdf is given by

$$f(x) = \lambda e^{-\lambda x}, \quad x \in [0, \infty).$$

- $X \sim \text{Uniform}(a, b)$ or $\text{Unif}(a, b)$ with $a < b$, if its pdf is given by

$$f(x) = \begin{cases} \frac{1}{b-a}, & \text{if } x \in [a, b], \\ 0, & \text{otherwise.} \end{cases}$$

- $\mathbb{E}[X] = \frac{b+a}{2}$, $\text{Var}(X) = \frac{(b-a)^2}{12}$.
- $X \sim \text{exponential}(\lambda)$ or $\text{Exp}(\lambda)$, with $\lambda > 0$, if its pdf is given by

$$f(x) = \lambda e^{-\lambda x}, \quad x \in [0, \infty).$$

- λ is called the rate parameter.

- $X \sim \text{Uniform}(a, b)$ or $\text{Unif}(a, b)$ with $a < b$, if its pdf is given by

$$f(x) = \begin{cases} \frac{1}{b-a}, & \text{if } x \in [a, b], \\ 0, & \text{otherwise.} \end{cases}$$

- $\mathbb{E}[X] = \frac{b+a}{2}$, $\text{Var}(X) = \frac{(b-a)^2}{12}$.
- $X \sim \text{exponential}(\lambda)$ or $\text{Exp}(\lambda)$, with $\lambda > 0$, if its pdf is given by

$$f(x) = \lambda e^{-\lambda x}, \quad x \in [0, \infty).$$

- λ is called the rate parameter.
- $F(x) = 1 - e^{-\lambda x}$, $\mathbb{P}(X > x) = 1 - F(x) = e^{-\lambda x}$.

- $X \sim \text{Uniform}(a, b)$ or $\text{Unif}(a, b)$ with $a < b$, if its pdf is given by

$$f(x) = \begin{cases} \frac{1}{b-a}, & \text{if } x \in [a, b], \\ 0, & \text{otherwise.} \end{cases}$$

- $\mathbb{E}[X] = \frac{b+a}{2}$, $\text{Var}(X) = \frac{(b-a)^2}{12}$.
- $X \sim \text{exponential}(\lambda)$ or $\text{Exp}(\lambda)$, with $\lambda > 0$, if its pdf is given by

$$f(x) = \lambda e^{-\lambda x}, \quad x \in [0, \infty).$$

- λ is called the rate parameter.
- $F(x) = 1 - e^{-\lambda x}$, $\mathbb{P}(X > x) = 1 - F(x) = e^{-\lambda x}$.
- $\mathbb{E}[X] = 1/\lambda$, $\text{Var}(X) = 1/\lambda^2$.

- $X \sim \text{Uniform}(a, b)$ or $\text{Unif}(a, b)$ with $a < b$, if its pdf is given by

$$f(x) = \begin{cases} \frac{1}{b-a}, & \text{if } x \in [a, b], \\ 0, & \text{otherwise.} \end{cases}$$

- $\mathbb{E}[X] = \frac{b+a}{2}$, $\text{Var}(X) = \frac{(b-a)^2}{12}$.
- $X \sim \text{exponential}(\lambda)$ or $\text{Exp}(\lambda)$, with $\lambda > 0$, if its pdf is given by

$$f(x) = \lambda e^{-\lambda x}, \quad x \in [0, \infty).$$

- λ is called the rate parameter.
- $F(x) = 1 - e^{-\lambda x}$, $\mathbb{P}(X > x) = 1 - F(x) = e^{-\lambda x}$.
- $\mathbb{E}[X] = 1/\lambda$, $\text{Var}(X) = 1/\lambda^2$.
- **Memoryless Property:** For $s > t \geq 0$,

$$\begin{aligned} \mathbb{P}(X > s | X > t) &= \frac{\mathbb{P}(X > s, X > t)}{\mathbb{P}(X > t)} = \frac{\mathbb{P}(X > s)}{\mathbb{P}(X > t)} = \frac{e^{-\lambda s}}{e^{-\lambda t}} = e^{-\lambda(s-t)} \\ &= \mathbb{P}(X > s - t). \end{aligned}$$

- If $X_1 \sim \text{Exp}(\lambda_1)$ and $X_2 \sim \text{Exp}(\lambda_2)$ are independent, then $\min\{X_1, X_2\} \sim \text{Exp}(\lambda_1 + \lambda_2)$.

- If $X_1 \sim \text{Exp}(\lambda_1)$ and $X_2 \sim \text{Exp}(\lambda_2)$ are independent, then $\min\{X_1, X_2\} \sim \text{Exp}(\lambda_1 + \lambda_2)$.
- If $X \sim \text{Exp}(\lambda)$, then for $\alpha > 0$, $Y := X^{1/\alpha} \sim \text{Weibull}(\alpha, \beta)$ in shape & scale parametrization with $\beta = (1/\lambda)^{1/\alpha}$, whose pdf is

$$f(y) = \alpha \beta^{-\alpha} y^{\alpha-1} e^{-(y/\beta)^\alpha}, \quad y \in (0, \infty).$$

- If $X_1 \sim \text{Exp}(\lambda_1)$ and $X_2 \sim \text{Exp}(\lambda_2)$ are independent, then $\min\{X_1, X_2\} \sim \text{Exp}(\lambda_1 + \lambda_2)$.
- If $X \sim \text{Exp}(\lambda)$, then for $\alpha > 0$, $Y := X^{1/\alpha} \sim \text{Weibull}(\alpha, \beta)$ in shape & scale parametrization with $\beta = (1/\lambda)^{1/\alpha}$, whose pdf is

$$f(y) = \alpha \beta^{-\alpha} y^{\alpha-1} e^{-(y/\beta)^\alpha}, \quad y \in (0, \infty).$$

- Erlang(k, λ) or $\text{Erl}(k, \lambda)$, with k being a positive integer, is a generalized version of $\text{Exp}(\lambda)$, whose pdf is

$$f(x) = \frac{\lambda^k}{(k-1)!} x^{k-1} e^{-\lambda x}, \quad x \in [0, \infty).$$



- If $X_1 \sim \text{Exp}(\lambda_1)$ and $X_2 \sim \text{Exp}(\lambda_2)$ are independent, then $\min\{X_1, X_2\} \sim \text{Exp}(\lambda_1 + \lambda_2)$.
- If $X \sim \text{Exp}(\lambda)$, then for $\alpha > 0$, $Y := X^{1/\alpha} \sim \text{Weibull}(\alpha, \beta)$ in shape & scale parametrization with $\beta = (1/\lambda)^{1/\alpha}$, whose pdf is

$$f(y) = \alpha \beta^{-\alpha} y^{\alpha-1} e^{-(y/\beta)^\alpha}, \quad y \in (0, \infty).$$

- Erlang(k, λ) or $\text{Erl}(k, \lambda)$, with k being a positive integer, is a generalized version of $\text{Exp}(\lambda)$, whose pdf is

$$f(x) = \frac{\lambda^k}{(k-1)!} x^{k-1} e^{-\lambda x}, \quad x \in [0, \infty).$$

- $\mathbb{E}[X] = k/\lambda$, $\text{Var}(X) = k/\lambda^2$.



- If $X_1 \sim \text{Exp}(\lambda_1)$ and $X_2 \sim \text{Exp}(\lambda_2)$ are independent, then $\min\{X_1, X_2\} \sim \text{Exp}(\lambda_1 + \lambda_2)$.
- If $X \sim \text{Exp}(\lambda)$, then for $\alpha > 0$, $Y := X^{1/\alpha} \sim \text{Weibull}(\alpha, \beta)$ in shape & scale parametrization with $\beta = (1/\lambda)^{1/\alpha}$, whose pdf is

$$f(y) = \alpha \beta^{-\alpha} y^{\alpha-1} e^{-(y/\beta)^\alpha}, \quad y \in (0, \infty).$$

- Erlang(k, λ) or $\text{Erl}(k, \lambda)$, with k being a positive integer, is a generalized version of $\text{Exp}(\lambda)$, whose pdf is

$$f(x) = \frac{\lambda^k}{(k-1)!} x^{k-1} e^{-\lambda x}, \quad x \in [0, \infty).$$

- $\mathbb{E}[X] = k/\lambda$, $\text{Var}(X) = k/\lambda^2$.
- $k = 1 \implies \text{Exp}(\lambda)$.



- If $X_1 \sim \text{Exp}(\lambda_1)$ and $X_2 \sim \text{Exp}(\lambda_2)$ are independent, then $\min\{X_1, X_2\} \sim \text{Exp}(\lambda_1 + \lambda_2)$.
- If $X \sim \text{Exp}(\lambda)$, then for $\alpha > 0$, $Y := X^{1/\alpha} \sim \text{Weibull}(\alpha, \beta)$ in shape & scale parametrization with $\beta = (1/\lambda)^{1/\alpha}$, whose pdf is

$$f(y) = \alpha \beta^{-\alpha} y^{\alpha-1} e^{-(y/\beta)^\alpha}, \quad y \in (0, \infty).$$

- Erlang(k, λ) or $\text{Erl}(k, \lambda)$, with k being a positive integer, is a generalized version of $\text{Exp}(\lambda)$, whose pdf is

$$f(x) = \frac{\lambda^k}{(k-1)!} x^{k-1} e^{-\lambda x}, \quad x \in [0, \infty).$$

- $\mathbb{E}[X] = k/\lambda$, $\text{Var}(X) = k/\lambda^2$.
 - $k = 1 \implies \text{Exp}(\lambda)$.
- If $X_1 \sim \text{Erl}(k_1, \lambda)$ and $X_2 \sim \text{Erl}(k_2, \lambda)$ are independent, then $X_1 + X_2 \sim \text{Erl}(k_1 + k_2, \lambda)$.

- If $X_1 \sim \text{Exp}(\lambda_1)$ and $X_2 \sim \text{Exp}(\lambda_2)$ are independent, then $\min\{X_1, X_2\} \sim \text{Exp}(\lambda_1 + \lambda_2)$.
- If $X \sim \text{Exp}(\lambda)$, then for $\alpha > 0$, $Y := X^{1/\alpha} \sim \text{Weibull}(\alpha, \beta)$ in shape & scale parametrization with $\beta = (1/\lambda)^{1/\alpha}$, whose pdf is

$$f(y) = \alpha \beta^{-\alpha} y^{\alpha-1} e^{-(y/\beta)^\alpha}, \quad y \in (0, \infty).$$

- Erlang(k, λ) or $\text{Erl}(k, \lambda)$, with k being a positive integer, is a generalized version of $\text{Exp}(\lambda)$, whose pdf is

$$f(x) = \frac{\lambda^k}{(k-1)!} x^{k-1} e^{-\lambda x}, \quad x \in [0, \infty).$$

- $\mathbb{E}[X] = k/\lambda$, $\text{Var}(X) = k/\lambda^2$.
 - $k = 1 \implies \text{Exp}(\lambda)$.
- If $X_1 \sim \text{Erl}(k_1, \lambda)$ and $X_2 \sim \text{Erl}(k_2, \lambda)$ are independent, then $X_1 + X_2 \sim \text{Erl}(k_1 + k_2, \lambda)$.
- If $X \sim \text{Erl}(k, \lambda)$, then $cX \sim \text{Erl}(k, \lambda/c)$ for $c > 0$.



- $X \sim \text{Gamma}(\alpha, \lambda)$ in shape & rate parametrization with $\alpha, \lambda > 0$, if its pdf is given by

$$f(x) = \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x}, \quad x \in [0, \infty).$$

- $X \sim \text{Gamma}(\alpha, \lambda)$ in shape & rate parametrization with $\alpha, \lambda > 0$, if its pdf is given by

$$f(x) = \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x}, \quad x \in [0, \infty).$$

- $\mathbb{E}[X] = \alpha/\lambda$, $\text{Var}(X) = \alpha/\lambda^2$.

- $X \sim \text{Gamma}(\alpha, \lambda)$ in shape & rate parametrization with $\alpha, \lambda > 0$, if its pdf is given by

$$f(x) = \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x}, \quad x \in [0, \infty).$$

- $\mathbb{E}[X] = \alpha/\lambda$, $\text{Var}(X) = \alpha/\lambda^2$.
- $\Gamma(\alpha) := \int_0^\infty t^{\alpha-1} e^{-t} dt$ is known as the gamma function.
 - $\Gamma(\alpha + 1) = \alpha\Gamma(\alpha)$; $\Gamma(n) = (n-1)!$, for integer $n > 0$.

- $X \sim \text{Gamma}(\alpha, \lambda)$ in shape & rate parametrization with $\alpha, \lambda > 0$, if its pdf is given by

$$f(x) = \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x}, \quad x \in [0, \infty).$$

- $\mathbb{E}[X] = \alpha/\lambda$, $\text{Var}(X) = \alpha/\lambda^2$.
- $\Gamma(\alpha) := \int_0^\infty t^{\alpha-1} e^{-t} dt$ is known as the gamma function.
 - $\Gamma(\alpha + 1) = \alpha\Gamma(\alpha)$; $\Gamma(n) = (n-1)!$, for integer $n > 0$.
- If $X_1 \sim \text{Gamma}(\alpha_1, \lambda)$ and $X_2 \sim \text{Gamma}(\alpha_2, \lambda)$ are independent, then $X_1 + X_2 \sim \text{Gamma}(\alpha_1 + \alpha_2, \lambda)$.

- $X \sim \text{Gamma}(\alpha, \lambda)$ in shape & rate parametrization with $\alpha, \lambda > 0$, if its pdf is given by

$$f(x) = \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x}, \quad x \in [0, \infty).$$

- $\mathbb{E}[X] = \alpha/\lambda$, $\text{Var}(X) = \alpha/\lambda^2$.
- $\Gamma(\alpha) := \int_0^\infty t^{\alpha-1} e^{-t} dt$ is known as the gamma function.
 - $\Gamma(\alpha + 1) = \alpha\Gamma(\alpha)$; $\Gamma(n) = (n-1)!$, for integer $n > 0$.
- If $X_1 \sim \text{Gamma}(\alpha_1, \lambda)$ and $X_2 \sim \text{Gamma}(\alpha_2, \lambda)$ are independent, then $X_1 + X_2 \sim \text{Gamma}(\alpha_1 + \alpha_2, \lambda)$.
- If $X \sim \text{Gamma}(\alpha, \lambda)$, then $cX \sim \text{Gamma}(\alpha, \lambda/c)$ for $c > 0$.

- $X \sim \text{Gamma}(\alpha, \lambda)$ in shape & rate parametrization with $\alpha, \lambda > 0$, if its pdf is given by

$$f(x) = \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x}, \quad x \in [0, \infty).$$

- $\mathbb{E}[X] = \alpha/\lambda$, $\text{Var}(X) = \alpha/\lambda^2$.
- $\Gamma(\alpha) := \int_0^\infty t^{\alpha-1} e^{-t} dt$ is known as the gamma function.
 - $\Gamma(\alpha + 1) = \alpha\Gamma(\alpha)$; $\Gamma(n) = (n-1)!$, for integer $n > 0$.
- If $X_1 \sim \text{Gamma}(\alpha_1, \lambda)$ and $X_2 \sim \text{Gamma}(\alpha_2, \lambda)$ are independent, then $X_1 + X_2 \sim \text{Gamma}(\alpha_1 + \alpha_2, \lambda)$.
- If $X \sim \text{Gamma}(\alpha, \lambda)$, then $cX \sim \text{Gamma}(\alpha, \lambda/c)$ for $c > 0$.
- Important special cases of $\text{Gamma}(\alpha, \lambda)$:
 - α is an integer $\implies \text{Erl}(\alpha, \lambda)$; $\alpha = 1 \implies \text{Exp}(\lambda)$;



- $X \sim \text{Gamma}(\alpha, \lambda)$ in shape & rate parametrization with $\alpha, \lambda > 0$, if its pdf is given by

$$f(x) = \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x}, \quad x \in [0, \infty).$$

- $\mathbb{E}[X] = \alpha/\lambda$, $\text{Var}(X) = \alpha/\lambda^2$.
- $\Gamma(\alpha) := \int_0^\infty t^{\alpha-1} e^{-t} dt$ is known as the gamma function.
 - $\Gamma(\alpha + 1) = \alpha\Gamma(\alpha)$; $\Gamma(n) = (n-1)!$, for integer $n > 0$.
- If $X_1 \sim \text{Gamma}(\alpha_1, \lambda)$ and $X_2 \sim \text{Gamma}(\alpha_2, \lambda)$ are independent, then $X_1 + X_2 \sim \text{Gamma}(\alpha_1 + \alpha_2, \lambda)$.
- If $X \sim \text{Gamma}(\alpha, \lambda)$, then $cX \sim \text{Gamma}(\alpha, \lambda/c)$ for $c > 0$.
- Important special cases of $\text{Gamma}(\alpha, \lambda)$:
 - α is an integer $\implies \text{Erl}(\alpha, \lambda)$; $\alpha = 1 \implies \text{Exp}(\lambda)$;
 - $\alpha = p/2$, where p is an integer, and $\lambda = 1/2 \implies$ **chi-square distribution with p degrees of freedom**, denoted as χ_p^2 .

- Beta distribution is a very flexible distribution that in a finite interval.

- Beta distribution is a very flexible distribution that in a finite interval.
- $X \sim \text{Beta}(\alpha, \beta)$ with $\alpha, \beta > 0$, if its pdf is given by

$$f(x) = \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha, \beta)}, \quad x \in [0, 1].$$

- Beta distribution is a very flexible distribution that in a finite interval.
- $X \sim \text{Beta}(\alpha, \beta)$ with $\alpha, \beta > 0$, if its pdf is given by

$$f(x) = \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha, \beta)}, \quad x \in [0, 1].$$

- $\mathbb{E}[X] = \alpha/(\alpha + \beta)$, $\text{Var}(X) = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$.



- Beta distribution is a very flexible distribution that in a finite interval.
- $X \sim \text{Beta}(\alpha, \beta)$ with $\alpha, \beta > 0$, if its pdf is given by


$$f(x) = \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha, \beta)}, \quad x \in [0, 1].$$

- $\mathbb{E}[X] = \alpha/(\alpha + \beta)$, $\text{Var}(X) = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$.
- $B(\alpha, \beta) := \int_0^1 t^{\alpha-1}(1-t)^{\beta-1}dt$ is known as the beta function.
 - $B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$.

- Beta distribution is a very flexible distribution that in a finite interval.
- $X \sim \text{Beta}(\alpha, \beta)$ with $\alpha, \beta > 0$, if its pdf is given by

$$f(x) = \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha, \beta)}, \quad x \in [0, 1].$$

- $\mathbb{E}[X] = \alpha/(\alpha + \beta)$, $\text{Var}(X) = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$.
- $B(\alpha, \beta) := \int_0^1 t^{\alpha-1}(1-t)^{\beta-1}dt$ is known as the beta function.
 - $B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$.
- The pdf is quite flexible
 - $\alpha = 1, \beta = 1 \implies \text{Unif}(0, 1)$
 - $\alpha > 1, \beta = 1 \implies$ strictly increasing
 - $\alpha = 1, \beta > 1 \implies$ strictly decreasing
 - $\alpha < 1, \beta < 1 \implies$ U-shaped
 - $\alpha > 1, \beta > 1 \implies$ unimodal



- $X \sim$ Student's t distribution with p degrees of freedom, denoted as t_p , where p is an integer, if its pdf is given by

$$f(x) = \frac{\Gamma(\frac{p+1}{2})}{\Gamma(\frac{p}{2})} \frac{1}{(p\pi)^{1/2}} \frac{1}{(1 + x^2/p)^{(p+1)/2}}, \quad x \in \mathbb{R}.$$

- $X \sim$ Student's t distribution with p degrees of freedom, denoted as t_p , where p is an integer, if its pdf is given by

$$f(x) = \frac{\Gamma(\frac{p+1}{2})}{\Gamma(\frac{p}{2})} \frac{1}{(p\pi)^{1/2}} \frac{1}{(1 + x^2/p)^{(p+1)/2}}, \quad x \in \mathbb{R}.$$

- $\mathbb{E}[X] = 0$ if $p > 1$;



- $X \sim$ Student's t distribution with p degrees of freedom, denoted as t_p , where p is an integer, if its pdf is given by

$$f(x) = \frac{\Gamma(\frac{p+1}{2})}{\Gamma(\frac{p}{2})} \frac{1}{(p\pi)^{1/2}} \frac{1}{(1 + x^2/p)^{(p+1)/2}}, \quad x \in \mathbb{R}.$$

- $\mathbb{E}[X] = 0$ if $p > 1$;
- $\text{Var}(X) = p/(p-2)$ if $p > 2$.



- $X \sim$ Student's t distribution with p degrees of freedom, denoted as t_p , where p is an integer, if its pdf is given by

$$f(x) = \frac{\Gamma(\frac{p+1}{2})}{\Gamma(\frac{p}{2})} \frac{1}{(p\pi)^{1/2}} \frac{1}{(1 + x^2/p)^{(p+1)/2}}, \quad x \in \mathbb{R}.$$

- $\mathbb{E}[X] = 0$ if $p > 1$;
- $\text{Var}(X) = p/(p-2)$ if $p > 2$.
- t_1 is also known as the standard Cauchy distribution, or Cauchy(0, 1), whose pdf is simply

$$f(x) = \frac{1}{\pi(1 + x^2)}, \quad x \in \mathbb{R}.$$



- The **normal distribution** (sometimes called the Gaussian distribution) plays a **central role** in a large body of statistics.

- The **normal distribution** (sometimes called the Gaussian distribution) plays a **central role** in a large body of statistics.
- $X \sim$ normal distribution with mean μ and variance σ^2 , denoted as $\mathcal{N}(\mu, \sigma^2)$, with $\sigma > 0$, if its pdf is given by

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, \quad x \in \mathbb{R}.$$

- The **normal distribution** (sometimes called the Gaussian distribution) plays a **central role** in a large body of statistics.
- $X \sim$ normal distribution with mean μ and variance σ^2 , denoted as $\mathcal{N}(\mu, \sigma^2)$, with $\sigma > 0$, if its pdf is given by

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, \quad x \in \mathbb{R}.$$

- $\mathbb{E}[X] = \mu$, $\text{Var}(X) = \sigma^2$.



- The **normal distribution** (sometimes called the Gaussian distribution) plays a **central role** in a large body of statistics.
- $X \sim$ normal distribution with mean μ and variance σ^2 , denoted as $\mathcal{N}(\mu, \sigma^2)$, with $\sigma > 0$, if its pdf is given by

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, \quad x \in \mathbb{R}.$$

- $\mathbb{E}[X] = \mu$, $\text{Var}(X) = \sigma^2$.
- If $X \sim \mathcal{N}(\mu, \sigma^2)$, then $Z := (X - \mu)/\sigma \sim \mathcal{N}(0, 1)$.
 - Z is also known as the **standard normal** RV.
 - We often use $\Phi(z)$ and $\phi(z)$ to denote the CDF and pdf of Z .
 - $\mathbb{P}(X \leq x) = \Phi((x - \mu)/\sigma)$.



- The **normal distribution** (sometimes called the Gaussian distribution) plays a **central role** in a large body of statistics.
- $X \sim$ normal distribution with mean μ and variance σ^2 , denoted as $\mathcal{N}(\mu, \sigma^2)$, with $\sigma > 0$, if its pdf is given by

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, \quad x \in \mathbb{R}.$$

- $\mathbb{E}[X] = \mu$, $\text{Var}(X) = \sigma^2$.
- If $X \sim \mathcal{N}(\mu, \sigma^2)$, then $Z := (X - \mu)/\sigma \sim \mathcal{N}(0, 1)$.
 - Z is also known as the **standard normal** RV.
 - We often use $\Phi(z)$ and $\phi(z)$ to denote the CDF and pdf of Z .
 - $\mathbb{P}(X \leq x) = \Phi((x - \mu)/\sigma)$.
- If $X \sim \mathcal{N}(\mu, \sigma^2)$, then $a + bX \sim \mathcal{N}(a + b\mu, b^2\sigma^2)$ for $b > 0$.

- The **normal distribution** (sometimes called the Gaussian distribution) plays a **central role** in a large body of statistics.
- $X \sim$ normal distribution with mean μ and variance σ^2 , denoted as $\mathcal{N}(\mu, \sigma^2)$, with $\sigma > 0$, if its pdf is given by

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, \quad x \in \mathbb{R}.$$

- $\mathbb{E}[X] = \mu$, $\text{Var}(X) = \sigma^2$.
- If $X \sim \mathcal{N}(\mu, \sigma^2)$, then $Z := (X - \mu)/\sigma \sim \mathcal{N}(0, 1)$.
 - Z is also known as the **standard normal** RV.
 - We often use $\Phi(z)$ and $\phi(z)$ to denote the CDF and pdf of Z .
 - $\mathbb{P}(X \leq x) = \Phi((x - \mu)/\sigma)$.
- If $X \sim \mathcal{N}(\mu, \sigma^2)$, then $a + bX \sim \mathcal{N}(a + b\mu, b^2\sigma^2)$ for $b > 0$.
- If $X_1 \sim \mathcal{N}(\mu_1, \sigma_1^2)$ and $X_2 \sim \mathcal{N}(\mu_2, \sigma_2^2)$ are independent, then $X_1 + X_2 \sim \mathcal{N}(\mu_1 + \mu_2, \sigma_1^2 + \sigma_2^2)$.

- If $Z \sim \mathcal{N}(0, 1)$, then $Z^2 \sim \chi_1^2$.



- If $Z \sim \mathcal{N}(0, 1)$, then $Z^2 \sim \chi_1^2$.

Proof. Let $Y := Z^2$. For $y \in [0, \infty)$,

$$\mathbb{P}(Y \leq y) = \mathbb{P}(Z^2 \leq y) = \mathbb{P}(-\sqrt{y} \leq Z \leq \sqrt{y}) = \int_{-\sqrt{y}}^{\sqrt{y}} \phi(t) dt =: F(y).$$



- If $Z \sim \mathcal{N}(0, 1)$, then $Z^2 \sim \chi_1^2$.

Proof. Let $Y := Z^2$. For $y \in [0, \infty)$,

$$\mathbb{P}(Y \leq y) = \mathbb{P}(Z^2 \leq y) = \mathbb{P}(-\sqrt{y} \leq Z \leq \sqrt{y}) = \int_{-\sqrt{y}}^{\sqrt{y}} \phi(t) dt =: F(y).$$

Then,

$$f(y) = \frac{d}{dy} F(y) = \phi(\sqrt{y}) \frac{d}{dy} \sqrt{y} - \phi(-\sqrt{y}) \frac{d}{dy} (-\sqrt{y})$$



- If $Z \sim \mathcal{N}(0, 1)$, then $Z^2 \sim \chi_1^2$.

Proof. Let $Y := Z^2$. For $y \in [0, \infty)$,

$$\mathbb{P}(Y \leq y) = \mathbb{P}(Z^2 \leq y) = \mathbb{P}(-\sqrt{y} \leq Z \leq \sqrt{y}) = \int_{-\sqrt{y}}^{\sqrt{y}} \phi(t) dt =: F(y).$$

Then,

$$\begin{aligned} f(y) &= \frac{d}{dy} F(y) = \phi(\sqrt{y}) \frac{d}{dy} \sqrt{y} - \phi(-\sqrt{y}) \frac{d}{dy} (-\sqrt{y}) \\ &= 2\phi(\sqrt{y}) \frac{d}{dy} \sqrt{y} = \frac{1}{\sqrt{2\pi}} e^{-\frac{y}{2}} y^{-\frac{1}{2}}. \end{aligned}$$



- If $Z \sim \mathcal{N}(0, 1)$, then $Z^2 \sim \chi_1^2$.

Proof. Let $Y := Z^2$. For $y \in [0, \infty)$,

$$\mathbb{P}(Y \leq y) = \mathbb{P}(Z^2 \leq y) = \mathbb{P}(-\sqrt{y} \leq Z \leq \sqrt{y}) = \int_{-\sqrt{y}}^{\sqrt{y}} \phi(t) dt =: F(y).$$

Then,

$$\begin{aligned} f(y) &= \frac{d}{dy} F(y) = \phi(\sqrt{y}) \frac{d}{dy} \sqrt{y} - \phi(-\sqrt{y}) \frac{d}{dy} (-\sqrt{y}) \\ &= 2\phi(\sqrt{y}) \frac{d}{dy} \sqrt{y} = \frac{1}{\sqrt{2\pi}} e^{-\frac{y}{2}} y^{-\frac{1}{2}}. \end{aligned}$$

If $Y \sim \chi_1^2$, i.e., $Y \sim \text{Gamma}(1/2, 1/2)$, it means its pdf is

$$f(y) = \frac{1}{\sqrt{2\pi}\Gamma(\frac{1}{2})} y^{-\frac{1}{2}} e^{-\frac{y}{2}}.$$



- If $Z \sim \mathcal{N}(0, 1)$, then $Z^2 \sim \chi_1^2$.

Proof. Let $Y := Z^2$. For $y \in [0, \infty)$,

$$\mathbb{P}(Y \leq y) = \mathbb{P}(Z^2 \leq y) = \mathbb{P}(-\sqrt{y} \leq Z \leq \sqrt{y}) = \int_{-\sqrt{y}}^{\sqrt{y}} \phi(t) dt =: F(y).$$

Then,

$$\begin{aligned} f(y) &= \frac{d}{dy} F(y) = \phi(\sqrt{y}) \frac{d}{dy} \sqrt{y} - \phi(-\sqrt{y}) \frac{d}{dy} (-\sqrt{y}) \\ &= 2\phi(\sqrt{y}) \frac{d}{dy} \sqrt{y} = \frac{1}{\sqrt{2\pi}} e^{-\frac{y}{2}} y^{-\frac{1}{2}}. \end{aligned}$$

If $Y \sim \chi_1^2$, i.e., $Y \sim \text{Gamma}(1/2, 1/2)$, it means its pdf is

$$f(y) = \frac{1}{\sqrt{2}\Gamma(\frac{1}{2})} y^{-\frac{1}{2}} e^{-\frac{y}{2}}.$$

The proof is completed by showing that $\Gamma(\frac{1}{2}) = \int_0^\infty t^{-\frac{1}{2}} e^{-t} dt = \sqrt{\pi}$, which can be seen if we convert to polar coordinates.

- If $Z \sim \mathcal{N}(0, 1)$ and $V \sim \chi_p^2$ are independent, then $\frac{Z}{\sqrt{V/p}} \sim t_p$.

- If $Z \sim \mathcal{N}(0, 1)$ and $V \sim \chi_p^2$ are independent, then $\frac{Z}{\sqrt{V/p}} \sim t_p$.

Proof. Since $V \sim \chi_p^2$, by definition, its pdf is

$$f_V(v) = \frac{\left(\frac{1}{2}\right)^{\frac{p}{2}}}{\Gamma\left(\frac{p}{2}\right)} v^{\frac{p}{2}-1} e^{-\frac{1}{2}v}, \quad v \in [0, \infty).$$

- If $Z \sim \mathcal{N}(0, 1)$ and $V \sim \chi_p^2$ are independent, then $\frac{Z}{\sqrt{V/p}} \sim t_p$.

Proof. Since $V \sim \chi_p^2$, by definition, its pdf is

$$f_V(v) = \frac{\left(\frac{1}{2}\right)^{\frac{p}{2}}}{\Gamma\left(\frac{p}{2}\right)} v^{\frac{p}{2}-1} e^{-\frac{1}{2}v}, \quad v \in [0, \infty).$$

Let $Y := \sqrt{V/p}$. For $y \in [0, \infty)$,

$$f_Y(y) = \frac{d}{dy} \mathbb{P}(Y \leq y) = \frac{d}{dy} \mathbb{P}(V \leq py^2) = \frac{d}{dy} \int_0^{py^2} f_V(v) dv = 2py f_V(py^2).$$

- If $Z \sim \mathcal{N}(0, 1)$ and $V \sim \chi_p^2$ are independent, then $\frac{Z}{\sqrt{V/p}} \sim t_p$.

Proof. Since $V \sim \chi_p^2$, by definition, its pdf is

$$f_V(v) = \frac{\left(\frac{1}{2}\right)^{\frac{p}{2}}}{\Gamma\left(\frac{p}{2}\right)} v^{\frac{p}{2}-1} e^{-\frac{1}{2}v}, \quad v \in [0, \infty).$$

Let $Y := \sqrt{V/p}$. For $y \in [0, \infty)$,

$$f_Y(y) = \frac{d}{dy} \mathbb{P}(Y \leq y) = \frac{d}{dy} \mathbb{P}(V \leq py^2) = \frac{d}{dy} \int_0^{py^2} f_V(v) dv = 2py f_V(py^2).$$

Let $T := \frac{Z}{\sqrt{V/p}} = \frac{Z}{Y}$. For $t \in \mathbb{R}$,

$$\mathbb{P}(T \leq t) = \mathbb{P}\left(\frac{Z}{Y} \leq t\right) = \mathbb{P}(Z \leq tY) = \int_0^\infty \mathbb{P}(Z \leq ty) f_Y(y) dy. \quad (\text{Why?})$$



- If $Z \sim \mathcal{N}(0, 1)$ and $V \sim \chi_p^2$ are independent, then $\frac{Z}{\sqrt{V/p}} \sim t_p$.

Proof. Since $V \sim \chi_p^2$, by definition, its pdf is

$$f_V(v) = \frac{\left(\frac{1}{2}\right)^{\frac{p}{2}}}{\Gamma\left(\frac{p}{2}\right)} v^{\frac{p}{2}-1} e^{-\frac{1}{2}v}, \quad v \in [0, \infty).$$

Let $Y := \sqrt{V/p}$. For $y \in [0, \infty)$,

$$f_Y(y) = \frac{d}{dy} \mathbb{P}(Y \leq y) = \frac{d}{dy} \mathbb{P}(V \leq py^2) = \frac{d}{dy} \int_0^{py^2} f_V(v) dv = 2py f_V(py^2).$$

Let $T := \frac{Z}{\sqrt{V/p}} = \frac{Z}{Y}$. For $t \in \mathbb{R}$,

$$\mathbb{P}(T \leq t) = \mathbb{P}\left(\frac{Z}{Y} \leq t\right) = \mathbb{P}(Z \leq tY) = \int_0^\infty \mathbb{P}(Z \leq ty) f_Y(y) dy. \quad (\text{Why?})$$

Then,

$$f_T(t) = \frac{d}{dt} \mathbb{P}(T \leq t) = \int_0^\infty \frac{d}{dt} \mathbb{P}(Z \leq ty) f_Y(y) dy.$$



Proof. (Cont'd) Note that $\frac{d}{dt} \mathbb{P}(Z \leq ty) = \frac{d}{dt} \int_{-\infty}^{ty} \phi(z) dz = y\phi(ty)$.



Proof. (Cont'd) Note that $\frac{d}{dt} \mathbb{P}(Z \leq ty) = \frac{d}{dt} \int_{-\infty}^{ty} \phi(z) dz = y\phi(ty)$. So,

$$f_T(t) = \int_0^\infty y\phi(ty) f_Y(y) dy = \int_0^\infty y\phi(ty) 2py f_V(py^2) dy$$



Proof. (Cont'd) Note that $\frac{d}{dt} \mathbb{P}(Z \leq ty) = \frac{d}{dt} \int_{-\infty}^{ty} \phi(z) dz = y\phi(ty)$. So,

$$\begin{aligned} f_T(t) &= \int_0^\infty y\phi(ty) f_Y(y) dy = \int_0^\infty y\phi(ty) 2py f_V(py^2) dy \\ &= \int_0^\infty 2py^2 \cdot \frac{1}{\sqrt{2\pi}} e^{-\frac{t^2 y^2}{2}} \cdot \frac{(\frac{1}{2})^{\frac{p}{2}}}{\Gamma(\frac{p}{2})} (py^2)^{\frac{p}{2}-1} e^{-\frac{1}{2}py^2} dy \end{aligned}$$



Proof. (Cont'd) Note that $\frac{d}{dt} \mathbb{P}(Z \leq ty) = \frac{d}{dt} \int_{-\infty}^{ty} \phi(z) dz = y\phi(ty)$. So,

$$\begin{aligned} f_T(t) &= \int_0^\infty y\phi(ty) f_Y(y) dy = \int_0^\infty y\phi(ty) 2py f_V(py^2) dy \\ &= \int_0^\infty 2py^2 \cdot \frac{1}{\sqrt{2\pi}} e^{-\frac{t^2 y^2}{2}} \cdot \frac{(\frac{1}{2})^{\frac{p}{2}}}{\Gamma(\frac{p}{2})} (py^2)^{\frac{p}{2}-1} e^{-\frac{1}{2}py^2} dy \\ &= \frac{1}{\Gamma(\frac{p}{2})} \frac{1}{(p\pi)^{1/2}} 2^{\frac{1-p}{2}} p^{\frac{p+1}{2}} \int_0^\infty y^p e^{-\frac{1}{2}(t^2+p)y^2} dy. \end{aligned}$$



Proof. (Cont'd) Note that $\frac{d}{dt} \mathbb{P}(Z \leq ty) = \frac{d}{dt} \int_{-\infty}^{ty} \phi(z) dz = y\phi(ty)$. So,

$$\begin{aligned} f_T(t) &= \int_0^\infty y\phi(ty) f_Y(y) dy = \int_0^\infty y\phi(ty) 2py f_V(py^2) dy \\ &= \int_0^\infty 2py^2 \cdot \frac{1}{\sqrt{2\pi}} e^{-\frac{t^2 y^2}{2}} \cdot \frac{(\frac{1}{2})^{\frac{p}{2}}}{\Gamma(\frac{p}{2})} (py^2)^{\frac{p}{2}-1} e^{-\frac{1}{2}py^2} dy \\ &= \frac{1}{\Gamma(\frac{p}{2})} \frac{1}{(p\pi)^{1/2}} 2^{\frac{1-p}{2}} p^{\frac{p+1}{2}} \int_0^\infty y^p e^{-\frac{1}{2}(t^2+p)y^2} dy. \end{aligned}$$

Let $x := y^2$. Then, integration by substitution shows that

$$\int_0^\infty y^p e^{-\frac{1}{2}(t^2+p)y^2} dy = \frac{1}{2} \int_0^\infty x^{\frac{p-1}{2}} e^{-\frac{1}{2}(t^2+p)x} dx =: \frac{1}{2} \int_0^\infty x^{\alpha-1} e^{-\lambda x} dx,$$

where $\alpha := \frac{p+1}{2}$ and $\lambda := \frac{1}{2}(t^2 + p)$.



Proof. (Cont'd) Note that $\frac{d}{dt} \mathbb{P}(Z \leq ty) = \frac{d}{dt} \int_{-\infty}^{ty} \phi(z) dz = y\phi(ty)$. So,

$$\begin{aligned} f_T(t) &= \int_0^\infty y\phi(ty)f_Y(y)dy = \int_0^\infty y\phi(ty)2pyf_V(py^2)dy \\ &= \int_0^\infty 2py^2 \cdot \frac{1}{\sqrt{2\pi}} e^{-\frac{t^2 y^2}{2}} \cdot \frac{(\frac{1}{2})^{\frac{p}{2}}}{\Gamma(\frac{p}{2})} (py^2)^{\frac{p}{2}-1} e^{-\frac{1}{2}py^2} dy \\ &= \frac{1}{\Gamma(\frac{p}{2})} \frac{1}{(p\pi)^{1/2}} 2^{\frac{1-p}{2}} p^{\frac{p+1}{2}} \int_0^\infty y^p e^{-\frac{1}{2}(t^2+p)y^2} dy. \end{aligned}$$

Let $x := y^2$. Then, integration by substitution shows that

$$\int_0^\infty y^p e^{-\frac{1}{2}(t^2+p)y^2} dy = \frac{1}{2} \int_0^\infty x^{\frac{p-1}{2}} e^{-\frac{1}{2}(t^2+p)x} dx =: \frac{1}{2} \int_0^\infty x^{\alpha-1} e^{-\lambda x} dx,$$

where $\alpha := \frac{p+1}{2}$ and $\lambda := \frac{1}{2}(t^2+p)$. Recalling the pdf of $\Gamma(\alpha, \lambda)$, it is easy to see that $\int_0^\infty x^{\alpha-1} e^{-\lambda x} dx = \Gamma(\alpha)/\lambda^\alpha$.



Proof. (Cont'd) Note that $\frac{d}{dt} \mathbb{P}(Z \leq ty) = \frac{d}{dt} \int_{-\infty}^{ty} \phi(z) dz = y\phi(ty)$. So,

$$\begin{aligned} f_T(t) &= \int_0^\infty y\phi(ty)f_Y(y)dy = \int_0^\infty y\phi(ty)2pyf_V(py^2)dy \\ &= \int_0^\infty 2py^2 \cdot \frac{1}{\sqrt{2\pi}} e^{-\frac{t^2 y^2}{2}} \cdot \frac{(\frac{1}{2})^{\frac{p}{2}}}{\Gamma(\frac{p}{2})} (py^2)^{\frac{p}{2}-1} e^{-\frac{1}{2}py^2} dy \\ &= \frac{1}{\Gamma(\frac{p}{2})} \frac{1}{(p\pi)^{1/2}} 2^{\frac{1-p}{2}} p^{\frac{p+1}{2}} \int_0^\infty y^p e^{-\frac{1}{2}(t^2+p)y^2} dy. \end{aligned}$$

Let $x := y^2$. Then, integration by substitution shows that

$$\int_0^\infty y^p e^{-\frac{1}{2}(t^2+p)y^2} dy = \frac{1}{2} \int_0^\infty x^{\frac{p-1}{2}} e^{-\frac{1}{2}(t^2+p)x} dx =: \frac{1}{2} \int_0^\infty x^{\alpha-1} e^{-\lambda x} dx,$$

where $\alpha := \frac{p+1}{2}$ and $\lambda := \frac{1}{2}(t^2+p)$. Recalling the pdf of $\Gamma(\alpha, \lambda)$, it is easy to see that $\int_0^\infty x^{\alpha-1} e^{-\lambda x} dx = \Gamma(\alpha)/\lambda^\alpha$. Finally,

$$\begin{aligned} f_T(t) &= \frac{1}{\Gamma(\frac{p}{2})} \frac{1}{(p\pi)^{1/2}} 2^{\frac{1-p}{2}} p^{\frac{p+1}{2}} \cdot \frac{1}{2} \frac{\Gamma(\frac{p+1}{2})}{(1/2)^{(p+1)/2} (t^2+p)^{(p+1)/2}} \\ &= \frac{\Gamma(\frac{p+1}{2})}{\Gamma(\frac{p}{2})} \frac{1}{(p\pi)^{1/2}} \frac{1}{(1+t^2/p)^{(p+1)/2}}. \end{aligned}$$



- $\mathbf{X} := (X_1, \dots, X_k)^\top$ is said to follow a k -variate normal distribution, if **every** linear combination of X_1, \dots, X_k follows a (univariate) normal distribution.

- $\mathbf{X} := (X_1, \dots, X_k)^\top$ is said to follow a k -variate normal distribution, if **every** linear combination of X_1, \dots, X_k follows a (univariate) normal distribution.
 - \mathbf{X} is also called a (k dimensional) normal random vector.

- $\mathbf{X} := (X_1, \dots, X_k)^\top$ is said to follow a k -variate normal distribution, if **every** linear combination of X_1, \dots, X_k follows a (univariate) normal distribution.
 - \mathbf{X} is also called a (k dimensional) normal random vector.
 - If $k = 2$, $\mathbf{X} = (X_1, X_2)^\top$ is also said to follow a *bivariate* normal distribution.

- $\mathbf{X} := (X_1, \dots, X_k)^\top$ is said to follow a k -variate normal distribution, if **every** linear combination of X_1, \dots, X_k follows a (univariate) normal distribution.
 - \mathbf{X} is also called a (k dimensional) normal random vector.
 - If $k = 2$, $\mathbf{X} = (X_1, X_2)^\top$ is also said to follow a *bivariate* normal distribution.
- $\mathbf{X} \sim$ a k -variate normal distribution, denoted as $\mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, if its joint pdf is given by

$$f(\mathbf{x}) = \frac{1}{(2\pi)^{k/2} |\boldsymbol{\Sigma}|^{1/2}} e^{-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{x}-\boldsymbol{\mu})}, \mathbf{x} \in \mathbb{R}^k,$$

where $|\boldsymbol{\Sigma}|$ is the determinant of $\boldsymbol{\Sigma}$.

- $\mathbf{X} := (X_1, \dots, X_k)^\top$ is said to follow a k -variate normal distribution, if **every** linear combination of X_1, \dots, X_k follows a (univariate) normal distribution.
 - \mathbf{X} is also called a (k dimensional) normal random vector.
 - If $k = 2$, $\mathbf{X} = (X_1, X_2)^\top$ is also said to follow a *bivariate* normal distribution.
- $\mathbf{X} \sim$ a k -variate normal distribution, denoted as $\mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, if its joint pdf is given by

$$f(\mathbf{x}) = \frac{1}{(2\pi)^{k/2} |\boldsymbol{\Sigma}|^{1/2}} e^{-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{x}-\boldsymbol{\mu})}, \mathbf{x} \in \mathbb{R}^k,$$

where $|\boldsymbol{\Sigma}|$ is the determinant of $\boldsymbol{\Sigma}$.

- $\boldsymbol{\mu} = (\mu_1, \dots, \mu_k)^\top = \mathbb{E}[\mathbf{X}] = (\mathbb{E}[X_1], \dots, \mathbb{E}[X_k])^\top \in \mathbb{R}^k$.
- $\boldsymbol{\Sigma} = (\Sigma_{ij}) = \text{Cov}(\mathbf{X}, \mathbf{X}) = (\text{Cov}(Z_i, Z_j)) \in \mathbb{R}^{k \times k}$.

- $\mathbf{X} := (X_1, \dots, X_k)^\top$ is said to follow a k -variate normal distribution, if **every** linear combination of X_1, \dots, X_k follows a (univariate) normal distribution.
 - \mathbf{X} is also called a (k dimensional) normal random vector.
 - If $k = 2$, $\mathbf{X} = (X_1, X_2)^\top$ is also said to follow a *bivariate* normal distribution.
- $\mathbf{X} \sim$ a k -variate normal distribution, denoted as $\mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, if its joint pdf is given by

$$f(\mathbf{x}) = \frac{1}{(2\pi)^{k/2} |\boldsymbol{\Sigma}|^{1/2}} e^{-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{x}-\boldsymbol{\mu})}, \mathbf{x} \in \mathbb{R}^k,$$

where $|\boldsymbol{\Sigma}|$ is the determinant of $\boldsymbol{\Sigma}$.

- $\boldsymbol{\mu} = (\mu_1, \dots, \mu_k)^\top = \mathbb{E}[\mathbf{X}] = (\mathbb{E}[X_1], \dots, \mathbb{E}[X_k])^\top \in \mathbb{R}^k$.
- $\boldsymbol{\Sigma} = (\Sigma_{ij}) = \text{Cov}(\mathbf{X}, \mathbf{X}) = (\text{Cov}(Z_i, Z_j)) \in \mathbb{R}^{k \times k}$.
- $\boldsymbol{\Sigma}$ is a symmetric and positive definite matrix.

- $\mathbf{X} := (X_1, \dots, X_k)^\top$ is said to follow a k -variate normal distribution, if **every** linear combination of X_1, \dots, X_k follows a (univariate) normal distribution.
 - \mathbf{X} is also called a (k dimensional) normal random vector.
 - If $k = 2$, $\mathbf{X} = (X_1, X_2)^\top$ is also said to follow a *bivariate* normal distribution.
- $\mathbf{X} \sim$ a k -variate normal distribution, denoted as $\mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, if its joint pdf is given by

$$f(\mathbf{x}) = \frac{1}{(2\pi)^{k/2} |\boldsymbol{\Sigma}|^{1/2}} e^{-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{x}-\boldsymbol{\mu})}, \mathbf{x} \in \mathbb{R}^k,$$

where $|\boldsymbol{\Sigma}|$ is the determinant of $\boldsymbol{\Sigma}$.

- $\boldsymbol{\mu} = (\mu_1, \dots, \mu_k)^\top = \mathbb{E}[\mathbf{X}] = (\mathbb{E}[X_1], \dots, \mathbb{E}[X_k])^\top \in \mathbb{R}^k$.
- $\boldsymbol{\Sigma} = (\Sigma_{ij}) = \text{Cov}(\mathbf{X}, \mathbf{X}) = (\text{Cov}(Z_i, Z_j)) \in \mathbb{R}^{k \times k}$.
- $\boldsymbol{\Sigma}$ is a symmetric and positive definite matrix.
- $X_i \sim \mathcal{N}(\mu_i, \sigma_i^2)$, $i = 1, \dots, k$.



- If $\mathbf{X} \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ is k dimensional, then
 - $\mathbf{Z} := \mathbf{A}^{-1}(\mathbf{X} - \boldsymbol{\mu}) \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$, where \mathbf{A} satisfies $\boldsymbol{\Sigma} = \mathbf{A}\mathbf{A}^\top$ (Cholesky decomposition), $\mathbf{0} \in \mathbb{R}^k$, and $\mathbf{I} \in \mathbb{R}^{k \times k}$ denotes the identity matrix.



- If $\mathbf{X} \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ is k dimensional, then
 - $\mathbf{Z} := \mathbf{A}^{-1}(\mathbf{X} - \boldsymbol{\mu}) \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$, where \mathbf{A} satisfies $\boldsymbol{\Sigma} = \mathbf{A}\mathbf{A}^\top$ (Cholesky decomposition), $\mathbf{0} \in \mathbb{R}^k$, and $\mathbf{I} \in \mathbb{R}^{k \times k}$ denotes the identity matrix.
 - $\mathbf{Z} = (Z_1, \dots, Z_k)^\top$, where $Z_i \sim \mathcal{N}(0, 1)$, $i = 1, \dots, k$, **iid**.



- If $\mathbf{X} \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ is k dimensional, then
 - $\mathbf{Z} := \mathbf{A}^{-1}(\mathbf{X} - \boldsymbol{\mu}) \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$, where \mathbf{A} satisfies $\boldsymbol{\Sigma} = \mathbf{A}\mathbf{A}^\top$ (Cholesky decomposition), $\mathbf{0} \in \mathbb{R}^k$, and $\mathbf{I} \in \mathbb{R}^{k \times k}$ denotes the identity matrix.
 - $\mathbf{Z} = (Z_1, \dots, Z_k)^\top$, where $Z_i \sim \mathcal{N}(0, 1)$, $i = 1, \dots, k$, **iid**.
 - $\mathbf{a} + \mathbf{B}\mathbf{X} \sim \mathcal{N}(\mathbf{a} + \mathbf{B}\boldsymbol{\mu}, \mathbf{B}\boldsymbol{\Sigma}\mathbf{B}^\top)$.[†]

[†]The multivariate normal distribution will be degenerate if \mathbf{B} does not have full row rank (\mathbf{B} 不行满秩).
SHANGHAI JIAO TONG UNIVERSITY

- If $\mathbf{X} \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ is k dimensional, then
 - $\mathbf{Z} := \mathbf{A}^{-1}(\mathbf{X} - \boldsymbol{\mu}) \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$, where \mathbf{A} satisfies $\boldsymbol{\Sigma} = \mathbf{A}\mathbf{A}^\top$ (Cholesky decomposition), $\mathbf{0} \in \mathbb{R}^k$, and $\mathbf{I} \in \mathbb{R}^{k \times k}$ denotes the identity matrix.
 - $\mathbf{Z} = (Z_1, \dots, Z_k)^\top$, where $Z_i \sim \mathcal{N}(0, 1)$, $i = 1, \dots, k$, **iid**.
 - $\mathbf{a} + \mathbf{B}\mathbf{X} \sim \mathcal{N}(\mathbf{a} + \mathbf{B}\boldsymbol{\mu}, \mathbf{B}\boldsymbol{\Sigma}\mathbf{B}^\top)$.[†]
- Suppose \mathbf{X} is a k dimensional random vector. Then,
 $\mathbf{X} \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma}) \iff$
 There exist $\boldsymbol{\mu} \in \mathbb{R}^k$ and $\mathbf{A} \in \mathbb{R}^{k \times \ell}$ such that $\mathbf{X} = \boldsymbol{\mu} + \mathbf{A}\mathbf{Z}$,
 where $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$ with $\mathbf{0} \in \mathbb{R}^\ell$ and $\mathbf{I} \in \mathbb{R}^{\ell \times \ell}$.

[†]The multivariate normal distribution will be degenerate if \mathbf{B} does not have full row rank (\mathbf{B} 不行满秩).

- If $\mathbf{X} \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ is k dimensional, then
 - $\mathbf{Z} := \mathbf{A}^{-1}(\mathbf{X} - \boldsymbol{\mu}) \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$, where \mathbf{A} satisfies $\boldsymbol{\Sigma} = \mathbf{A}\mathbf{A}^\top$ (Cholesky decomposition), $\mathbf{0} \in \mathbb{R}^k$, and $\mathbf{I} \in \mathbb{R}^{k \times k}$ denotes the identity matrix.
 - $\mathbf{Z} = (Z_1, \dots, Z_k)^\top$, where $Z_i \sim \mathcal{N}(0, 1)$, $i = 1, \dots, k$, **iid**.
 - $\mathbf{a} + \mathbf{B}\mathbf{X} \sim \mathcal{N}(\mathbf{a} + \mathbf{B}\boldsymbol{\mu}, \mathbf{B}\boldsymbol{\Sigma}\mathbf{B}^\top)$.[†]
- Suppose \mathbf{X} is a k dimensional random vector. Then,
 $\mathbf{X} \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma}) \iff$
 There exist $\boldsymbol{\mu} \in \mathbb{R}^k$ and $\mathbf{A} \in \mathbb{R}^{k \times \ell}$ such that $\mathbf{X} = \boldsymbol{\mu} + \mathbf{A}\mathbf{Z}$,
 where $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$ with $\mathbf{0} \in \mathbb{R}^\ell$ and $\mathbf{I} \in \mathbb{R}^{\ell \times \ell}$.
 - Such \mathbf{A} must satisfy $\boldsymbol{\Sigma} = \mathbf{A}\mathbf{A}^\top$.

[†]The multivariate normal distribution will be degenerate if \mathbf{B} does not have full row rank (\mathbf{B} 不行满秩).

- Bivariate normal distribution: $(X_1, X_2)^T \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, where $\boldsymbol{\mu} = (\mu_1, \mu_2)^T$, and

$$\boldsymbol{\Sigma} = \begin{bmatrix} \text{Cov}(X_1, X_1) & \text{Cov}(X_1, X_2) \\ \text{Cov}(X_2, X_1) & \text{Cov}(X_2, X_2) \end{bmatrix} =: \begin{bmatrix} \sigma_1^2 & \rho\sigma_1\sigma_2 \\ \rho\sigma_1\sigma_2 & \sigma_2^2 \end{bmatrix},$$

and the joint pdf is

$$f(x_1, x_2) = \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} \times e^{-\frac{1}{2(1-\rho^2)} \left[\left(\frac{x_1-\mu_1}{\sigma_1} \right)^2 - 2\rho \left(\frac{x_1-\mu_1}{\sigma_1} \right) \left(\frac{x_2-\mu_2}{\sigma_2} \right) + \left(\frac{x_2-\mu_2}{\sigma_2} \right)^2 \right]}.$$



- Bivariate normal distribution: $(X_1, X_2)^T \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, where $\boldsymbol{\mu} = (\mu_1, \mu_2)^T$, and

$$\boldsymbol{\Sigma} = \begin{bmatrix} \text{Cov}(X_1, X_1) & \text{Cov}(X_1, X_2) \\ \text{Cov}(X_2, X_1) & \text{Cov}(X_2, X_2) \end{bmatrix} =: \begin{bmatrix} \sigma_1^2 & \rho\sigma_1\sigma_2 \\ \rho\sigma_1\sigma_2 & \sigma_2^2 \end{bmatrix},$$

and the joint pdf is

$$f(x_1, x_2) = \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} \times e^{-\frac{1}{2(1-\rho^2)} \left[\left(\frac{x_1-\mu_1}{\sigma_1} \right)^2 - 2\rho \left(\frac{x_1-\mu_1}{\sigma_1} \right) \left(\frac{x_2-\mu_2}{\sigma_2} \right) + \left(\frac{x_2-\mu_2}{\sigma_2} \right)^2 \right]}.$$

- To see $\rho = 0 \implies X_1 \perp X_2$, let $\rho = 0$, and note

$$\begin{aligned} f(x_1, x_2) &= \frac{1}{2\pi\sigma_1\sigma_2} e^{-\frac{1}{2} \left[\left(\frac{x_1-\mu_1}{\sigma_1} \right)^2 + \left(\frac{x_2-\mu_2}{\sigma_2} \right)^2 \right]} \\ &= \frac{1}{\sqrt{2\pi}\sigma_1} e^{-\frac{(x_1-\mu_1)^2}{2\sigma_1^2}} \times \frac{1}{\sqrt{2\pi}\sigma_2} e^{-\frac{(x_2-\mu_2)^2}{2\sigma_2^2}} = f_{X_1}(x_1)f_{X_2}(x_2). \end{aligned}$$

- If $(X_1, X_2)^\top \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ and $X_i \sim \mathcal{N}(\mu_i, \sigma^2)$, $i = 1, 2$, then $X_1 + X_2 \perp X_1 - X_2$.



- If $(X_1, X_2)^\top \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ and $X_i \sim \mathcal{N}(\mu_i, \sigma^2)$, $i = 1, 2$, then $X_1 + X_2 \perp X_1 - X_2$.

Proof. Note that

$$\mathbf{Y} := \begin{bmatrix} X_1 + X_2 \\ X_1 - X_2 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} X_1 \\ X_2 \end{bmatrix} =: \mathbf{B} \begin{bmatrix} X_1 \\ X_2 \end{bmatrix}.$$

- If $(X_1, X_2)^\top \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ and $X_i \sim \mathcal{N}(\mu_i, \sigma^2)$, $i = 1, 2$, then $X_1 + X_2 \perp X_1 - X_2$.

Proof. Note that

$$\mathbf{Y} := \begin{bmatrix} X_1 + X_2 \\ X_1 - X_2 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} X_1 \\ X_2 \end{bmatrix} =: \mathbf{B} \begin{bmatrix} X_1 \\ X_2 \end{bmatrix}.$$

Since \mathbf{B} has full row rank, $\mathbf{Y} \sim \mathcal{N}(\mathbf{B}\boldsymbol{\mu}, \mathbf{B}\boldsymbol{\Sigma}\mathbf{B}^\top)$, which is non-degenerate.

- If $(X_1, X_2)^\top \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ and $X_i \sim \mathcal{N}(\mu_i, \sigma^2)$, $i = 1, 2$, then $X_1 + X_2 \perp X_1 - X_2$.

Proof. Note that

$$\mathbf{Y} := \begin{bmatrix} X_1 + X_2 \\ X_1 - X_2 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} X_1 \\ X_2 \end{bmatrix} =: \mathbf{B} \begin{bmatrix} X_1 \\ X_2 \end{bmatrix}.$$

Since \mathbf{B} has full row rank, $\mathbf{Y} \sim \mathcal{N}(\mathbf{B}\boldsymbol{\mu}, \mathbf{B}\boldsymbol{\Sigma}\mathbf{B}^\top)$, which is non-degenerate. Hence, to prove $X_1 + X_2 \perp X_1 - X_2$, it suffices to show $\text{Cov}(X_1 + X_2, X_1 - X_2) = 0$.

- If $(X_1, X_2)^\top \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ and $X_i \sim \mathcal{N}(\mu_i, \sigma^2)$, $i = 1, 2$, then $X_1 + X_2 \perp X_1 - X_2$.

Proof. Note that

$$\mathbf{Y} := \begin{bmatrix} X_1 + X_2 \\ X_1 - X_2 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} X_1 \\ X_2 \end{bmatrix} =: \mathbf{B} \begin{bmatrix} X_1 \\ X_2 \end{bmatrix}.$$

Since \mathbf{B} has full row rank, $\mathbf{Y} \sim \mathcal{N}(\mathbf{B}\boldsymbol{\mu}, \mathbf{B}\boldsymbol{\Sigma}\mathbf{B}^\top)$, which is non-degenerate. Hence, to prove $X_1 + X_2 \perp X_1 - X_2$, it suffices to show $\text{Cov}(X_1 + X_2, X_1 - X_2) = 0$. Note that

$$\begin{aligned} \text{Cov}(X_1 + X_2, X_1 - X_2) &= \text{Cov}(X_1, X_1) - \text{Cov}(X_2, X_2) \\ &= \sigma^2 - \sigma^2 = 0. \end{aligned}$$



-
- The diagram is a complex flowchart titled "A Comprehensive Classification of Probability Distributions". It maps various probability distributions to their parent distributions and associated mathematical functions. The chart is organized into several columns and rows, with arrows indicating the relationships between distributions. Key distributions include Negative Hypergeometric, Hypergeometric, Bernoulli, Poisson, Discrete Uniform, Geometric, Negative Binomial, Binomial, Poisson, and Liquid-Spin Uniform. The chart also includes various mathematical functions like the Gamma function, Beta function, and Hypergeometric function, and their relationships to the distributions. The chart is a comprehensive reference for probability distributions and their mathematical properties.
- Top Row (Parent Distributions):**
- Negative Hypergeometric (N, n, k)
 - Hypergeometric (N, K, n)
 - Bernoulli (p)
 - Polya (n, p, β)
 - Discrete Uniform (x_1, \dots, x_n)
- Second Row (Distributions and Functions):**
- Geometric (p)
 - Negative Binomial, Pascal (μ, p)
 - Binomial (n, p)
 - Poisson (λ)
 - Liquid-Spin Uniform (A, λ)
- Third Row (Mathematical Functions and Parameters):**
- $\Gamma(x) = \frac{\Gamma(x-1)}{x-1}$
 - $\Gamma(x) = \frac{\Gamma(x-1)}{x-1}$
 - $\Gamma(x) = \frac{\Gamma(x-1)}{x-1}$
 - $\Gamma(x) = \frac{\Gamma(x-1)}{x-1}$
 - $\Gamma(x) = \frac{\Gamma(x-1)}{x-1}$
- Fourth Row (Distributions and Functions):**
- Normal (μ, σ^2)
 - Standard Normal
 - T (ν)
 - F (ν_1, ν_2)
 - Chi-Squared (ν)
- Fifth Row (Distributions and Functions):**
- Exponential (λ)
 - Lognormal (μ, σ)
 - Standard Cauchy
 - Cauchy (μ, σ)
 - Any-Site
- Sixth Row (Distributions and Functions):**
- Weibull (k, λ)
 - Generalized Gamma (α, β, γ)
 - Gamma (α, β)
 - Beta (α, β)
 - Triangular (a, b, c)
- Seventh Row (Distributions and Functions):**
- Ricefish (λ, μ)
 - Exponential (λ)
 - Exponential (λ)
 - Exponential (λ)
 - Exponential (λ)
- Eighth Row (Distributions and Functions):**
- Double Exponential, Laplace (μ, σ)
 - Standard Logistic
 - Logistic (μ, σ)
 - Loglogistic (μ, σ)
 - Parao (μ, σ)
- Ninth Row (Distributions and Functions):**
- X_1, \dots, X_n iid, $\mu = \gamma, \sigma = \gamma$
 - X_1, \dots, X_n iid, $\mu = \gamma, \sigma = \gamma$
 - X_1, \dots, X_n iid, $\mu = \gamma, \sigma = \gamma$
 - X_1, \dots, X_n iid, $\mu = \gamma, \sigma = \gamma$
 - X_1, \dots, X_n iid, $\mu = \gamma, \sigma = \gamma$
- Bottom Row (Distributions and Functions):**
- X_1, \dots, X_n iid, $\mu = \gamma, \sigma = \gamma$
 - X_1, \dots, X_n iid, $\mu = \gamma, \sigma = \gamma$
 - X_1, \dots, X_n iid, $\mu = \gamma, \sigma = \gamma$
 - X_1, \dots, X_n iid, $\mu = \gamma, \sigma = \gamma$
 - X_1, \dots, X_n iid, $\mu = \gamma, \sigma = \gamma$

[illegible]

上海交通大学
SHANGHAI JIAO TONG UNIVERSITY

- 1 Probability Space
- 2 Random Variables & Distributions
- 3 Expectations
- 4 Common Distributions
- 5 Useful Inequalities**
- 6 Convergence
- 7 Properties of a Random Sample



Markov's Inequality

Let X be a RV. If $\mathbb{P}(X \geq 0) = 1$ and $\mathbb{P}(X = 0) < 1$, then, for any $r > 0$,

$$\mathbb{P}(X \geq r) \leq \frac{\mathbb{E}[X]}{r},$$

with equality if and only if

$$X = \begin{cases} r, & \text{with probability } p, \\ 0, & \text{with probability } 1 - p. \end{cases}$$



Markov's Inequality

Let X be a RV. If $\mathbb{P}(X \geq 0) = 1$ and $\mathbb{P}(X = 0) < 1$, then, for any $r > 0$,

$$\mathbb{P}(X \geq r) \leq \frac{\mathbb{E}[X]}{r},$$

with equality if and only if

$$X = \begin{cases} r, & \text{with probability } p, \\ 0, & \text{with probability } 1 - p. \end{cases}$$

- Markov's Inequality has many variations, which are usually called Chebyshev's Inequality.



Chebyshev's Inequality

Let X be a RV and $g(x)$ be a nonnegative function. Then, for any $r > 0$,

$$\mathbb{P}(g(X) \geq r) \leq \frac{\mathbb{E}[g(X)]}{r}.$$

Chebyshev's Inequality

Let X be a RV and $g(x)$ be a nonnegative function. Then, for any $r > 0$,

$$\mathbb{P}(g(X) \geq r) \leq \frac{\mathbb{E}[g(X)]}{r}.$$

Chebyshev's Inequality

Let X be a RV. Then, for any $r, p > 0$,

$$\mathbb{P}(|X| \geq r) \leq \frac{\mathbb{E}[|X|^p]}{r^p},$$

$$\mathbb{P}(|X - \mu| \geq r) \leq \frac{\sigma^2}{r^2},$$

where $\mu := \mathbb{E}[X]$, and $\sigma^2 := \text{Var}(X)$.

- Chebyshev's Inequality is typically very conservative.

- Chebyshev's Inequality is typically very conservative.
- If $Z \sim \mathcal{N}(0, 1)$, a tighter bound is available: For any $t > 0$,

$$2\Phi(-t) = \mathbb{P}(|Z| \geq t) \leq \sqrt{\frac{2}{\pi}} \frac{1}{t} e^{-t^2/2},$$


$$2\Phi(-t) = \mathbb{P}(|Z| \geq t) \geq \sqrt{\frac{2}{\pi}} \frac{t}{1+t^2} e^{-t^2/2}.$$



- Chebyshev's Inequality is typically very conservative.
- If $Z \sim \mathcal{N}(0, 1)$, a tighter bound is available: For any $t > 0$,

$$2\Phi(-t) = \mathbb{P}(|Z| \geq t) \leq \sqrt{\frac{2}{\pi}} \frac{1}{t} e^{-t^2/2},$$

$$2\Phi(-t) = \mathbb{P}(|Z| \geq t) \geq \sqrt{\frac{2}{\pi}} \frac{t}{1+t^2} e^{-t^2/2}.$$



- A function $g(x)$ is **convex** if

$$g(\lambda x + (1 - \lambda)y) \leq \lambda g(x) + (1 - \lambda)g(y),$$

for all x and y , and $\lambda \in (0, 1)$.



- A function $g(x)$ is **convex** if

$$g(\lambda x + (1 - \lambda)y) \leq \lambda g(x) + (1 - \lambda)g(y),$$

for all x and y , and $\lambda \in (0, 1)$.

- A function $g(x)$ is concave if $-g(x)$ is convex.



- A function $g(x)$ is **convex** if

$$g(\lambda x + (1 - \lambda)y) \leq \lambda g(x) + (1 - \lambda)g(y),$$

for all x and y , and $\lambda \in (0, 1)$.

- A function $g(x)$ is concave if $-g(x)$ is convex.

Jensen's Inequality

Let X be a RV. If $g(x)$ is a convex function, then

$$\mathbb{E}[g(X)] \geq g(\mathbb{E}[X]),$$

with equality if and only if $g(x)$ is a linear function on some set A such that $\mathbb{P}(X \in A) = 1$.



Hölder's Inequality

Let X and Y be any two RVs, and let p and q be any two positive numbers (necessarily greater than 1) satisfying

$$\frac{1}{p} + \frac{1}{q} = 1.$$

Then,

$$|\mathbb{E}[XY]| \leq \mathbb{E}[|XY|] \leq \{\mathbb{E}[|X|^p]\}^{1/p} \{\mathbb{E}[|Y|^q]\}^{1/q}.$$



Cauchy-Schwarz Inequality ($p = q = 2$)

Let X and Y be any two RVs, then

$$|\mathbb{E}[XY]| \leq \mathbb{E}[|XY|] \leq \{\mathbb{E}[|X|^2]\}^{1/2} \{\mathbb{E}[|Y|^2]\}^{1/2}.$$



Cauchy-Schwarz Inequality ($p = q = 2$)

Let X and Y be any two RVs, then

$$|\mathbb{E}[XY]| \leq \mathbb{E}[|XY|] \leq \{\mathbb{E}[|X|^2]\}^{1/2} \{\mathbb{E}[|Y|^2]\}^{1/2}.$$

Liapounov's Inequality ($Y \equiv 1$)

Let X be a RV, then for any $s > r > 1$,

$$\{\mathbb{E}[|X|^r]\}^{1/r} \leq \{\mathbb{E}[|X|^s]\}^{1/s}.$$



Minkowski's Inequality

Let X and Y be any two RVs. Then, for $p \geq 1$,

$$\{\mathbb{E}[|X + Y|^p]\}^{1/p} \leq \{\mathbb{E}[|X|^p]\}^{1/p} + \{\mathbb{E}[|Y|^p]\}^{1/p}.$$



Minkowski's Inequality

Let X and Y be any two RVs. Then, for $p \geq 1$,

$$\{\mathbb{E}[|X + Y|^p]\}^{1/p} \leq \{\mathbb{E}[|X|^p]\}^{1/p} + \{\mathbb{E}[|Y|^p]\}^{1/p}.$$

- **Remark:** The preceding Hölder's Inequality (including its special cases) and Minkowski's Inequality also apply to numerical sums where there is no explicit reference to an expectation.

- 1 Probability Space
- 2 Random Variables & Distributions
- 3 Expectations
- 4 Common Distributions
- 5 Useful Inequalities
- 6 Convergence**
- 7 Properties of a Random Sample



Consider a sequence of RVs $\{X_n : n \geq 1\}$ and another RV X .

Consider a sequence of RVs $\{X_n : n \geq 1\}$ and another RV X .

- **Convergence Almost Surely** (a.s.), $X_n \xrightarrow{a.s.} X$:

$$\mathbb{P}(\lim_{n \rightarrow \infty} X_n = X) = 1.$$

Consider a sequence of RVs $\{X_n : n \geq 1\}$ and another RV X .

- **Convergence Almost Surely** (a.s.), $X_n \xrightarrow{a.s.} X$:

$$\mathbb{P}(\lim_{n \rightarrow \infty} X_n = X) = 1.$$

- **Convergence in Probability**, $X_n \xrightarrow{p} X$:

$$\lim_{n \rightarrow \infty} \mathbb{P}(|X_n - X| > \epsilon) = 0, \text{ for any } \epsilon > 0.$$



Consider a sequence of RVs $\{X_n : n \geq 1\}$ and another RV X .

- **Convergence Almost Surely** (a.s.), $X_n \xrightarrow{a.s.} X$:

$$\mathbb{P}(\lim_{n \rightarrow \infty} X_n = X) = 1.$$

- **Convergence in Probability**, $X_n \xrightarrow{p} X$:

$$\lim_{n \rightarrow \infty} \mathbb{P}(|X_n - X| > \epsilon) = 0, \text{ for any } \epsilon > 0.$$

- **Convergence in Distribution**, $X_n \xrightarrow{d} X$ or $X_n \Rightarrow X$:

$$\lim_{n \rightarrow \infty} F_n(x) = F(x), \text{ for any continuous point } x \text{ of } F(x),$$

where F_n and F are CDF of X_n and X , respectively.



Consider a sequence of RVs $\{X_n : n \geq 1\}$ and another RV X .

- **Convergence Almost Surely** (a.s.), $X_n \xrightarrow{a.s.} X$:

$$\mathbb{P}(\lim_{n \rightarrow \infty} X_n = X) = 1.$$

- **Convergence in Probability**, $X_n \xrightarrow{p} X$:

$$\lim_{n \rightarrow \infty} \mathbb{P}(|X_n - X| > \epsilon) = 0, \text{ for any } \epsilon > 0.$$

- **Convergence in Distribution**, $X_n \xrightarrow{d} X$ or $X_n \Rightarrow X$:

$$\lim_{n \rightarrow \infty} F_n(x) = F(x), \text{ for any continuous point } x \text{ of } F(x),$$

where F_n and F are CDF of X_n and X , respectively.

- **Convergence in L^r Norm** ($r \in [1, \infty)$), $X_n \xrightarrow{L^r} X$:

$$\lim_{n \rightarrow \infty} \mathbb{E}(|X_n - X|^r) = 0,$$

given $\mathbb{E}[|X_n|^r] < \infty$ for any $n \geq 1$ and $\mathbb{E}[|X|^r] < \infty$.



- Simple relationships:

$$\begin{array}{ccccc}
 X_n \xrightarrow{a.s.} X & \implies & X_n \xrightarrow{p} X & \implies & X_n \Rightarrow X \\
 & & \uparrow & & \\
 X_n \xrightarrow{L^s} X & \xRightarrow{s > r \geq 1} & X_n \xrightarrow{L^r} X & \implies & \mathbb{E}[|X_n|^r] \rightarrow \mathbb{E}[|X|^r]
 \end{array}$$



- Simple relationships:

$$\begin{array}{ccccc}
 X_n \xrightarrow{a.s.} X & \implies & X_n \xrightarrow{p} X & \implies & X_n \Rightarrow X \\
 & & \uparrow & & \\
 X_n \xrightarrow{L^s} X & \xRightarrow{s > r \geq 1} & X_n \xrightarrow{L^r} X & \implies & \mathbb{E}[|X_n|^r] \rightarrow \mathbb{E}[|X|^r]
 \end{array}$$

- $X_n \Rightarrow \text{a constant } c \implies X_n \xrightarrow{p} c.$



- Simple relationships:

$$\begin{array}{ccccc}
 X_n \xrightarrow{a.s.} X & \implies & X_n \xrightarrow{p} X & \implies & X_n \Rightarrow X \\
 & & \uparrow & & \\
 X_n \xrightarrow{L^s} X & \xRightarrow{s > r \geq 1} & X_n \xrightarrow{L^r} X & \implies & \mathbb{E}[|X_n|^r] \rightarrow \mathbb{E}[|X|^r]
 \end{array}$$

- $X_n \Rightarrow \text{a constant } c \implies X_n \xrightarrow{p} c.$
- $X_n \xrightarrow{L^1} X \implies \mathbb{E}[X_n] \rightarrow \mathbb{E}[X].$



- Simple relationships:

$$\begin{array}{ccccc}
 X_n \xrightarrow{a.s.} X & \implies & X_n \xrightarrow{p} X & \implies & X_n \Rightarrow X \\
 & & \uparrow & & \\
 X_n \xrightarrow{L^s} X & \xRightarrow{s > r \geq 1} & X_n \xrightarrow{L^r} X & \implies & \mathbb{E}[|X_n|^r] \rightarrow \mathbb{E}[|X|^r]
 \end{array}$$

- $X_n \Rightarrow \text{a constant } c \implies X_n \xrightarrow{p} c.$
- $X_n \xrightarrow{L^1} X \implies \mathbb{E}[X_n] \rightarrow \mathbb{E}[X].$
- $X_n \xrightarrow{a.s.} X \iff \sup_{j \geq n} |X_j - X| \xrightarrow{p} 0.$



- Simple relationships:

$$\begin{array}{ccccc}
 X_n \xrightarrow{a.s.} X & \implies & X_n \xrightarrow{p} X & \implies & X_n \Rightarrow X \\
 & & \uparrow & & \\
 X_n \xrightarrow{L^s} X & \xRightarrow{s > r \geq 1} & X_n \xrightarrow{L^r} X & \implies & \mathbb{E}[|X_n|^r] \rightarrow \mathbb{E}[|X|^r]
 \end{array}$$

- $X_n \Rightarrow \text{a constant } c \implies X_n \xrightarrow{p} c.$
- $X_n \xrightarrow{L^1} X \implies \mathbb{E}[X_n] \rightarrow \mathbb{E}[X].$
- $X_n \xrightarrow{a.s.} X \iff \sup_{j \geq n} |X_j - X| \xrightarrow{p} 0.$
- $X_n \xrightarrow{p} X \iff \text{For every subsequence } X_n(m) \text{ there is a further subsequence } X_n(m_k) \text{ such that } X_n(m_k) \xrightarrow{a.s.} X.$

- Question: If $X_n \Rightarrow X$ or $X_n \xrightarrow{p} X$ or $X_n \xrightarrow{a.s.} X$, does it imply $\mathbb{E}[X_n] \rightarrow \mathbb{E}[X]$?

- Question: If $X_n \Rightarrow X$ or $X_n \xrightarrow{p} X$ or $X_n \xrightarrow{a.s.} X$, does it imply $\mathbb{E}[X_n] \rightarrow \mathbb{E}[X]$?

Fatou's Lemma

Suppose $X_n \geq Y$ a.s. for all n where $\mathbb{E}[|Y|] < \infty$. Then $\mathbb{E}[\liminf_{n \rightarrow \infty} X_n] \leq \liminf_{n \rightarrow \infty} \mathbb{E}[X_n]$. In particular, if $X_n \geq 0$ a.s. for all n , then the result holds.



- Question: If $X_n \Rightarrow X$ or $X_n \xrightarrow{p} X$ or $X_n \xrightarrow{a.s.} X$, does it imply $\mathbb{E}[X_n] \rightarrow \mathbb{E}[X]$?

Fatou's Lemma

Suppose $X_n \geq Y$ a.s. for all n where $\mathbb{E}[|Y|] < \infty$. Then $\mathbb{E}[\liminf_{n \rightarrow \infty} X_n] \leq \liminf_{n \rightarrow \infty} \mathbb{E}[X_n]$. In particular, if $X_n \geq 0$ a.s. for all n , then the result holds.

Monotone Convergence Theorem (MCT)

Suppose $X_n \xrightarrow{a.s.} X$, and $0 \leq X_1 \leq X_2 \leq \dots$ a.s.. Then $\mathbb{E}[X_n] \rightarrow \mathbb{E}[X]$.



Dominated Convergence Theorem (DCT)

Suppose $X_n \xrightarrow{a.s.} X$, $|X_n| \leq Y$ a.s. for all n , and $\mathbb{E}[|Y|] < \infty$. Then $\mathbb{E}[X_n] \rightarrow \mathbb{E}[X]$.



Dominated Convergence Theorem (DCT)

Suppose $X_n \xrightarrow{a.s.} X$, $|X_n| \leq Y$ a.s. for all n , and $\mathbb{E}[|Y|] < \infty$. Then $\mathbb{E}[X_n] \rightarrow \mathbb{E}[X]$.

- The DCT is still true if $\xrightarrow{a.s.}$ is replaced by \xrightarrow{p} .

Dominated Convergence Theorem (DCT)

Suppose $X_n \xrightarrow{a.s.} X$, $|X_n| \leq Y$ a.s. for all n , and $\mathbb{E}[|Y|] < \infty$. Then $\mathbb{E}[X_n] \rightarrow \mathbb{E}[X]$.

- The DCT is still true if $\xrightarrow{a.s.}$ is replaced by \xrightarrow{p} .
- An **even more general** result:
Suppose $X_n \xrightarrow{p} X$, $|X_n| \leq Y$ a.s. for all n , and $\mathbb{E}[|Y|^r] < \infty$ with $r \geq 1$. Then, $\mathbb{E}[|X_n|^r] < \infty$, $\mathbb{E}[|X|] < \infty$, and $X_n \xrightarrow{L^r} X$.

- If $X_n \xrightarrow{a.s.} X$ and $Y_n \xrightarrow{a.s.} Y$, then
 - ① $X = Y$ a.s.;
 - ② Random vector $(X_n, Y_n)^\top \xrightarrow{a.s.} (X, Y)^\top$.



- If $X_n \xrightarrow{a.s.} X$ and $Y_n \xrightarrow{a.s.} Y$, then
 - ① $X = Y$ a.s.;
 - ② Random vector $(X_n, Y_n)^\top \xrightarrow{a.s.} (X, Y)^\top$.
 $\implies aX_n + bY_n \xrightarrow{a.s.} aX + bY$; $X_n Y_n \xrightarrow{a.s.} XY$.



- If $X_n \xrightarrow{a.s.} X$ and $Y_n \xrightarrow{a.s.} Y$, then
 - ① $X = Y$ a.s.;
 - ② Random vector $(X_n, Y_n)^\top \xrightarrow{a.s.} (X, Y)^\top$.
 $\implies aX_n + bY_n \xrightarrow{a.s.} aX + bY; X_n Y_n \xrightarrow{a.s.} XY$.
- If $X_n \xrightarrow{p} X$ and $Y_n \xrightarrow{p} Y$, then
 - ① $X = Y$ a.s.;
 - ② Random vector $(X_n, Y_n)^\top \xrightarrow{p} (X, Y)^\top$.
 $\implies aX_n + bY_n \xrightarrow{p} aX + bY; X_n Y_n \xrightarrow{p} XY$.



- If $X_n \xrightarrow{a.s.} X$ and $Y_n \xrightarrow{a.s.} Y$, then
 - ① $X = Y$ a.s.;
 - ② Random vector $(X_n, Y_n)^\top \xrightarrow{a.s.} (X, Y)^\top$.
 $\implies aX_n + bY_n \xrightarrow{a.s.} aX + bY$; $X_n Y_n \xrightarrow{a.s.} XY$.
- If $X_n \xrightarrow{p} X$ and $Y_n \xrightarrow{p} Y$, then
 - ① $X = Y$ a.s.;
 - ② Random vector $(X_n, Y_n)^\top \xrightarrow{p} (X, Y)^\top$.
 $\implies aX_n + bY_n \xrightarrow{p} aX + bY$; $X_n Y_n \xrightarrow{p} XY$.
- If $X_n \xrightarrow{L^r} X$ and $Y_n \xrightarrow{L^r} Y$, then
 - ① $X = Y$ a.s.;
 - ② $aX_n + bY_n \xrightarrow{L^r} aX + bY$.



- If $X_n \xrightarrow{a.s.} X$ and $Y_n \xrightarrow{a.s.} Y$, then
 - ① $X = Y$ a.s.;
 - ② Random vector $(X_n, Y_n)^\top \xrightarrow{a.s.} (X, Y)^\top$.
 $\implies aX_n + bY_n \xrightarrow{a.s.} aX + bY$; $X_n Y_n \xrightarrow{a.s.} XY$.
- If $X_n \xrightarrow{p} X$ and $Y_n \xrightarrow{p} Y$, then
 - ① $X = Y$ a.s.;
 - ② Random vector $(X_n, Y_n)^\top \xrightarrow{p} (X, Y)^\top$.
 $\implies aX_n + bY_n \xrightarrow{p} aX + bY$; $X_n Y_n \xrightarrow{p} XY$.
- If $X_n \xrightarrow{L^r} X$ and $Y_n \xrightarrow{L^r} Y$, then
 - ① $X = Y$ a.s.;
 - ② $aX_n + bY_n \xrightarrow{L^r} aX + bY$.
- None of the above are true for convergence in distribution.



- If $X_n \xrightarrow{a.s.} X$ and $Y_n \xrightarrow{a.s.} Y$, then
 - ① $X = Y$ a.s.;
 - ② Random vector $(X_n, Y_n)^\top \xrightarrow{a.s.} (X, Y)^\top$.
 $\implies aX_n + bY_n \xrightarrow{a.s.} aX + bY$; $X_n Y_n \xrightarrow{a.s.} XY$.
- If $X_n \xrightarrow{p} X$ and $Y_n \xrightarrow{p} Y$, then
 - ① $X = Y$ a.s.;
 - ② Random vector $(X_n, Y_n)^\top \xrightarrow{p} (X, Y)^\top$.
 $\implies aX_n + bY_n \xrightarrow{p} aX + bY$; $X_n Y_n \xrightarrow{p} XY$.
- If $X_n \xrightarrow{L^r} X$ and $Y_n \xrightarrow{L^r} Y$, then
 - ① $X = Y$ a.s.;
 - ② $aX_n + bY_n \xrightarrow{L^r} aX + bY$.
- None of the above are true for convergence in distribution.
- If $X_n \Rightarrow X$ and $Y_n \Rightarrow$ a constant c , then
 - Random vector $(X_n, Y_n)^\top \Rightarrow (X, c)^\top$.
 $\implies aX_n + bY_n \Rightarrow aX + bc$; $X_n Y_n \Rightarrow cX$. (Slutsky's theorem)

Continuous Mapping Theorem

Consider a sequence of RVs $\{X_n : n \geq 1\}$ and another RV X . Suppose g is a function that has the set of discontinuity points D such that $\mathbb{P}(X \in D) = 0$. Then,

$$X_n \xrightarrow{a.s.} X \implies g(X_n) \xrightarrow{a.s.} g(X);$$

$$X_n \xrightarrow{p} X \implies g(X_n) \xrightarrow{p} g(X);$$

$$X_n \Rightarrow X \implies g(X_n) \Rightarrow g(X).$$



- 1 Probability Space
- 2 Random Variables & Distributions
- 3 Expectations
- 4 Common Distributions
- 5 Useful Inequalities
- 6 Convergence
- 7 Properties of a Random Sample**

Properties of a Random Sample

- Let X_1, \dots, X_n be a **random sample** from a distribution with mean μ and variance σ^2 , i.e., X_1, \dots, X_n are iid, and $\mathbb{E}[X_i] = \mu$ and $\text{Var}(X_i) = \sigma^2$, $i = 1, \dots, n$.

Properties of a Random Sample

- Let X_1, \dots, X_n be a **random sample** from a distribution with mean μ and variance σ^2 , i.e., X_1, \dots, X_n are iid, and $\mathbb{E}[X_i] = \mu$ and $\text{Var}(X_i) = \sigma^2$, $i = 1, \dots, n$.

- Define

$$\bar{X} := \frac{1}{n} \sum_{i=1}^n X_i, \text{ and } S^2 := \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}.$$



Properties of a Random Sample

- Let X_1, \dots, X_n be a **random sample** from a distribution with mean μ and variance σ^2 , i.e., X_1, \dots, X_n are iid, and $\mathbb{E}[X_i] = \mu$ and $\text{Var}(X_i) = \sigma^2$, $i = 1, \dots, n$.

- Define

$$\bar{X} := \frac{1}{n} \sum_{i=1}^n X_i, \text{ and } S^2 := \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}.$$

- For a **general** distribution, the following is true:



Properties of a Random Sample

- Let X_1, \dots, X_n be a **random sample** from a distribution with mean μ and variance σ^2 , i.e., X_1, \dots, X_n are iid, and $\mathbb{E}[X_i] = \mu$ and $\text{Var}(X_i) = \sigma^2$, $i = 1, \dots, n$.

- Define

$$\bar{X} := \frac{1}{n} \sum_{i=1}^n X_i, \text{ and } S^2 := \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}.$$

- For a **general** distribution, the following is true:
 - 1 \bar{X} is an **unbiased** estimator of μ , i.e., $\mathbb{E}[\bar{X}] = \mu$;



Properties of a Random Sample

- Let X_1, \dots, X_n be a **random sample** from a distribution with mean μ and variance σ^2 , i.e., X_1, \dots, X_n are iid, and $\mathbb{E}[X_i] = \mu$ and $\text{Var}(X_i) = \sigma^2$, $i = 1, \dots, n$.

- Define

$$\bar{X} := \frac{1}{n} \sum_{i=1}^n X_i, \text{ and } S^2 := \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}.$$

- For a **general** distribution, the following is true:

- ① \bar{X} is an **unbiased** estimator of μ , i.e., $\mathbb{E}[\bar{X}] = \mu$;
- ② S^2 is an **unbiased** estimator of σ^2 , i.e., $\mathbb{E}[S^2] = \sigma^2$;



Properties of a Random Sample

- Let X_1, \dots, X_n be a **random sample** from a distribution with mean μ and variance σ^2 , i.e., X_1, \dots, X_n are iid, and $\mathbb{E}[X_i] = \mu$ and $\text{Var}(X_i) = \sigma^2$, $i = 1, \dots, n$.

- Define

$$\bar{X} := \frac{1}{n} \sum_{i=1}^n X_i, \text{ and } S^2 := \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}.$$

- For a **general** distribution, the following is true:

- ① \bar{X} is an **unbiased** estimator of μ , i.e., $\mathbb{E}[\bar{X}] = \mu$;
- ② S^2 is an **unbiased** estimator of σ^2 , i.e., $\mathbb{E}[S^2] = \sigma^2$;
- ③ $\text{Var}(\bar{X}) = \sigma^2/n$.



Properties of a Random Sample

- Let X_1, \dots, X_n be a **random sample** from a distribution with mean μ and variance σ^2 , i.e., X_1, \dots, X_n are iid, and $\mathbb{E}[X_i] = \mu$ and $\text{Var}(X_i) = \sigma^2$, $i = 1, \dots, n$.

- Define

$$\bar{X} := \frac{1}{n} \sum_{i=1}^n X_i, \text{ and } S^2 := \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}.$$

- For a **general** distribution, the following is true:

- ① \bar{X} is an **unbiased** estimator of μ , i.e., $\mathbb{E}[\bar{X}] = \mu$;
- ② S^2 is an **unbiased** estimator of σ^2 , i.e., $\mathbb{E}[S^2] = \sigma^2$;
- ③ $\text{Var}(\bar{X}) = \sigma^2/n$.

- If the distribution is $\mathcal{N}(\mu, \sigma^2)$, we further have:



Properties of a Random Sample

- Let X_1, \dots, X_n be a **random sample** from a distribution with mean μ and variance σ^2 , i.e., X_1, \dots, X_n are iid, and $\mathbb{E}[X_i] = \mu$ and $\text{Var}(X_i) = \sigma^2$, $i = 1, \dots, n$.

- Define

$$\bar{X} := \frac{1}{n} \sum_{i=1}^n X_i, \text{ and } S^2 := \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}.$$

- For a **general** distribution, the following is true:

- ① \bar{X} is an **unbiased** estimator of μ , i.e., $\mathbb{E}[\bar{X}] = \mu$;
- ② S^2 is an **unbiased** estimator of σ^2 , i.e., $\mathbb{E}[S^2] = \sigma^2$;
- ③ $\text{Var}(\bar{X}) = \sigma^2/n$.

- If the distribution is $\mathcal{N}(\mu, \sigma^2)$, we further have:

- ④ $\bar{X} \sim \mathcal{N}(\mu, \sigma^2/n)$, i.e., $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim \mathcal{N}(0, 1)$;



Properties of a Random Sample

- Let X_1, \dots, X_n be a **random sample** from a distribution with mean μ and variance σ^2 , i.e., X_1, \dots, X_n are iid, and $\mathbb{E}[X_i] = \mu$ and $\text{Var}(X_i) = \sigma^2$, $i = 1, \dots, n$.

- Define

$$\bar{X} := \frac{1}{n} \sum_{i=1}^n X_i, \text{ and } S^2 := \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}.$$

- For a **general** distribution, the following is true:

- ① \bar{X} is an **unbiased** estimator of μ , i.e., $\mathbb{E}[\bar{X}] = \mu$;
- ② S^2 is an **unbiased** estimator of σ^2 , i.e., $\mathbb{E}[S^2] = \sigma^2$;
- ③ $\text{Var}(\bar{X}) = \sigma^2/n$.

- If the distribution is $\mathcal{N}(\mu, \sigma^2)$, we further have:

- ④ $\bar{X} \sim \mathcal{N}(\mu, \sigma^2/n)$, i.e., $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim \mathcal{N}(0, 1)$;
- ⑤ $\bar{X} \perp S^2$;

Properties of a Random Sample

- Let X_1, \dots, X_n be a **random sample** from a distribution with mean μ and variance σ^2 , i.e., X_1, \dots, X_n are iid, and $\mathbb{E}[X_i] = \mu$ and $\text{Var}(X_i) = \sigma^2$, $i = 1, \dots, n$.

- Define

$$\bar{X} := \frac{1}{n} \sum_{i=1}^n X_i, \text{ and } S^2 := \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}.$$

- For a **general** distribution, the following is true:

- ① \bar{X} is an **unbiased** estimator of μ , i.e., $\mathbb{E}[\bar{X}] = \mu$;
- ② S^2 is an **unbiased** estimator of σ^2 , i.e., $\mathbb{E}[S^2] = \sigma^2$;
- ③ $\text{Var}(\bar{X}) = \sigma^2/n$.

- If the distribution is $\mathcal{N}(\mu, \sigma^2)$, we further have:

- ④ $\bar{X} \sim \mathcal{N}(\mu, \sigma^2/n)$, i.e., $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim \mathcal{N}(0, 1)$;
- ⑤ $\bar{X} \perp S^2$;
- ⑥ $(n-1)S^2/\sigma^2 \sim \chi_{n-1}^2$;

Properties of a Random Sample

- Let X_1, \dots, X_n be a **random sample** from a distribution with mean μ and variance σ^2 , i.e., X_1, \dots, X_n are iid, and $\mathbb{E}[X_i] = \mu$ and $\text{Var}(X_i) = \sigma^2$, $i = 1, \dots, n$.

- Define

$$\bar{X} := \frac{1}{n} \sum_{i=1}^n X_i, \text{ and } S^2 := \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}.$$

- For a **general** distribution, the following is true:

- ① \bar{X} is an **unbiased** estimator of μ , i.e., $\mathbb{E}[\bar{X}] = \mu$;
- ② S^2 is an **unbiased** estimator of σ^2 , i.e., $\mathbb{E}[S^2] = \sigma^2$;
- ③ $\text{Var}(\bar{X}) = \sigma^2/n$.

- If the distribution is $\mathcal{N}(\mu, \sigma^2)$, we further have:

- ④ $\bar{X} \sim \mathcal{N}(\mu, \sigma^2/n)$, i.e., $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim \mathcal{N}(0, 1)$;
- ⑤ $\bar{X} \perp S^2$;
- ⑥ $(n-1)S^2/\sigma^2 \sim \chi_{n-1}^2$;
- ⑦ $\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t_{n-1}$.



- For a **general** distribution, *what can we say about the distribution of \bar{X} ?*

- For a **general** distribution, *what can we say about the distribution of \bar{X}* ?
- $\text{Var}(\bar{X}) = \sigma^2/n$ intuitively means that the randomness of \bar{X} vanishes and \bar{X} concentrates around μ when n gets large.



- For a **general** distribution, *what can we say about the distribution of \bar{X}* ?
- $\text{Var}(\bar{X}) = \sigma^2/n$ intuitively means that the randomness of \bar{X} vanishes and \bar{X} concentrates around μ when n gets large.
- Denote \bar{X} as \bar{X}_n , to explicitly indicate the effect of **sample size** n .



- For a **general** distribution, *what can we say about the distribution of \bar{X} ?*
- $\text{Var}(\bar{X}) = \sigma^2/n$ intuitively means that the randomness of \bar{X} vanishes and \bar{X} concentrates around μ when n gets large.
- Denote \bar{X} as \bar{X}_n , to explicitly indicate the effect of **sample size** n .

Weak Law of Large Numbers (WLLN)

Suppose X_1, \dots, X_n are iid with mean μ and variance $\sigma^2 < \infty$.[†] Then, $\bar{X}_n \xrightarrow{p} \mu$.

[†] Mutual independence can be weakened to pairwise independence; $\sigma^2 < \infty$ can be weakened to $\mathbb{E}[|X_i|] \leq \infty$.



- For a **general** distribution, *what can we say about the distribution of \bar{X} ?*
- $\text{Var}(\bar{X}) = \sigma^2/n$ intuitively means that the randomness of \bar{X} vanishes and \bar{X} concentrates around μ when n gets large.
- Denote \bar{X} as \bar{X}_n , to explicitly indicate the effect of **sample size** n .

Weak Law of Large Numbers (WLLN)

Suppose X_1, \dots, X_n are iid with mean μ and variance $\sigma^2 < \infty$.[†] Then, $\bar{X}_n \xrightarrow{p} \mu$.

Strong Law of Large Numbers (SLLN)

Suppose X_1, \dots, X_n are iid with mean μ and variance $\sigma^2 < \infty$.[†] Then, $\bar{X}_n \xrightarrow{a.s.} \mu$.

[†] Mutual independence can be weakened to pairwise independence; $\sigma^2 < \infty$ can be weakened to $\mathbb{E}[|X_i|] \leq \infty$.

- Note that for **normal** distribution, $\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \sim \mathcal{N}(0, 1)$, regardless of the value of n .
- For a **general** distribution, *what can we say about the distribution of $\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}}$?*



- Note that for **normal** distribution, $\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \sim \mathcal{N}(0, 1)$, regardless of the value of n .
- For a **general** distribution, *what can we say about the distribution of $\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}}$?*
- Note that $\mathbb{E} \left[\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \right] = 0$ and $\text{Var} \left(\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \right) = 1$, regardless of the distribution and the value of n .

- Note that for **normal** distribution, $\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \sim \mathcal{N}(0, 1)$, regardless of the value of n .
- For a **general** distribution, *what can we say about the distribution of $\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}}$?*
- Note that $\mathbb{E} \left[\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \right] = 0$ and $\text{Var} \left(\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \right) = 1$, regardless of the distribution and the value of n .

Central Limit Theorem (CLT)

Suppose X_1, \dots, X_n are iid with mean μ and variance $\sigma^2 \in (0, \infty)$. Then,

$$\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \Rightarrow \mathcal{N}(0, 1).$$