

Calibration and PnP Solutions

Lecturer: Martial Hebert

TA: Xiaofang Wang & Xinshuo Weng Scribes: Tianyi Shen¹ & Xueqiang Wang²

1 Recap




Figure 1: Affine planar rectification

Given a projective version of the world scene, as shown in Fig. 1, we may find a homography H_a that preserves the parallelism and thus recovers affine properties as in the original plane. So $H_a H$ is an affine transformation. Actually, H_a can be computed if we know the projection of the line at infinity. In this case, we may denote $l_\infty = [l_1 \ l_2 \ l_3]^T$. H_a can be expressed as

$$H_a = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ l_1 & l_2 & l_3 \end{bmatrix}$$

2 Metric rectification

2.1 For 2D plane




Figure 2: Metric planar rectification

¹ Tianyi Shen, tshen1 ² Xueqiang Wang, xueqianw

For metric rectification, the euclidean properties of the original plane can be recovered if we find a homography H_m , which preserves the angle between two lines on the plane. We first denote two lines $l = [a \ b \ c]^\top$ and $l' = [a' \ b' \ c']^\top$. The angle between these two lines is

$$\cos \theta = \frac{aa' + bb'}{\sqrt{a^2 + b^2} \sqrt{a'^2 + b'^2}}$$

Given $aa' + bb' = l' TC^* l$ and $C^* = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$, we may derive that

$$\cos \theta = \frac{l' TC^* l}{\sqrt{l^\top C^* l} \sqrt{l'^\top C^* l'}}$$

If l' and l are orthogonal, we have

$$l' TC^* l = 0$$

In Fig. 2, assume we have a line l_0 in the original plane and its homography transformation l . After metric rectification, the line became l' and right angle preserved. We get the following equation

$$l' = H_m^{-\top} H^{-\top} l_0 = H^{-\top} l \quad (1)$$

$$l_0^\top C^* l_0^j = l_0^\top H_m^{-1} C^* H_m^{-\top} l_0^j = 0 \quad (2)$$

To solve for H_m , we may need at least 4 pairs of lines.

2.2 For 3D space

Assume that we have two perpendicular planes π_1 and π_2 in 3D space and π_∞ at infinity. Ω_∞ is an absolute conic on π_∞ . The dual of the absolute conic Ω_∞ is called the absolute dual quadric, denoted Ω_∞^*

$$\Omega_\infty^* = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad (3)$$

Similarly, the properties are the same as in the plane. The angle between planes is

$$\cos \theta = \frac{\pi_1^\top \Omega_\infty^* \pi_2}{\sqrt{\pi_1^\top \Omega_\infty^* \pi_1} \sqrt{\pi_2^\top \Omega_\infty^* \pi_2}} \quad (4)$$

Note that C_∞^* is invariant if and only if H is a similarity transformation.

3 Calibration

3.1 Calibration from 3D points (PnP)

For the camera matrix, $M = K[R \ t]$. We want to solve this intrinsic matrix K from calibration.




Figure 3: Calibration from plane homographies: The image of the absolute conic

Given a plane at infinity Π_∞ as in Fig. 3, the absolute conic Ω_∞ is the conic of equation

$$x^2 + y^2 + z^2 = 0, \quad w = 0 \quad (5)$$

where $P = [x \ y \ z \ 0]^\top$. The point p at the image plane can be expressed as

$$\begin{aligned} p &= k [R \ t] \begin{bmatrix} x \\ y \\ z \\ 0 \end{bmatrix} = KR \begin{bmatrix} x \\ y \\ z \end{bmatrix} \\ \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} &= R^\top K^{-1} p \end{aligned} \quad (6)$$

From Eqns. 5 and 6, we have

$$\begin{aligned} [x \ y \ z] \begin{bmatrix} x \\ y \\ z \end{bmatrix} &= 0 \\ p^\top K^{-\top} R R^\top K^{-1} p &= 0 \\ p^\top K^{-\top} K^{-1} p &= 0 \\ p^\top \omega_\infty p &= 0 \end{aligned} \quad (7)$$

Therefore, ω_∞ can be represented by the matrix $K^{-\top} K^{-1}$.

3.2 Outline of the calibration solution




Figure 4: Point I, J at plane Π_∞

Now we assume that a plane Π intersects the conic Ω_∞ at point $I, J = [x \ y \ 0]^\top$. Since $x^2 + y^2 = 0$ does not have a real solution. I, J can be expressed as

$$I = \begin{bmatrix} 1 \\ i \\ 0 \end{bmatrix} \quad J = \begin{bmatrix} i \\ -1 \\ 0 \end{bmatrix} \quad (8)$$

Since $I \in \Omega_\infty$ and $HI \in \omega_\infty$, we know

$$\begin{aligned} I^\top H^\top \omega_\infty H I &= 0 \\ J^\top H^\top \omega_\infty H J &= 0 \end{aligned}$$

Thus we have 2 equations per plane. Note that ω_∞ has 5 dof since a 3×3 matrix gives 6 dof while it is also up to scale.

4 PnP solutions

Suppose we have 3 world coordinates of point i : $p_i^w = [x_i^w \ y_i^w \ z_i^w]^\top$, and 3 coordinates of point i with respect to camera coordinate system: $p_i^c = [x_i^c \ y_i^c \ z_i^c]^\top$.

Now let $M = K [R \ t]$, in this case K is a known variable. Since we have $p = MP$, $M = \begin{bmatrix} \mathbf{m}_1^\top \\ \mathbf{m}_2^\top \\ \mathbf{m}_3^\top \end{bmatrix}$,

$$\begin{aligned} x &= \frac{\mathbf{m}_1^\top P}{\mathbf{m}_3^\top P}, \quad y = \frac{\mathbf{m}_2^\top P}{\mathbf{m}_3^\top P} \\ \Rightarrow \begin{cases} P_i^\top \mathbf{m}_1 - x P_i^\top \mathbf{m}_3 = 0 \\ P_i^\top \mathbf{m}_2 - y P_i^\top \mathbf{m}_3 = 0 \end{cases} \end{aligned} \quad (9)$$

Suppose we have N sets of Eqn. 9, we may write in the matrix form

$$\begin{bmatrix} P_i^\top & 0 & -x P_i^\top \\ 0 & P_i^\top & -y P_i^\top \\ \vdots & \vdots & \vdots \end{bmatrix} \begin{bmatrix} \mathbf{m}_1 \\ \mathbf{m}_2 \\ \mathbf{m}_3 \end{bmatrix} = 0 \quad (10)$$