

Modify By Confidence Level

Xiaolin Shen

July 7, 2018

1 The BTL Model —ModifyByConfidenceLevel

Suppose g is a K -dim vector, with one and only one component to be equal to 1, for each pair, $p(< w, v > | g_k = 1) = p^k(w \succ v) \times \prod_{k' \neq k} p^{k'}(w \succeq v)$. then the probability of generating a session observation d given the hidden aspect a is defined as:

$$p(d|g, \theta, V, U) = p(d|\Theta^t, g) = \prod_{w \in W^d, v \in L^d, w \succ_R v} a^K \frac{w_k}{w_k + \theta^t v_k} \prod_{k' \neq k} \frac{\theta^t w_{k'}}{v_{k'} + \theta^t w_{k'}} \quad (1)$$

where $w \succ_R v$ is the ranking pair in R relation existing in session d , a is the given value.

Besides,

$$\begin{aligned} p(g|d, \Theta^t) &\propto p(g|u, \Theta^t) p(d|g, \Theta^t) \\ p(g_k = 1|d, \Theta^t) &\propto u_k^t \frac{w_k^t}{w_k^t + \theta^t l_k^t} \prod_{k' \neq k} [l_{k'}^t + \theta^t w_{k'}^t] \end{aligned} \quad (2)$$

First, let's use $\gamma(d, k, \Theta^t)$ to denote the conditional probability $p(g_k = 1|d, \Theta^t)$ given parameters in the t -th round, when the current session specific favorite aspect is $g_k = 1$, defined as follows

$$\begin{aligned} \gamma(d, k, \Theta^t) &= \frac{p(d, g|\Theta^t)}{\sum_g p(d, g|\Theta^t)} = \frac{p(g|\Theta^t) p(d|\Theta^t, g)}{\sum_g p(g|\Theta^t) p(d|\Theta^t, g)} \\ &= \frac{u_k \prod_{w \in W^d, v \in L^d, w \succ_R v} a^K \frac{w_k}{w_k + \theta^t v_k} \prod_{k' \neq k} \frac{\theta^t w_{k'}}{v_{k'} + \theta^t w_{k'}}}{\sum_{k=1}^K u_k \prod_{w \in W^d, v \in L^d, w \succ_R v} a^K \frac{w_k}{w_k + \theta^t v_k} \prod_{k' \neq k} \frac{\theta^t w_{k'}}{v_{k'} + \theta^t w_{k'}}} \end{aligned} \quad (3)$$

Note that $\forall d, \sum_k \gamma(d, k, \Theta^t) = 1$.

1.1 E-step

In the E-step of t -th EM round, compute the expectation $Q(\Theta^t) = E_G \ln p(D, G|\Theta)$

$$\begin{aligned}
E_G \ln p(D, G|\Theta) &= \sum_d \sum_{k=1}^K \gamma(d, k, \Theta^t) \ln p(d, g|\Theta) \\
&= \sum_d \sum_{k=1}^K \gamma(d, k, \Theta^t) \{ \ln u_k \\
&\quad + \sum_{w \in W_d, v \in V_d, w \succ_R v} [\ln a^K + \ln \frac{w_k}{w_k + \theta v_k} + \sum_{k' \neq k} \ln \frac{\theta w_{k'}}{v_{k'} + \theta w_{k'}}] \}
\end{aligned} \tag{4}$$

1.2 M-step

1.2.1 For u

first maximize $Q(\Theta^t)$ with respect to U . For each $u \in U$, eliminating constant terms, we have:

$$\begin{aligned}
\min -\sum_{u(d)=u} \sum_{k=1}^K \gamma(d, k, \Theta^t) \ln u_k \\
w.r.t \sum_k u_k = 1
\end{aligned} \tag{5}$$

Solving the above Lagrange function Equ. 5, we get

$$u_k = \frac{\sum_{u(d)=u} \gamma(d, k, \Theta^t)}{\sum_{s=1}^K \sum_{u(d)=u} \gamma(d, s, \Theta^t)} \tag{6}$$

1.2.2 For v

for the :

$$\ln \frac{y}{x} \geq 1 - \frac{x}{y}$$

we can derive a lower bound for the log-likelihood over the complete data, given the parameters learnt from previous round. hence, we obtain a minorization function of $\tilde{Q}(\Theta^t)$.

$$\begin{aligned}
\tilde{Q}(\Theta^t) &= \sum_d \sum_k \gamma(d, k, \Theta^t) \sum_{w \in W_d, v \in L_d} \{ \ln a^K + [\ln w_k + 1 - \ln(w_k^t + \theta^t v_k^t) - \frac{w_k + \theta v_k}{w_k^t + \theta^t v_k^t}] + \\
&\quad \sum_{k' \neq k} [\ln(\theta w_{k'}) + 1 - \ln(v_{k'}^t + \theta^t w_{k'}^t) - \frac{v_{k'} + \theta w_{k'}}{v_{k'}^t + \theta^t w_{k'}^t}] \} \dots \dots \dots (7)
\end{aligned}$$

$\tilde{Q}(\Theta^t)$ can be separated for each item v . Considering only the k -th component v_k , $\tilde{Q}(v_k, \Theta^t)$ involves two terms, one of which is relevant to observations $d \in W(v)$ where v acts as skyline object, the other is relevant to observations $d \in L(v)$ where v acts as comparisons, $\tilde{Q}(v_k, \Theta^t) = \tilde{Q}^1(v_k, \Theta^t) + \tilde{Q}^2(v_k, \Theta^t)$. Removing all constants and irrelevant terms for v_k , we have the following minorizing function:

$$\begin{aligned}
\tilde{Q}^1(v_k, \Theta^t) &= \sum_{d \in W(v)} |L_d| \ln v_k - v_k \sum_{d \in W(v)} \sum_{v' \in L_d} \left[\frac{\gamma(d, k, \Theta^t)}{\alpha(v, v', k, \Theta^t)} + \sum_{k' \neq k} \frac{\theta^t \gamma(d, k', \Theta^t)}{\alpha(v', v, k, \Theta^t)} \right] \\
\tilde{Q}^2(v_k, \Theta^t) &= -v_k \sum_{d \in L(v)} \sum_{v' \in W_d} \left[\frac{\theta^t \gamma(d, k, \Theta^t)}{\alpha(v', v, k, \Theta^t)} + \sum_{k' \neq k} \frac{\gamma(d, k', \Theta^t)}{\alpha(v, v', k, \Theta^t)} \right]
\end{aligned}$$

where:

$|L_d|$ is the number of objects being dominated in d ,
 $\alpha(v, v', k, \Theta^t) = v_k^t + \theta^t v_k'^t$.

By setting the partial derivative of $\frac{\partial \tilde{Q}(v_k, \Theta^t)}{\partial v_k} = 0$, we have:

$$\begin{aligned} \frac{1}{v_k} = & \frac{\sum_{d \in W(v)} \sum_{v' \in L_d} \left[\frac{\gamma(d, k, \Theta^t)}{\alpha(v, v', k, \Theta^t)} + \sum_{k' \neq k} \frac{\theta^t \gamma(d, k', \Theta^t)}{\alpha(v', v, k, \Theta^t)} \right]}{\sum_{d \in W(v)} |L_d|} \\ & + \frac{\sum_{d \in L(v)} \sum_{v' \in W_d} \left[\frac{\theta^t \gamma(d, k, \Theta^t)}{\alpha(v', v, k, \Theta^t)} + \sum_{k' \neq k} \frac{\gamma(d, k', \Theta^t)}{\alpha(v, v', k, \Theta^t)} \right]}{\sum_{d \in W(v)} |L_d|} \end{aligned}$$

1.2.3 For θ

Fix u, v , update θ by:

Fix $v \in V$ and $u \in U$, rearranging *Equ.7*, we have the solution for $\frac{\partial \tilde{Q}(\Theta^t)}{\partial \theta} = 0$ as:

$$\theta = \frac{(K-1) \sum_d |W_d| |L_d|}{\sum_d \sum_k \gamma(d, k, \Theta^t) \sum_{w, v} \left[\frac{v_k}{\alpha(w, v, k, \Theta^t)} + \sum_{k' \neq k} \frac{w_{k'}}{\alpha(v, w, k', \Theta^t)} \right]} \quad (7)$$