

The BTL Model

Xiaolin Shen

August 7, 2018

1 The Standard BTL Model

Suppose that there are K underlying aspects, the latent preference vector for each user is denoted as $u \in \mathbb{R}^K$, and $\sum_k u_k = 1$, latent item feature vector $i, j \in \mathbb{R}^K$.

Base on the standard BTL model:

$$p(i \succ j|u) = \frac{u^T i}{u^T i + u^T j}$$
$$p(i \prec j|u) = 1 - p(i \succ j|u)$$

We use the maximal likelihood estimation. The likelihood function can be written as follows. The model parameters are denoted as $\Theta = \{v \in V, u \in U\}$, \succ_d represents a pairwise ranking observation in a session d .

$$L(\Theta) = \prod_d \prod_{i \succ_d j} \frac{\sum_{k=1}^K u_k i_k}{\sum_{k=1}^K u_k i_k + \sum_{k=1}^K u_k j_k} \quad (1)$$

Thus, the log likelihood and its lower bound is:

$$l(\Theta) = \log L(\Theta) \quad (2)$$

$$\begin{aligned} &= \sum_u \sum_{d \in D_u} \sum_{i \succ_d j} \log \frac{\sum_k u_k i_k}{\sum_k u_k i_k + \sum_k u_k j_k} \\ &\geq \sum_u \sum_{d \in D_u} \sum_{i \succ_d j} \left[1 - \frac{\sum_k u_k i_k + \sum_k u_k j_k}{\sum_k u_k^t i_k^t + \sum_k u_k^t j_k^t} + \log \frac{\sum_k u_k i_k}{\sum_k u_k^t i_k^t + \sum_k u_k^t j_k^t} \right] \end{aligned} \quad (3)$$

where we apply

$$\ln \frac{y}{x} \geq 1 - \frac{x}{y}$$

To maximize the lower-bound in Equ. ?? we employ conjugate gradient method. We first fix all vs , and apply $\log \sum_k u_k i_k \geq \frac{\sum_k i_k \log u_k}{\sum_k i_k} + \log \sum_k i_k, \forall i_k \geq$

0. denote $c^t(i, j, d) = \sum_k u_k^t i_k^t + \sum_k u_k^t j_k^t$ in the t -th round of the iteration, we have

$$\frac{\partial l}{\partial u_k} = \sum_{d \in D_u} \sum_{i \succ_d j} \left(\frac{i_k}{u_k \sum_k i_k} - \frac{i_k + j_k}{c^t(i, j, d)} \right) = 0 \quad (4)$$

$$u_k = \frac{\sum_{d \in D_u} \sum_{i \succ_d j} \frac{i_k}{\sum_k i_k}}{\sum_{d \in D_u} \sum_{i \succ_d j} \frac{i_k + j_k}{c^t(i, j, d)}} \quad (5)$$

Next we fix all u s, and again apply the Jensen's inequality $\log \sum_k u_k i_k \geq \sum_k u_k \log i_k, \forall u_k \geq 0, \sum_k u_k = 1$. For convenience, we use v to denote a arbitral item, and O_u is the set of all observed pairwise ranking of user u , if $v \succ j$ is the winning item then we say $o \in W(v)$.

$$\frac{\partial l}{\partial v_k} = \sum_u \sum_{o \in W(v) \& \& o \in O_u} \frac{u_k}{i_k} - \sum_u \sum_{o \in O_u} \frac{u_k}{c^t(i, j, d)} = 0 \quad (6)$$

$$i_k = \frac{\sum_u \sum_{o \in O_u} \frac{u_k}{c^t(i, j, d)}}{\sum_u \sum_{o \in W(v) \& \& o \in O_u} u_k} \quad (7)$$