

The BTL Model

Xiaolin Shen

August 10, 2018

1 The Standard BTL Model

Suppose that there are K underlying aspects, the latent preference vector for each user is denoted as $u \in R^{+K}$, and $\sum_k u_k = 1$, latent item feature vector $i, j \in R^{+K}$.

Base on the standard BTL model:

$$p(i \succ j|u) = \frac{u^T i}{u^T i + u^T j}$$

$$p(i \prec j|u) = 1 - p(i \succ j|u)$$

We use the maximal likelihood estimation. The likelihood function can be written as follows. The model parameters are denoted as $\Theta = \{v \in V, u \in U\}$, \succ_d represents a pairwise ranking observation in a session d .

$$L(\Theta) = \prod_{d \in D} \prod_{i \succ_d j} \frac{\sum_{k=1}^K u_k i_k}{\sum_{k=1}^K u_k i_k + \sum_{k=1}^K u_k j_k} \quad (1)$$

Thus, the log likelihood and its lower bound is:

$$l(\Theta) = \log L(\Theta) \quad (2)$$

$$= \sum_u \sum_{d \in D_u} \sum_{i \succ_d j} \log \frac{\sum_k u_k i_k}{\sum_k u_k i_k + \sum_k u_k j_k} \quad (3)$$

$$\geq \sum_u \sum_{d \in D_u} \sum_{i \succ_d j} \left[1 - \frac{\sum_k u_k i_k + \sum_k u_k j_k}{\sum_k u_k^t i_k^t + \sum_k u_k^t j_k^t} + \log \frac{\sum_k u_k i_k}{\sum_k u_k^t i_k^t + \sum_k u_k^t j_k^t} \right]$$

where we apply

$$\ln \frac{y}{x} \geq 1 - \frac{x}{y}$$

To maximize the lower-bound in *Equ.3* we employ conjugate gradient method.

We first fix all latent vectors for the item ¹, and apply $\log \sum_k u_k i_k \geq \frac{\sum_k i_k \log u_k}{\sum_k i_k} + \log \sum_k i_k, \forall i_k \geq 0$. Let's compute $c^t(d, i, j) = \sum_k u_k^t i_k^t + \sum_k u_k^t j_k^t, c_k^t(d, i, j) =$

¹We use the supscript t for parameters in the t^{th} round

$\frac{i_k^t + j_k^t}{c^t(d, i, j)}$, and $f_k^t(d, i) = \frac{i_k^t}{\sum_k i_k^t}$ for all pairwise ranking observations in d using the t -th round parameters, we have

$$\frac{\partial l}{\partial u_k} = \sum_{d \in D_u} \sum_{i \succ_d j} \left(\frac{f_k^t(d, i)}{u_k} - c_k^t(d, i, j) \right) = 0 \quad (4)$$

$$(5)$$

Here we omit the constraint that $u_k \geq 0$ and hope that this constraint will be satisfied by the solution.

$$u_k = \frac{\sum_{d \in D_u} \sum_{i \succ_d j} \frac{i_k}{\sum_k i_k}}{\sum_{k=1}^K \sum_{d \in D_u} \sum_{i \succ_d j} \frac{i_k + j_k}{c^t(i, j, d)}} \quad (6)$$

Next we fix all u s, and again apply the Jensen's inequality $\log \sum_k u_k i_k \geq \sum_k u_k \log i_k, \forall u_k \geq 0, \sum_k u_k = 1$. For convenience, we use v to denote a arbitral item, and O_u is the set of all observed pairwise ranking of user u , if $v \succ j$ is the winning item then we say $o \in W(v)$.

$$\frac{\partial l(\Theta)}{\partial v_k} = \sum_u \sum_{o \in W(v) \& o \in O_u} \frac{u_k}{i_k} - \sum_u \sum_{o \in O_u} \frac{u_k}{c^t(i, j, d)} = 0 \quad (7)$$

$$i_k = \frac{\sum_u \sum_{o \in O_u} \frac{u_k}{c^t(i, j, d)}}{\sum_u \sum_{o \in W(v) \& o \in O_u} u_k} \quad (8)$$