

The BTL Model

Xiaolin Shen

August 5, 2018

1 The Standard BTL Model –

Base on the standard BTL model:

$$\begin{aligned} p(i \succ j|u) &= \frac{u_i}{u_i + u_j} \\ p(i \prec j|u) &= \frac{u_j}{u_j + u_i} \end{aligned}$$

Suppose that there are K underlying aspects. for each pair, $p(< w, v >) = p^k(w \succ v)$. then the probability of generating a session observation d is defined as:

$$p(< w, l > |V, U) = \prod_{k=1}^K \left[\frac{u_k w_k}{u_k w_k + u_k v_k} \right] \quad (1)$$

The likelihood function can be written as follows. The model parameters are denoted as $\Theta = \{v \in V, u \in U\}$

$$L(\Theta) = p(D|\Theta) = \prod_{d \in D} \prod_{w \in W^d, v \in L^d} \prod_{k=1}^K \left[\frac{u_k w_k}{u_k w_k + u_k v_k} \right] \quad (2)$$

Thus, the log likelihood is:

$$l(\Theta) = \log L(\Theta) \quad (3)$$

$$\begin{aligned} &= \log \prod_{d \in D} \prod_{w \in W^d, v \in L^d} \prod_{k=1}^K \left[\frac{u_k w_k}{u_k w_k + u_k v_k} \right] \\ &= \sum_{d \in D} \log \prod_{w \in W^d, v \in L^d} \prod_{k=1}^K \left[\frac{u_k w_k}{u_k w_k + u_k v_k} \right] \end{aligned} \quad (4)$$

afterwards,