

STA 602 - Intro to Bayesian Statistics

Lecture 13

Li Ma

Duke University

Bayesian inference under independent Normal-inverse-Wishart priors

- Recall our sampling model

$$\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n \mid \boldsymbol{\theta}, \Sigma \stackrel{\text{iid}}{\sim} \mathcal{N}(\boldsymbol{\mu}, \Sigma).$$

- Independent prior specification:

$$p(\boldsymbol{\theta}, \Sigma) = p(\boldsymbol{\theta})p(\Sigma)$$

with

$$\boldsymbol{\theta} \sim \mathcal{N}(\boldsymbol{\mu}_0, \Lambda_0)$$

and

$$\Sigma \sim \text{IW}(v_0, \mathbf{S}_0).$$

Find the full conditionals

- ▶ Let's derive the full conditionals so we can do Gibbs sampling
- ▶ The likelihood is

$$L(\boldsymbol{\theta}, \Sigma; \mathbf{y}_1, \dots, \mathbf{y}_n) = p(\mathbf{y}_1, \dots, \mathbf{y}_n | \boldsymbol{\mu}, \Sigma) \propto |\Sigma|^{-n/2} e^{-\frac{1}{2} \sum_i (\mathbf{y}_i - \boldsymbol{\theta})' \Sigma^{-1} (\mathbf{y}_i - \boldsymbol{\theta})}.$$

- ▶ The prior is given by

$$p(\boldsymbol{\theta}) \propto e^{-\frac{1}{2} (\boldsymbol{\theta} - \boldsymbol{\mu}_0)' \Lambda_0^{-1} (\boldsymbol{\theta} - \boldsymbol{\mu}_0)}$$

and

$$p(\Sigma) \propto |\Sigma|^{-\frac{\nu_0 + p + 1}{2}} e^{-\frac{1}{2} \text{tr}(\mathbf{S}_0 \Sigma^{-1})}$$

The full joint probability

- The full joint probability is then

$$\begin{aligned} p(\mathbf{y}_1, \dots, \mathbf{y}_n, \boldsymbol{\theta}, \Sigma) \propto & |\Sigma|^{-n/2} e^{-\frac{1}{2} \sum_i (\mathbf{y}_i - \boldsymbol{\theta})' \Sigma^{-1} (\mathbf{y}_i - \boldsymbol{\theta})} \times \\ & e^{-\frac{1}{2} (\boldsymbol{\theta} - \boldsymbol{\mu}_0)' \Lambda_0^{-1} (\boldsymbol{\theta} - \boldsymbol{\mu}_0)} \times \\ & |\Sigma|^{-\frac{\nu_0 + p + 1}{2}} e^{-\frac{1}{2} \text{tr}(\mathbf{S}_0 \Sigma^{-1})} \end{aligned}$$

The full conditional of $\boldsymbol{\theta}$

- ▶ Viewing the joint probability as a function in $\boldsymbol{\theta}$, we have

$$\begin{aligned} p(\boldsymbol{\theta}|\mathbf{y}_1, \dots, \mathbf{y}_n, \Sigma) &\propto e^{-\frac{1}{2} \sum_i (\mathbf{y}_i - \boldsymbol{\theta})' \Sigma^{-1} (\mathbf{y}_i - \boldsymbol{\theta}) - \frac{1}{2} (\boldsymbol{\theta} - \boldsymbol{\mu}_0)' \Lambda_0^{-1} (\boldsymbol{\theta} - \boldsymbol{\mu}_0)} \\ &\propto e^{-\frac{1}{2} [n\boldsymbol{\theta}' \Sigma^{-1} \boldsymbol{\theta} - 2n\bar{\mathbf{y}}' \Sigma^{-1} \boldsymbol{\theta} + \boldsymbol{\theta}' \Lambda_0^{-1} \boldsymbol{\theta} + 2\boldsymbol{\mu}_0' \Lambda_0^{-1} \boldsymbol{\theta}]} \\ &\propto e^{-\frac{1}{2} [\boldsymbol{\theta}' (n\Sigma^{-1} + \Lambda_0^{-1}) \boldsymbol{\theta} - 2(n\bar{\mathbf{y}}' \Sigma^{-1} + \boldsymbol{\mu}_0' \Lambda_0^{-1}) \boldsymbol{\theta}]} \end{aligned}$$

- ▶ Now let Λ_n be such that

$$\Lambda_n^{-1} = n\Sigma^{-1} + \Lambda_0^{-1}$$

and

$$\boldsymbol{\mu}_n = \Lambda_n (n\Sigma^{-1} \bar{\mathbf{y}} + \Lambda_0^{-1} \boldsymbol{\mu}_0).$$

- ▶ Then by completion of squares we have

$$\begin{aligned} p(\boldsymbol{\theta}|\mathbf{y}_1, \dots, \mathbf{y}_n, \Sigma) &\propto e^{-\frac{1}{2} (\boldsymbol{\theta}' \Lambda_n^{-1} \boldsymbol{\theta} - 2\boldsymbol{\mu}_n' \Lambda_n^{-1} \boldsymbol{\theta})} \\ &\propto e^{-\frac{1}{2} (\boldsymbol{\theta} - \boldsymbol{\mu}_n)' \Lambda_n^{-1} (\boldsymbol{\theta} - \boldsymbol{\mu}_n)}, \end{aligned}$$

which is exactly a $N(\boldsymbol{\mu}_n, \Lambda_n)$.

- ▶ Compare this with univariate case — similar interpretation.

Full conditional of Σ

- ▶ Viewing the joint probability as a function in Σ , we have

$$p(\Sigma | \mathbf{y}_1, \dots, \mathbf{y}_n, \boldsymbol{\theta}) \propto |\Sigma|^{-(n+v_0+p+1)/2} e^{-\frac{1}{2} [\sum_i (\mathbf{y}_i - \boldsymbol{\theta})' \Sigma^{-1} (\mathbf{y}_i - \boldsymbol{\theta}) + \text{tr}(\mathbf{S}_0 \Sigma^{-1})]}.$$

- ▶ Two useful properties of $\text{tr}(\cdot)$:

- (i) For two matrix $A_{k \times m}$ and $B_{m \times k}$, we can prove (by definition of trace) that

$$\text{tr}(AB) = \text{tr}(BA).$$

Note that AB is $k \times k$ and BA is $m \times m$.

- (ii) Also by definition of trace, we have for two square matrix A and B ,

$$\text{tr}(A + B) = \text{tr}(A) + \text{tr}(B).$$

Full conditional of Σ

- Now note that $(\mathbf{y}_i - \boldsymbol{\theta})' \Sigma^{-1} (\mathbf{y}_i - \boldsymbol{\theta})$ is a scalar (i.e., a 1×1), and so it is equal to its trace

$$(\mathbf{y}_i - \boldsymbol{\theta})' \Sigma^{-1} (\mathbf{y}_i - \boldsymbol{\theta}) = \text{tr} \left((\mathbf{y}_i - \boldsymbol{\theta}) (\mathbf{y}_i - \boldsymbol{\theta})' \Sigma^{-1} \right)$$

by Property (i).

- Then by Property (ii)

$$\begin{aligned} \sum_i (\mathbf{y}_i - \boldsymbol{\theta})' \Sigma^{-1} (\mathbf{y}_i - \boldsymbol{\theta}) &= \sum_i \text{tr} \left((\mathbf{y}_i - \boldsymbol{\theta}) (\mathbf{y}_i - \boldsymbol{\theta})' \Sigma^{-1} \right) \\ &= \text{tr} \left(\sum_i (\mathbf{y}_i - \boldsymbol{\theta}) (\mathbf{y}_i - \boldsymbol{\theta})' \Sigma^{-1} \right) \\ &= \text{tr} (\mathbf{S}_{\boldsymbol{\theta}} \Sigma^{-1}) \end{aligned}$$

where

$$\mathbf{S}_{\boldsymbol{\theta}} = \sum_i (\mathbf{y}_i - \boldsymbol{\theta}) (\mathbf{y}_i - \boldsymbol{\theta})',$$

which is the *residual sum of squared matrix* of the observations.

Full conditional of Σ

- ▶ By Property (ii) again,

$$\text{tr}(\mathbf{S}_\theta \Sigma^{-1}) + \text{tr}(\mathbf{S}_0 \Sigma^{-1}) = \text{tr}((\mathbf{S}_\theta + \mathbf{S}_0) \Sigma^{-1})$$

- ▶ Therefore, putting all pieces together, we have

$$p(\Sigma | \mathbf{y}_1, \dots, \mathbf{y}_n, \boldsymbol{\theta}) \propto |\Sigma|^{-(n+v_0+p+1)/2} e^{-\frac{1}{2} \text{tr}((\mathbf{S}_\theta + \mathbf{S}_0) \Sigma^{-1})},$$

which we recognize as again an inverse-Wishart($v_n, \mathbf{S}_0 + \mathbf{S}_\theta$) distribution where $v_n = v_0 + n$.

- ▶ That is

$$\Sigma | \mathbf{y}_1, \dots, \mathbf{y}_n, \boldsymbol{\theta} \sim \text{IW}(v_n, \mathbf{S}_0 + \mathbf{S}_\theta).$$

- ▶ v_0 is the prior degrees of freedom and \mathbf{S}_0 the prior sum of squares matrix.
- ▶ The full conditional expectation of Σ is

$$\begin{aligned} E(\Sigma | \mathbf{y}_1, \dots, \mathbf{y}_n, \boldsymbol{\theta}) &= \frac{\mathbf{S}_0 + \mathbf{S}_\theta}{v_n - p - 1} \\ &= \frac{v_0 - p - 1}{v_n - p - 1} \cdot \underbrace{\frac{\mathbf{S}_0}{v_0 - p - 1}}_{\text{prior mean of } \Sigma} + \frac{n}{v_n - p - 1} \cdot \underbrace{\frac{\mathbf{S}_\theta}{n}}_{\text{Empirical estimate of } \Sigma}. \end{aligned}$$

Gibbs sampling

- ▶ Initialize the chain at $(\boldsymbol{\theta}^{(0)}, \Sigma^{(0)})$.
- ▶ For $t = 1, 2, \dots$,
 - ▶ Update $\boldsymbol{\theta}$:
 - ▶ Compute

$$\begin{aligned}\left(\Lambda_n^{(t)}\right)^{-1} &= n\left(\Sigma^{(t-1)}\right)^{-1} + \Lambda_0^{-1}, \\ \boldsymbol{\mu}_n^{(t)} &= \Lambda_n^{(t)}\left(n\left(\Sigma^{(t-1)}\right)^{-1} \bar{y} + \Lambda_0^{-1} \boldsymbol{\mu}_0\right)\end{aligned}$$

- ▶ Draw

$$\boldsymbol{\theta}^{(t)} \sim \text{N}(\boldsymbol{\mu}_n^{(t)}, \Lambda_n^{(t)})$$

- ▶ Update Σ :
 - ▶ Compute

$$\mathbf{S}_{\boldsymbol{\theta}^{(t)}} = \sum_i (\mathbf{y}_i - \boldsymbol{\theta}^{(t)})(\mathbf{y}_i - \boldsymbol{\theta}^{(t)})'.$$

- ▶ Draw

$$\Sigma^{(t)} \sim \text{IW}\left(v_n, \mathbf{S}_0 + \mathbf{S}_{\boldsymbol{\theta}^{(t)}}\right).$$

- ▶ Discard suitable burn-in steps.

Back to air pollutant example

- Suppose now we measure two pollutants (e.g., PM2.5 and SO2) concurrently 16 times on a day, so our data are bivariate

(104, 100), (105, 102), (103, 101), (102, 104), (105, 108), (107, 108),
(106, 103), (104, 104), (103, 106), (106, 107), (105, 105), (102, 101),
(102, 100), (108, 106), (105, 105), (104, 105)

Prior specification

- ▶ Suppose based on historical data, both pollutants have average level of 100. So we set

$$\boldsymbol{\mu}_0 = (100, 100)'.$$

- ▶ Suppose the historical distribution of PM2.5 has a standard deviation of about 10, and SO2 has a standard deviation of about 5, and their values tend to rise or fall together with a correlation of about 0.3. Turning this into our prior,

$$\Lambda_0 = \begin{pmatrix} 10^2 & 0.3 \times 10 \times 5 \\ 0.3 \times 10 \times 5 & 5^2 \end{pmatrix} = \begin{pmatrix} 100 & 15 \\ 15 & 25 \end{pmatrix}.$$

Prior specification on Σ

- ▶ Suppose we have very little idea about the value of Σ , so we want to choose a weak prior.
- ▶ This can be achieved by setting ν_0 to a small value, such as $\nu_0 = p + 2$. (Note that the inverse-Wishart prior requires $\nu_0 > p + 1$ to have a finite mean.)
- ▶ As for the prior mean \mathbf{S}_0 , (e.g., $\mathbf{S}_0/(\nu_0 - p - 1)$), for simplicity, we use a diagonal matrix

$$\mathbf{S}_0 = \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix}$$

where the diagonal is chosen to be 4, corresponding to marginal standard deviations of 2 for the device readings.


- ▶ The off-diagonals are set to 0 because we don't really know anything about how the device reading errors are correlated for the two pollutants.

Reading the data

```
library(mvtnorm) # for drawing multivariate normal
library(MCMCpack) # for drawing inverse-Wishart
library(coda) # for MCMC diagnostics
options(digits=4) # tidy up print outs

y <- matrix(c(104,100,105,102,103,101,102,104,105,108,107,108,106,103,104,104,
              103,106,106,107,105,105,102,101,102,100,108,106,105,105,104,105),
            ncol=2,byrow=TRUE)
n <- nrow(y) # sample size
p <- ncol(y) # dimensionality

plot(y[,1],y[,2],xlab="PM2.5",ylab="SO2")
```



Prior specification

```
# prior for theta
mu.0 <- c(100,100)
Lambda.0 = matrix(c(100,15,15,25),ncol=2,byrow=TRUE)

# prior for Sigma
nu.0 <- p + 2 # a very weak prior
S0 <- (nu.0-p-1)*matrix(c(4,0,0,4),ncol=2,byrow=TRUE)

ybar <- apply(y,2,mean)
nu.n <- nu.0 + n
```

Gibbs sampling

```
niter <- 10000 # total number of iterations
nburnin <- 1000 # 1000 burn-in steps

THETA <- matrix(NA,nrow=niter,ncol=p) # matrix for storing the draws for theta
colnames(THETA) <- c("theta1","theta2")

THETA.init <- ybar # Initial values set to sample mean
THETA.curr <- THETA.init # the theta value at current iteration

SIGMA <- matrix(NA,nrow=niter,ncol=p*p) # matrix for storing the draws for Sigma
colnames(SIGMA) <- c("sigma11","sigma12","sigma21","sigma22")

SIGMA.init <- cov(y) # initial value set to sample covariance
SIGMA.curr <- SIGMA.init # the Sigma value at current iteration

### Start Gibbs sampling
for (t in 1:niter) {

  ## Update theta
  Lambda.n <- solve(n*solve(SIGMA.curr)+solve(Lambda.0))
  mu.n <- Lambda.n %*% (n*solve(SIGMA.curr,ybar)+solve(Lambda.0,mu.0))
  THETA.curr <- rmvnorm(1,mean=mu.n,sigma=Lambda.n)


  ## Update Sigma
  S.theta <- (t(y)-c(THETA.curr))%*% t(t(y)-c(THETA.curr))
  SIGMA.curr <- riwish(v=nu.n,S=S0+S.theta)

  ## Save the current iteration
  THETA[t,] <- THETA.curr
  SIGMA[t,] <- SIGMA.curr

}
```

Histogram of MCMC draws for θ

```
ggplot(data.frame(THETA), aes(x=theta1, y=theta2) ) +  
  labs(x=expression(theta_1), y=expression(theta_2)) +  
  geom_bin2d(bins=70) +  
  scale_fill_continuous(type = "viridis") +  
  lims(x=c(101,107), y=c(101,107))
```







MCMC diagnostics

```
THETA.mcmc <- mcmc(THETA[-(1:nburnin),], start=nburnin+1)
summary(THETA.mcmc)
```

```
##
## Iterations = 1001:10000
## Thinning interval = 1
## Number of chains = 1
## Sample size per chain = 9000
##
## 1. Empirical mean and standard deviation for each variable,
##    plus standard error of the mean:
##
##           Mean      SD Naive SE Time-series SE
## theta1  104 0.448  0.00472      0.00472
## theta2  104 0.658  0.00694      0.00694
##
## 2. Quantiles for each variable:
##
##           2.5% 25% 50% 75% 97.5%
## theta1  104 104 104 105   105
## theta2  103 104 104 104   105
```

Trace plots for θ

```
plot (THETA.mcmc)
```





ESS for θ

```
effectiveSize (THETA.mcmc)
```

```
## theta1 theta2  
##    9000    9000
```

Autocorrelation plot for θ

```
autocorr.plot(THETA.mcmc, lag.max=100)
```











MCMC diagnostics

```
SIGMA.mcmc <- mcmc(SIGMA[-(1:nburnin),], start=nburnin+1)
summary(SIGMA.mcmc)
```

```
##
## Iterations = 1001:10000
## Thinning interval = 1
## Number of chains = 1
## Sample size per chain = 9000
##
## 1. Empirical mean and standard deviation for each variable,
##    plus standard error of the mean:
##
##           Mean      SD Naive SE Time-series SE
## sigma11 3.23 1.21   0.0128      0.0128
## sigma12 2.65 1.42   0.0150      0.0157
## sigma21 2.65 1.42   0.0150      0.0157
## sigma22 6.92 2.59   0.0273      0.0291
##
## 2. Quantiles for each variable:
##
##           2.5%  25%  50%  75% 97.5%
## sigma11 1.644 2.40 2.99 3.79 6.33
## sigma12 0.635 1.71 2.39 3.31 6.07
## sigma21 0.635 1.71 2.39 3.31 6.07
## sigma22 3.553 5.13 6.38 8.10 13.34
```

Trace plots for Σ

```
plot(SIGMA.mcmc)
```







ESS for Σ

```
effectiveSize(SIGMA.mcmc)
```

```
## sigma11 sigma12 sigma21 sigma22  
##      8916      8200      8200      7948
```

Autocorrelation plot for Σ

```
autocorr.plot (SIGMA.mcmc, lag.max=100)
```



With a stronger prior on Σ with $\nu_0 = 50$

- ▶ Suppose we have a better idea about the value of Σ , so we want to choose a stronger prior.
- ▶ For example we set $\nu_0 = 50$.
- ▶ Now the prior mean $\mathbf{S}_0/(\nu_0 - p - 1) = \mathbf{S}_0/47$. Thus if we want the same prior mean as before, we set

$$\mathbf{S}_0 = 47 \cdot \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix} = \begin{pmatrix} 188 & 0 \\ 0 & 188 \end{pmatrix}$$


```
# prior for theta
mu.0 <- c(100,100)
Lambda.0 = matrix(c(100,15,15,25),ncol=2,byrow=TRUE)

# prior for Sigma
nu.0 <- 50 # a stronger prior on covariance now
S0 <- (nu.0-p-1) * matrix(c(4,0,0,4),ncol=2,byrow=TRUE) # maintaining the same prior mean

ybar <- apply(y,2,mean)
nu.n <- nu.0 + n
```

Histogram of MCMC draws for θ

```
ggplot(data.frame(THETA), aes(x=theta1, y=theta2)) +  
  labs(x=expression(theta_1), y=expression(theta_2)) +  
  geom_bin2d(bins=70) +  
  scale_fill_continuous(type = "viridis") +  
  lims(x=c(101,107), y=c(101,107))
```



- ▶ How does this compare to the case with $v_0 = 4$?
- ▶ Why is the posterior spread of θ smaller?





MCMC diagnostics

```
THETA.mcmc <- mcmc(THETA[-(1:nburnin),], start=nburnin+1)
summary(THETA.mcmc)
```

```
##
## Iterations = 1001:10000
## Thinning interval = 1
## Number of chains = 1
## Sample size per chain = 9000
##
## 1. Empirical mean and standard deviation for each variable,
##    plus standard error of the mean:
##
##           Mean      SD Naive SE Time-series SE
## theta1  104 0.480  0.00506      0.00506
## theta2  104 0.538  0.00568      0.00568
##
## 2. Quantiles for each variable:
##
##           2.5% 25% 50% 75% 97.5%
## theta1  103 104 104 105  105
## theta2  103 104 104 104  105
```

Trace plots for θ

```
plot (THETA.mcmc)
```





ESS for θ

```
effectiveSize(THETA.mcmc)
```

```
## theta1 theta2  
##    9000    9000
```

Autocorrelation plot for θ

```
autocorr.plot(THETA.mcmc, lag.max=100)
```











MCMC diagnostics

```
SIGMA.mcmc <- mcmc(SIGMA[-(1:nburnin),], start=nburnin+1)
summary(SIGMA.mcmc)
```

```
##
## Iterations = 1001:10000
## Thinning interval = 1
## Number of chains = 1
## Sample size per chain = 9000
##
## 1. Empirical mean and standard deviation for each variable,
##    plus standard error of the mean:
##
##           Mean      SD Naive SE Time-series SE
## sigma11  3.798 0.700  0.00738      0.00738
## sigma12  0.685 0.546  0.00576      0.00576
## sigma21  0.685 0.546  0.00576      0.00576
## sigma22  4.760 0.872  0.00919      0.00919
##
## 2. Quantiles for each variable:
##
##           2.5%   25%   50%   75%  97.5%
## sigma11  2.661 3.300 3.724 4.20  5.39
## sigma12 -0.337 0.324 0.668 1.02  1.82
## sigma21 -0.337 0.324 0.668 1.02  1.82
## sigma22  3.323 4.138 4.658 5.28  6.71
```

Trace plots for Σ

```
plot(SIGMA.mcmc)
```







ESS for Σ

```
effectiveSize(SIGMA.mcmc)
```

```
## sigma11 sigma12 sigma21 sigma22  
##      9000      9000      9000      9000
```

Autocorrelation plot for Σ

```
autocorr.plot(SIGMA.mcmc, lag.max=100)
```




Stronger prior on θ

- Suppose the historical distribution of PM2.5 has a standard deviation of about **1** (vs 10 before), and SO2 has a standard deviation of about **0.5** vs (5 before), still with a correlation of about 0.3. Turning this into our prior,

$$\Lambda_0 = \begin{pmatrix} 1^2 & 0.3 \times 1 \times 0.5 \\ 0.3 \times 1 \times 0.5 & 0.5^2 \end{pmatrix} = \begin{pmatrix} 1 & 0.15 \\ 0.15 & 0.25 \end{pmatrix}.$$

With weak prior on Σ : $v_0 = p + 2$

```
ggplot(data.frame(THETA), aes(x=theta1, y=theta2)) +  
  labs(x=expression(theta_1), y=expression(theta_2)) +  
  geom_bin2d(bins=70) +  
  scale_fill_continuous(type = "viridis") +  
  lims(x=c(99,105), y=c(99,105))
```



- Why is there a lot *more* shrinkage toward prior mean?





MCMC diagnostics

```
THETA.mcmc <- mcmc(THETA[-(1:nburnin),], start=nburnin+1)
summary(THETA.mcmc)
```

```
##
## Iterations = 1001:10000
## Thinning interval = 1
## Number of chains = 1
## Sample size per chain = 9000
##
## 1. Empirical mean and standard deviation for each variable,
##    plus standard error of the mean:
##
##           Mean      SD Naive SE Time-series SE
## theta1  103 0.618  0.00651      0.01197
## theta2  101 0.513  0.00540      0.00707
##
## 2. Quantiles for each variable:
##
##           2.5% 25% 50% 75% 97.5%
## theta1 101.2 102 103 103 104
## theta2  99.9 101 101 101 102
```

Trace plots for θ

```
plot (THETA.mcmc)
```



ESS for θ



```
effectiveSize (THETA.mcmc)
```

```
## theta1 theta2  
## 2662 5259
```

- ▶ This stronger prior induced much stronger autocorrelation and smaller ESS.
- ▶ Why?
 - ▶ The prior concentrated near a value far from the empirical estimate.
 - ▶ Stronger posterior dependency between θ and Σ .

Autocorrelation plot for θ

```
autocorr.plot(THETA.mcmc, lag.max=100)
```











MCMC diagnostics

```
SIGMA.mcmc <- mcmc(SIGMA[-(1:nburnin)],,start=nburnin+1)
summary(SIGMA.mcmc)
```

```
##
## Iterations = 1001:10000
## Thinning interval = 1
## Number of chains = 1
## Sample size per chain = 9000
##
## 1. Empirical mean and standard deviation for each variable,
##    plus standard error of the mean:
##
##           Mean      SD Naive SE Time-series SE
## sigma11  6.80 3.55   0.0374         0.0626
## sigma12  8.24 4.24   0.0447         0.0645
## sigma21  8.24 4.24   0.0447         0.0645
## sigma22 16.00 6.63   0.0699         0.0842
##
## 2. Quantiles for each variable:
##
##           2.5%   25%   50%   75%  97.5%
## sigma11  2.63   4.35   5.93   8.3   15.9
## sigma12  2.73   5.30   7.32  10.2   18.9
## sigma21  2.73   5.30   7.32  10.2   18.9
## sigma22  7.33  11.38  14.59  19.1   32.3
```

Trace plots for Σ

```
plot(SIGMA.mcmc)
```







ESS for Σ

```
effectiveSize(SIGMA.mcmc)
```

```
## sigma11 sigma12 sigma21 sigma22  
##      3212      4310      4310      6200
```


Autocorrelation plot for Σ

```
autocorr.plot (SIGMA.mcmc, lag.max=100)
```



With a strong prior on Σ : $v_0 = 50$

```
ggplot(data.frame(THETA), aes(x=theta1, y=theta2)) +  
  labs(x=expression(theta_1), y=expression(theta_2)) +  
  geom_bin2d(bins=70) +  
  scale_fill_continuous(type = "viridis") +  
  lims(x=c(99,105), y=c(99,105))
```



- Why is there a lot *less* shrinkage toward prior mean?





MCMC diagnostics

```
THETA.mcmc <- mcmc(THETA[-(1:nburnin),], start=nburnin+1)
summary(THETA.mcmc)
```

```
##
## Iterations = 1001:10000
## Thinning interval = 1
## Number of chains = 1
## Sample size per chain = 9000
##
## 1. Empirical mean and standard deviation for each variable,
##    plus standard error of the mean:
##
##           Mean      SD Naive SE Time-series SE
## theta1  103 0.513  0.00540      0.00681
## theta2  102 0.421  0.00444      0.00537
##
## 2. Quantiles for each variable:
##
##           2.5% 25% 50% 75% 97.5%
## theta1  102 103 103 104 104
## theta2  101 102 102 102 103
```

Trace plots for θ

```
plot (THETA.mcmc)
```





ESS for θ

```
effectiveSize(THETA.mcmc)
```

```
## theta1 theta2  
##    5673    6139
```


Autocorrelation plot for θ

```
autocorr.plot(THETA.mcmc, lag.max=100)
```











MCMC diagnostics

```
SIGMA.mcmc <- mcmc(SIGMA[-(1:nburnin),], start=nburnin+1)
summary(SIGMA.mcmc)
```

```
##
## Iterations = 1001:10000
## Thinning interval = 1
## Number of chains = 1
## Sample size per chain = 9000
##
## 1. Empirical mean and standard deviation for each variable,
##    plus standard error of the mean:
##
##           Mean      SD Naive SE Time-series SE
## sigma11  4.12 0.827  0.00872      0.0101
## sigma12  1.32 0.757  0.00798      0.0101
## sigma21  1.32 0.757  0.00798      0.0101
## sigma22  6.03 1.200  0.01265      0.0148
##
## 2. Quantiles for each variable:
##
##           2.5%   25%   50%   75%  97.5%
## sigma11  2.8141 3.534 4.01 4.58  6.06
## sigma12 -0.0119 0.803 1.26 1.76  2.95
## sigma21 -0.0119 0.803 1.26 1.76  2.95
## sigma22  4.1017 5.188 5.88 6.72  8.76
```

Trace plots for Σ

```
plot(SIGMA.mcmc)
```



ESS for Σ

```
effectiveSize(SIGMA.mcmc)
```

```
## sigma11 sigma12 sigma21 sigma22  
##      6675      5604      5604      6590
```

Autocorrelation plot for Σ

```
autocorr.plot (SIGMA.mcmc, lag.max=100)
```

