

Lower bounds on error probability of quantum channel discrimination by the Bures angle and the trace distance

Ryo Ito and Ryuhei Mori

Tokyo Institute of Technology

Sept. 7, 2023

Quantum channel discrimination problem

Definition (Quantum channel discrimination problem)

For fixed finite set of quantum channels $\{\Phi_1, \Phi_2, \dots, \Phi_k\}$ and a probability distribution $(p_i)_{i=1,2,\dots,k}$,

Input: An oracle $\mathcal{O} = \Phi_i$ for $i \in \{1, 2, \dots, k\}$ drawn w.p. p_i


Output: $i \in \{1, 2, \dots, k\}$ satisfying $\mathcal{O} = \Phi_i$

- A fundamental problem in quantum information science

Algorithm for quantum channel discrimination problem

General algorithm calling the oracle n times.

- ▶ $\mathcal{O} \in \{\Phi_1, \dots, \Phi_k\}$
- ▶ $\Gamma_1, \dots, \Gamma_n$: Quantum channels
- ▶ M : Measurement



Discrimination of quantum states

Definition (Quantum state discrimination problem)

For fixed finite set of quantum states $\{\rho_1, \rho_2, \dots, \rho_k\}$ and a probability distribution $(p_i)_{i=1,2,\dots,k}$,

Input: A quantum state $\rho = \rho_i$ for $i \in \{1, 2, \dots, k\}$ drawn w.p. p_i

Output: $i \in \{1, 2, \dots, k\}$ satisfying $\rho = \rho_i$

Theorem (Holevo–Helstrom theorem

[Helstrom 1967], [Holevo 1972])

The largest success probability of discrimination of two quantum states is

$$\frac{1 + \|\rho_1 \rho_1 - \rho_2 \rho_2\|_1}{2}.$$

For more than two quantum states, the largest success probability is a solution of semidefinite programming (SDP).

Discrimination of two quantum channels

Theorem (Discrimination of two quantum channels)

The largest success probability of discrimination of *two* quantum channels with a *single* query is

$$p_{\text{succ}} = \frac{1 + \|p_1\Phi_1 - p_2\Phi_2\|_{\diamond}}{2}$$

where

$$\|\Gamma_A\|_{\diamond} := \max_{\rho_{AR}} \|(\Gamma_A \otimes \text{Id}_R)(\rho_{AR})\|_1.$$

Corollary (Discrimination by non-adaptive algorithms)

The largest success probability of discrimination of *two* quantum channels with *n non-adaptive (parallel)* queries is

$$p_{\text{succ}} = \frac{1 + \|p_1\Phi_1^{\otimes n} - p_2\Phi_2^{\otimes n}\|_{\diamond}}{2}.$$

Previous qualitative result

- ▶ There are two quantum channels that cannot be distinguished exactly by any non-adaptive algorithm, but **can be distinguished exactly** by an adaptive algorithm with two queries [Harrow, Hassidim, Leung, Watrous, 2010].
- ▶ For any two quantum channels, either of the followings is true [Yu and Zhou 2021]
 - ▶ The quantum channels can be distinguished **exactly** by finite queries.
 - ▶ The error probability of quantum channel discrimination decays **exponentially** in the number of queries.

Previous result and our result

# of channels	2	≥ 2
QUD	[Kawachi, Kawano, Le Gall, Tamaki 2019] Port-based teleportation	Grover search $\ell = 1$ [Zalka 1999]
QCD	[Pirandola, Laurenza, Lupo, Pereira, 2019] [This work]	[Zhuang, Pirandola, 2020]
QCGD		[This work]

Discrimination of two quantum channels

Quantum channel discrimination of $\Phi_A^{(1)}$ and $\Phi_A^{(2)}$.

$$\rho_1 = \Phi_A^{(1)} \Gamma_{AR}^{(n)} \Phi_A^{(1)} \Gamma_{AR}^{(n-1)} \cdots \Phi_A^{(1)} \Gamma_{AR}^{(1)} |0^n\rangle \langle 0^n|$$

$$\rho_2 = \Phi_A^{(2)} \Gamma_{AR}^{(n)} \Phi_A^{(2)} \Gamma_{AR}^{(n-1)} \cdots \Phi_A^{(2)} \Gamma_{AR}^{(1)} |0^n\rangle \langle 0^n|$$

From Holevo–Helstrom theorem,

$$\frac{1 - \frac{1}{2} \|\rho_1 - \rho_2\|_1}{2}.$$

The trace distance $D(\rho, \sigma) = \|\rho - \sigma\|_1$ satisfies

1. (Triangle inequality) $\forall \rho, \sigma, \tau \quad D(\rho, \sigma) \leq D(\rho, \tau) + D(\tau, \sigma)$
2. (Monotonicity) $\forall \rho, \sigma, \mathcal{E} \quad D(\mathcal{E}(\rho), \mathcal{E}(\sigma)) \leq D(\rho, \sigma)$

Hybrid argument

$$\rho^{(0)} = \Phi_A^{(1)} \Gamma_{AR}^{(n)} \Phi_A^{(1)} \Gamma_{AR}^{(n-1)} \dots \Phi_A^{(1)} \Gamma_{AR}^{(2)} \Phi_A^{(1)} \Gamma_{AR}^{(1)} |0^n\rangle \langle 0^n|$$

$$\rho^{(1)} = \Phi_A^{(1)} \Gamma_{AR}^{(n)} \Phi_A^{(1)} \Gamma_{AR}^{(n-1)} \dots \Phi_A^{(1)} \Gamma_{AR}^{(2)} \Phi_A^{(2)} \Gamma_{AR}^{(1)} |0^n\rangle \langle 0^n|$$

$$\rho^{(2)} = \Phi_A^{(1)} \Gamma_{AR}^{(n)} \Phi_A^{(1)} \Gamma_{AR}^{(n-1)} \dots \Phi_A^{(2)} \Gamma_{AR}^{(2)} \Phi_A^{(2)} \Gamma_{AR}^{(1)} |0^n\rangle \langle 0^n|$$

\vdots

$$\rho^{(n)} = \Phi_A^{(2)} \Gamma_{AR}^{(n)} \Phi_A^{(2)} \Gamma_{AR}^{(n-1)} \dots \Phi_A^{(2)} \Gamma_{AR}^{(2)} \Phi_A^{(2)} \Gamma_{AR}^{(1)} |0^n\rangle \langle 0^n|$$

$$\begin{aligned} \|\rho^{(0)} - \rho^{(n)}\|_1 &= \|\rho^{(0)} - \rho^{(n)}\|_1 \\ &\leq \sum_{i=0}^{n-1} \|\rho^{(i)} - \rho^{(i+1)}\|_1 \end{aligned}$$

Upper bound for the single step

$$\begin{aligned}
 \|\rho^{(i)} - \rho^{(i+1)}\|_1 &= \left\| \Phi_A^{(1)} \Gamma_{AR}^{(n)} \cdots \Phi_A^{(1)} \Gamma_{AR}^{(i+1)} \Phi_A^{(2)} \Gamma_{AR}^{(i)} \cdots \Phi_A^{(2)} \Gamma_{AR}^{(1)} |0^n\rangle \langle 0^n| \right. \\
 &\quad \left. - \Phi_A^{(1)} \Gamma_{AR}^{(n)} \cdots \Phi_A^{(1)} \Gamma_{AR}^{(i+2)} \Phi_A^{(2)} \Gamma_{AR}^{(i+1)} \cdots \Phi_A^{(2)} \Gamma_{AR}^{(1)} |0^n\rangle \langle 0^n| \right\|_1 \\
 &\leq \left\| \Phi_A^{(1)} \Gamma_{AR}^{(i+1)} \Phi_A^{(2)} \Gamma_{AR}^{(i)} \cdots \Phi_A^{(2)} \Gamma_{AR}^{(1)} |0^n\rangle \langle 0^n| \right. \\
 &\quad \left. - \Phi_A^{(2)} \Gamma_{AR}^{(i+1)} \Phi_A^{(2)} \Gamma_{AR}^{(i)} \cdots \Phi_A^{(2)} \Gamma_{AR}^{(1)} |0^n\rangle \langle 0^n| \right\|_1 \\
 &\leq \max_{\rho_{AR}} \left\| \Phi_A^{(1)}(\rho_{AR}) - \Phi_A^{(2)}(\rho_{AR}) \right\|_1 \\
 &= \left\| \Phi_A^{(1)} - \Phi_A^{(2)} \right\|_\diamond
 \end{aligned}$$

A simple lower bound of the error probability


Lemma (A simple lower bound)

The quantum channel discrimination of $\Phi^{(1)}$, $\Phi^{(2)}$

$$p_{\text{err}}(n) \geq \frac{1 - \frac{n}{2} \left\| \Phi_A^{(1)} - \Phi_A^{(2)} \right\|_{\diamond}}{2}$$

Triangle inequalities are not tight

For pure states with two real state vectors $|\psi_0\rangle$ and $|\psi_2\rangle$, the trace distance is equal to the **Euclidean distance** up to a constant factor.



Obviously, $\| |\psi_0\rangle - |\psi_2\rangle \|_2 < \| |\psi_0\rangle - |\psi_1\rangle \|_2 + \| |\psi_1\rangle - |\psi_2\rangle \|_2$.

Bures angle

- ▶ The fidelity: $F(\rho, \sigma) := \|\sqrt{\rho}\sqrt{\sigma}\|_1$.
- ▶ The **Bures angle**: $A(\rho, \sigma) := \arccos F(\rho, \sigma)$.

Lemma

$$\|p\rho - (1-p)\sigma\|_1 \leq \sqrt{1 - 4p(1-p)F(\rho, \sigma)^2}$$

The equality is satisfied if ρ and σ are pure states.

Bures angle satisfies **the triangle inequality** and **the monotonicity**.

From the triangle inequality of the Bures angle

$$A(\rho^{(0)}, \rho^{(n)}) \leq \sum_{i=0}^{n-1} A(\rho^{(i)}, \rho^{(i+1)})$$

Upper bound for the single step

$$\begin{aligned}
 A(\rho^{(i)}, \rho^{(i+1)}) &= A\left(\Phi_A^{(1)} \Gamma_{AR}^{(n)} \cdots \Phi_A^{(1)} \Gamma_{AR}^{(i+1)} \Phi_A^{(2)} \Gamma_{AR}^{(i)} \cdots \Phi_A^{(2)} \Gamma_{AR}^{(1)} |0^n\rangle \langle 0^n|, \right. \\
 &\quad \left. \Phi_A^{(1)} \Gamma_{AR}^{(n)} \cdots \Phi_A^{(1)} \Gamma_{AR}^{(i+2)} \Phi_A^{(2)} \Gamma_{AR}^{(i+1)} \cdots \Phi_A^{(2)} \Gamma_{AR}^{(1)} |0^n\rangle \langle 0^n| \right) \\
 &\leq A\left(\Phi_A^{(1)} \Gamma_{AR}^{(i+1)} \Phi_A^{(2)} \Gamma_{AR}^{(i)} \cdots \Phi_A^{(2)} \Gamma_{AR}^{(1)} |0^n\rangle \langle 0^n|, \right. \\
 &\quad \left. \Phi_A^{(2)} \Gamma_{AR}^{(i+1)} \cdots \Phi_A^{(2)} \Gamma_{AR}^{(1)} |0^n\rangle \langle 0^n| \right) \\
 &\leq \max_{\rho_{AR}} A\left(\Phi_A^{(1)}(\rho_{AR}), \Phi_A^{(2)}(\rho_{AR}) \right) \\
 &= \arccos \min_{\rho_{AR}} F\left(\Phi_A^{(1)}(\rho_{AR}), \Phi_A^{(2)}(\rho_{AR}) \right).
 \end{aligned}$$

The minimum of the fidelity can be evaluated by SDP.

Our result using the Bures angle

Theorem ([Ito and Mori 2021])

$$p_{\text{err}}(\textcolor{red}{n}) \geq \frac{1}{2} \left(1 - \sqrt{1 - 4p_1 p_2 \cos^2(\textcolor{red}{n}\tau)} \right)$$

where

$$\tau := \arccos \min_{\rho_{AR}} F \left(\Phi_A^{(1)}(\rho_{AR}), \Phi_A^{(2)}(\rho_{AR}) \right).$$

Corollary (Discrimination of two amplitude damping channels)

For the discrimination of two amplitude damping channels with parameters r_1 and r_2 ,


$$p_{\text{err}}(\textcolor{red}{n}) \geq \frac{1}{2} \left(1 - \sqrt{1 - 4p_1 p_2 \cos^2(\textcolor{red}{n}\Delta)} \right)$$

where $\Delta := \arccos(\sqrt{r_1 r_2} + \sqrt{(1-r_1)(1-r_2)})$.

Especially, when $p_1 = p_2 = 1/2$,

$$p_{\text{err}}(\textcolor{red}{n}) \geq \frac{1}{2} (1 - \sin(\textcolor{red}{n}\Delta)).$$

Improvement of the triangle inequality



Let $k \in \{0, 1, \dots, n\}$, $\alpha \in (0, 1]$, $\beta \in (0, 1]$ satisfying $\alpha^k = \beta^{n-k}$.

The weights are chosen as
 $1, \alpha, \alpha^2, \alpha^3, \dots, \alpha^k = \beta^{n-k}, \beta^{n-k-1}, \dots, \beta, 1$.

Our result using the trace distance with the weights

Theorem

Let $k \in \{0, 1, \dots, n\}$ be an integer. Let α and β be non-negative real numbers that satisfy $p_1 \alpha^k = p_2 \beta^{n-k}$. Let τ_\diamond^1 and τ_\diamond^2 be


$$\tau_\diamond^1 := \max_{\rho_{AR}} \left\| \left(\Phi_A^{(1)} - \alpha \Phi_A^{(2)} \right) (\rho_{AR}) \right\|_1,$$
$$\tau_\diamond^2 := \max_{\rho_{AR}} \left\| \left(\beta \Phi_A^{(1)} - \Phi_A^{(2)} \right) (\rho_{AR}) \right\|_1.$$

The following then holds:

$$p_{\text{err}}(n) \geq \frac{1}{2} \left[1 - p_1 \left(\sum_{i=0}^{k-1} \alpha^i \right) \tau_\diamond^1 - p_2 \left(\sum_{i=0}^{n-k-1} \beta^i \right) \tau_\diamond^2 \right],$$

for arbitrary α , β and, k .

Numerical results



Quantum channel group discrimination

Quantum channel **group** discrimination:

Problem to find a **group** including the channel (group may intersect).

We also derive lower bounds of the error probability for the quantum channel **group** discrimination problem using the **Bures angle** and the **trece distance with weight**.

The lower bound using Bures angle proves the **optimality of Grover's search** (This was proven for the case with exactly one marked element by [Zalka 1999]).

The lower bound using the trace distance with weight is numerically better than previously known lower bounds for some cases.