

# Aerobic anoxygenic phototrophic bacteria in Antarctic sea ice and seawater

Eileen Y. Koh,<sup>††</sup> William Phua<sup>‡</sup> and Ken G. Ryan

School of Biological Sciences, Victoria University of Wellington, PO Box 600, Wellington 6140, New Zealand.

## Summary

**Aerobic anoxygenic phototrophs are obligate aerobes with unusually high concentrations of carotenoids, low cellular contents of bacteriochlorophyll-*a* and they lack light-harvesting complex II. In this study, sea ice and seawater samples were collected from six different sites in the Ross Sea, Antarctica. Using a combination of primers for *pufM* (which encodes a pigment-binding protein subunit of the reaction centre complex), clone libraries of DNA and cDNA were created and a total of 63 positive clones were obtained from three sites, all clustering within the  $\alpha$ -*Proteobacteria*. Fifty-three of these clones were from seawater. The remaining clones were from sea ice and all were found in the middle and bottom sections of the ice. These sea ice bacteria may favour the lower part of the ice matrix where irradiance is low. This report highlights the first findings of AAnPs in antarctic sea ice and seawater within the Ross Sea Region.**

## Introduction

Recent large-scale environmental metagenomic surveys have revealed a wide distribution and abundance of bacterial phototrophic genes throughout the world's oceans (Venter *et al.*, 2004; Rusch *et al.*, 2007; Yutin *et al.*, 2007). Until 30 years ago when the first aerobic anoxygenic phototroph (AAnP) was reported (Shiba *et al.*, 1979), these phototrophic bacteria had been thought to be strictly anaerobic. Since 1979, the AAnPs have been found to represent a functional group of bacteria, defined by their aerobic, heterotrophic and phototrophic abilities. AAnPs are obligate aerobes with unusually high concentrations

of carotenoids, low cellular contents of bchl-*a* and they lack light-harvesting complex II (Kolber *et al.*, 2001; Rathgeber *et al.*, 2004). Given their larger cellular size and faster growth rates compared with the other heterotrophic bacteria (Schwalbach and Fuhrman, 2005; Jiao *et al.*, 2007; Lami *et al.*, 2007), AAnP might contribute up to 50% of the total bacterial production in oligotrophic waters and thus are a large sink for dissolved organic matter (Koblizek *et al.*, 2007). In fact, we still do not understand their contribution to the carbon cycle, their true abundance and their distribution in the global marine environment (Kolber *et al.*, 2000; 2001; Béjà *et al.*, 2002; Jiao *et al.*, 2010).

A part of the photosynthetic gene cluster in AAnPs is a 45 kilobase photosynthetic unit forming (*puf*) super-operon that codes for the reaction centre structural proteins (Béjà *et al.*, 2002; Tank *et al.*, 2009). This contiguous super-operon may be located on the bacterial chromosome or on extra-chromosomal elements (Pradella *et al.*, 2004). In AAnP, the expression of the *puf* operon is insensitive to oxygen (Nishimura *et al.*, 1996; Masuda *et al.*, 2000). One of these genes, *pufM* has been frequently used as a marker in the investigation of diversity in AAnPs (Béjà *et al.*, 2002; Lambeck *et al.*, 2002; Hu *et al.*, 2006; Waidner and Kirchman, 2008; Cottrell and Kirchman, 2009; Jiao *et al.*, 2010).

During ice-covered seasons in the Southern Ocean, light levels in the water columns are governed by the thickness and transparency of sea ice (Becquevort *et al.*, 2009). Microbes growing near the surface of sea ice experience very low temperatures, and high salinities and light levels that together comprise physical conditions close to the limits of life. Lower sections of sea ice are characterized by low light intensity (McMinn *et al.*, 2003), stable temperatures and readily available nutrients (Thomas and Dieckmann, 2002; Lazzara *et al.*, 2007; Ralph *et al.*, 2007). The highest numbers of cells, both algal and bacterial, are therefore often found at the bottom 10 cm of the ice (McMinn *et al.*, 1999; 2010; Lizotte, 2003; Ryan *et al.*, 2011). The microbial communities in sea ice exhibit significantly higher single-cell metabolic activity than those in many other marine ecosystems despite the extreme environment (Junge *et al.*, 2004; Martin *et al.*, 2008; 2009), and they are highly diverse (Brown and Bowman, 2001; Brinkmeyer *et al.*, 2003; Murray and Grzymalski, 2007; Bowman *et al.*, 2011). Very recently, the presence of

Received 9 March, 2011; revised 3 August, 2011; accepted 9 August, 2011. \*For correspondence. E-mail koh\_eileen@yahoo.com; Tel. (+65) 65166666; Fax (+65) 67791635. Present addresses: <sup>†</sup>Department of Civil and Environmental Engineering, The National University of Singapore, Block E1A #07-03, no. 1 Engineering Drive 2, Singapore 117576, Singapore; <sup>‡</sup>National Junior College, 37 Hillcrest Road, Singapore 288913, Singapore.

**Table 1.** Primers used in this study.

Primer	Sequence (5'→3')	Target gene	Reference
<i>puf</i> M.557F	CGCACCTGGACTGGAC	<i>pufM</i>	Achenbach <i>et al.</i> (2001)
<i>puf</i> M.750R	CCCATGGTCCAGCGCCAGAA		
<i>puf</i> M_uniF	GGNAAYYTNTWYTAYAAAYCCNTTYCA		Yutin <i>et al.</i> (2005)
<i>puf</i> M_uniR	YCCATNGTCCANCKCCARAA		
<i>puf</i> M_WAW <sup>R</sup>	AYNGCRAACCACCANGCCCA		

AAnPs were amplified from both DNA and cDNA extracts using *pufM* primers (Table 1), as recommended by Achenbach and colleagues (2001) and Yutin and colleagues (2005), and in combinations. Amplification conditions were at 94°C (5 min), 40 cycles of 94°C (45 s), 53.1°C (1 min) and 72°C (1 min), with a final extension at 72°C for 5 min.

F, forward; R, reverse.

Archaea has been reported in both the Arctic (Collins *et al.*, 2010) and Antarctic sea ice (Cowie *et al.*, 2011), and the newly described copiotrophic *Coraliomargarita* was found in sufficient abundance to suggest a niche occupation in sea ice (Bowman *et al.*, 2011). In fact, substantial research has been conducted on the ecology and biochemistry of sea ice (Krembs *et al.*, 2002; 2011; Collins *et al.*, 2008).

The majority of the sea ice bacterial research reported to date has focused on the heterotrophs (Delille, 2004; Martin *et al.*, 2009; Moreau *et al.*, 2010; Piquet *et al.*, 2011). Little attention has been placed on novel phototrophs such as AAnPs (Karr *et al.*, 2003); however, some other phototrophs have been found in Antarctic lakes and in the Southern Ocean (Giebel *et al.*, 2009) and we have recently described the presence of proteorhodopsin bacteria in sea ice communities (Koh *et al.*, 2010). Here, the *pufM* gene was used to describe for the first time the distribution of AAnP in the Antarctic seawater and ice environment.

Seawater was collected from the Ross Sea through the Scott Base water supply at ~ 5 m depth in a 20 l carboy, and transported back to New Zealand for filtration and further analyses. After thawing at 4°C, approximately 10 l of seawater was first filtered through a 3.0 µm nitrocellulose membrane filter (Millipore, USA) before passing the filtrate through a 0.22 µm mixed nitrocellulose acetate membrane filter (Pall Life Sciences, USA). This filtration was done in triplicates and both 3.0 and 0.22 µm filter papers were stored separately in a sterile 50 ml Falcon tube (BD Biosciences, USA) with sufficient 1 × Phosphate Buffer Solution (PBS) added to submerge the filter at the end of the filtration. The tubes were vortexed and 1 ml of suspension was transferred to a sterile microcentrifuge tube. The microcentrifuge tube was spun at 8000 *g* for 10 cycles of 5 min each (Eppendorf, Germany). The supernatant was then decanted before adding another aliquot of cell suspension. A total of 10 ml was taken from each tube to be used as starting material for subsequent DNA extraction using the High Pure PCR Template Preparation Kit (Roche Diagnostics GmbH, Germany), according to the manufacturer's instructions.


Sea ice core samples were collected at five sites across the Victoria Land Coast. All procedures pertaining to samples collection, processing, nucleic acids preparation and extraction were carried out as previously described (Koh *et al.*, 2010).

In this study, *pufM* primers targeting the *bchl-a*-containing bacteria published by Achenbach and colleagues (2001) and Yutin and colleagues (2005) were used to screen the sample set (Table 1). These primers are specific to the *pufM* gene in the light-harvesting reaction centre of the phototrophic *Proteobacteria*, and they provide a qualitative estimate of the diversity of bacteria with this gene in the environment (Achenbach *et al.*, 2001).

## Results and discussion

The goal of this study was to demonstrate the presence of *bchl-a* based phototrophy in the antarctic seawater and sea ice environment. No PCR products were detected after amplification with *pufM* primers from Yutin and colleagues (2005). Only the *pufM* primers from Achenbach and colleagues (2001) gave positive PCR amplicons and 30 plasmid clones from each sample were chosen for further *EcoRI*-digestion before sequencing. Sixty-three positive *pufM* clones from seawater (53) and sea ice (10) formed two clusters that showed phylogenetic relatedness to the *α-Proteobacteria* AAnP and none to the *β*- and *γ-Proteobacteria* (Fig. 1). This observation is similar to the Antarctic Lake Fryxell study by Karr and colleagues (2003) who also found an ice-bound community of AAnPs dominated by *α-Proteobacteria*. The similarity of the sea ice and seawater derived clones likely reflects the fact that sea ice is derived from seawater (Thomas and Dieckmann, 2002).

Within cluster A, 29 DNA clones from the sea water were grouped with the cDNA clones from sea ice and had 100% bootstrap support with minimal genetic distance between them (Fig. 1). Clones from TNB\_Brine\_CG200m65, on the other hand, were genetically distinct from the seawater DNA and other sea ice cDNA clones. Even within the TNB\_Brine\_CG200m65 clones, the



**Fig. 1.** Neighbour-joining tree based on the ~270 bp nucleotide sequence of the DNA fragment coding for the *putM* gene. ● = seawater DNA; ▲ = DNA; △ = cDNA. 3 K ≥ 3.0 µm filter; 22 K and F ≤ 3.0 µm; CE = Cape Evans; CR = Cape Roberts; TNB = Terra Nova Bay; SB SW = Scott Base Seawater. CG200m65 = Brine collected at Campbell Glacier at 200 m distance 65 cm depth. The phylogeny of the DNA sequences was constructed in MEGA4.0 (Tamura *et al.*, 2007) based on neighbour-joining algorithms with 1000 bootstrap replicates and *p*-distance as substitution model. Bootstrap values ≥ 50% are presented at the corresponding nodes. Green non-sulfur bacterium *Chloroflexus aurantiacus* (X07847) was used as out-group. The sequences reported in this study have been deposited in GenBank under the following accession numbers HQ142788-HQ142850.

genetic substitution rate was at 0.03 and the branch was supported by a 66% bootstrap value. Clones in cluster A have 97% bootstrap values to non-cultured *putM* relatives from the Delaware estuary (Waidner and Kirchman, 2008) and Lake Fryxell in Antarctica (Karr *et al.*, 2003), and also well-supported bootstrap values to environmental *putM*-containing *Roseobacter* (Oz *et al.*, 2005) and *Alphaproteobacterium* R2A62 (Béjà *et al.*, 2002).

Cluster B contained DNA clones from seawater and the same TNB\_Brine\_CG200m65 sample from Cluster A (Fig. 1). All of the sea-ice-derived sequences in this cluster were from brine collected in the middle portion of the sea ice. There was 100% bootstrap support to *Roseobacter denitrificans* and 99% to *Erythrobacter longus*, both of which are  $\alpha$ -*Proteobacteria* AAnPs from cultured samples (Fig. 1), implying that clones in both clusters could be *Roseobacter*-clade affiliated (RCA). *Roseobacter* are well represented across diverse marine habitats (reviewed by Buchan *et al.*, 2005; Wagner-Dobler and Biebl, 2006) and RCA bacteria in particular, have been observed to constitute about 20% of the total bacterial

community in the Southern Ocean (Selje *et al.*, 2004; Giebel *et al.*, 2009; 2011). Biodiversity studies based on the 16S rDNA gene conducted on sea ice from other Antarctic study sites within our research group have also yielded clones with high similarities to *Roseobacter* (E.W. Maas, A.M. Simpson, A.R. Martin, K.G. Ryan and R. O'Toole, unpubl. data) and *Erythrobacter* sp. (R.O.M. Cowie, E.W. Maas and K.G. Ryan, unpubl. data).

The 10 positive clones from sea ice were extracted from three different sites separated by ~300 km: Cape Evans (CE; 1), Cape Roberts (CR; 2) and Terra Nova Bay (TNB; 7). Three of these were of cDNA origin (represented by open triangles in Fig. 1) and seven of DNA origin (filled triangles in Fig. 1). Only two clones were from the bottom sea ice at CR, while the rest were from the middle portion. No *putM* clones were observed from the top portion of the sea ice from any location.

Most of the sea ice AAnPs were detected in mid sections of the ice, where the salinity was more than twice (~87‰) that of seawater. Only two clones were found in bottom sections of the ice where the salinity matches that

of the seawater. While some AAnP bacteria have been reported in Canadian hypersaline lakes (Csotonyi *et al.*, 2008) and saline lakes on the Tibetan Plateau (Jiang *et al.*, 2009; 2010), in other studies, the diversity of the *puM* gene is inversely proportional to salinity (Jiang *et al.*, 2009). Waidner and Kirchman (2008) and Jiang and colleagues (2010) have reported that high salinity reduced AAnP bacterial community diversity and composition, and Giebel and colleagues (2009) noted the inverse correlation of salinity and RCA cluster abundance from the subtropics to the Southern Ocean. The role of salinity in AAnP distribution is equivocal at present and more studies in hypersaline environments such as sea ice are needed.

One important factor that contributes to the successful competitive adaption of AAnP bacteria to the environment may be detritus particle attachment (Waidner and Kirchman, 2008; Lami *et al.*, 2009; Cottrell *et al.*, 2010), as observed in the Delaware estuary where AAnP attached to detritus particles accounted for > 10% of the total prokaryotic numbers (Waidner and Kirchman, 2008). In sea ice, detritus particles are distributed throughout the ice column (Sullivan and Palmisano, 1984), and these could serve as an attachment surface for AAnP. *Roseobacter* spp., for instance, are known to encode genes that support attachment (Slightom and Buchan, 2009). Thus, the sea ice and seawater clones in this study that clustered close to RCA sequences (Fig. 1) might favour a particle attachment lifestyle, via sea ice crystals and dendritus particles.

Marine  $\gamma$ -*Proteobacteria* with AAnP genes have also been reported recently (Hu *et al.*, 2006; Cho *et al.*, 2007; Fuchs *et al.*, 2007; Spring *et al.*, 2009). However, no *puM*-containing clones from the  $\gamma$ -*Proteobacteria* group were found in this study, although we have found other  $\gamma$ - and  $\alpha$ -*Proteobacteria* phototrophs such as proteorhodopsin-bearing bacteria in antarctic sea ice in a separate study (Koh *et al.*, 2010). These findings suggest that phototrophic *Proteobacteria* may have varied strategies to obtain energy for cellular needs, either using *bchl-a* (Shiba and Harashima, 1986; Kolber *et al.*, 2000) or proteorhodopsin (Béjà *et al.*, 2000; Koh *et al.*, 2010) to supplement their heterotrophic existence.

Metabolic gene probes, such as the *puM* gene, have several advantages over rDNA-based probes, in that they provide an association to an exclusive physiology in a phylogenetically diverse group. And when coupled with mRNA detection techniques, phototrophic metabolism can be inferred. Reverse-transcription PCR was used to produce cDNA from mRNA transcripts stored in RNAlater to determine whether *puM* was being transcribed at the time of sampling. In this study, only three cDNA transcripts from cluster A were apparently actively transcribing *puM* in the sea ice at the time of sampling. Given that the *puf* operon of AAnP is strongly repressed by high irradiance (Rathgeber *et al.*, 2004; Yurkov and Csotonyi, 2009), and

**Table 2.** Distribution of brine, sea ice and seawater clones within each clonal group.

Clonal group	Seawater DNA	Sea ice cDNA	Brine	Total
A	30	3	3	36
B	23	0	4	27
Total	53	3	7	63

All clones derived were derived from amplification with *puM* primers reported by Achenbach and colleagues (2001). The seawater analysed in this study was collected from the sea water supply line at Scott Base (SB SW). DNA was extracted using the High Pure PCR Template Preparation Kit (Roche Diagnostics GmbH, Germany), according to the manufacturer's instructions (S1). The collection and DNA extraction of sea ice and brine were carried out as documented in Koh and colleagues (2010).

that our samples were collected near the mid-austral summer, it is possible that many sea ice AAnP bacteria were not active at the time of collection, and instead have later or earlier onset of photosynthetic activity when ambient light levels are lower. In addition, this could explain why *puM* cDNA transcripts were only detected from the mid-bottom ice matrix and not near the surface where irradiance can be up to a thousand-fold higher than at the bottom of the ice (Buckley and Trodahl, 1987; Harashima *et al.*, 1987; Yurkov and Beatty, 1998). In addition, the higher amount of nutrients present in the underlying seawater (Thomas and Dieckmann, 2002) may further support AAnP growth. More abundance data from polar marine environments of AAnP are urgently required.

Schwalbach and Fuhrman (2005) reported that in the Western Antarctic, AAnP only comprised ~ 0.05% of the total bacterial abundance, while Cottrell and Kirchman (2009) reported up to 8% in the Arctic Ocean. In this study, we observed a higher number of clones from seawater than from the sea ice (Table 2) and this may indicate that at least in ice-covered regions of the Ross Sea region, AAnPs are more abundant in the open water than in the ice although further quantitative work is needed to confirm this observation.

## Conclusion

Using a combination of standard and reverse-transcription PCR to screen antarctic sea ice and seawater samples collected over 300 km apart in the Ross Sea region, we present the first report of AAnP bacteria in antarctic seawater and sea ice. The seawater clones were closely grouped to previously identified environmental clones and to *Roseobacter dentrificans*, with the latter suggesting presence of the RCA cluster in the Ross Sea region. The sea ice clones, on the other hand, exhibited stronger grouping to environmental clones than to cultured representatives. Not only did we find that the majority of the *bchl-a*-bearing bacteria were found in the middle section of



the sea ice and within the water column, we also observed cDNA regenerated from *puM*-RNA in sea ice bacterial cells, which suggests that the cells are actively synthesizing the *bchl-a* gene within this extreme ecosystem.

Bacteria in sea ice are highly metabolically active (Martin *et al.*, 2008) and perhaps bacterial phototrophy provides a selective advantage in such extreme conditions. Microbial energetics within the sea ice is still poorly understood and the ecological importance of the AAnPs in the sea ice environment needs further verification to provide a better understanding of their functions and their contributions to the sea ice 'microbial loop'.

## Acknowledgements

The authors thank Event K043 for their help with field sampling during the field seasons of 2007 and 2008, and acknowledge the logistical and field support of Antarctica New Zealand, in particular S. Gordon, Project Manager of the Latitudinal Gradient Project (LGP). We thank Bex Cowie for the preparation of the cDNAs. We also thank both anonymous reviewers for their suggestions to improve the manuscript. This research was supported by the New Zealand Foundation of Research Science and Technology Grant number VICX0706 to KGR, the VUW SFRG Grants 103167/2585 and 96429/2661 to EYK and WP respectively. The authors declare no conflict of interests.

## References

- Achenbach, L.A., Carey, J., and Madigan, M.T. (2001) Photosynthetic and Phylogenetic Primers for Detection of Anoxygenic Phototrophs in Natural Environments. *Appl Environ Microbiol* **67**: 2922–2926.
- Becquevort, S., Dumont, I., Tison, J.L., Lannuzel, D., Sauvee, M.L., Chou, L., and Schoemann, V. (2009) Biogeochemistry and microbial community composition in sea ice and underlying seawater off East Antarctica during early spring. *Polar Biol* **32**: 879–895.
- Béjå, O., Aravind, L., Koonin, E.V., Suzuki, M.T., Hadd, A., Nguyen, L.P., *et al.* (2000) Bacterial rhodopsin: evidence for a new type of phototrophy in the sea. *Science* **289**: 1902–1906.
- Béjå, O., Suzuki, M.T., Heidelberg, J.F., Nelson, W.C., Preston, C.M., Hamada, T., *et al.* (2002) Unsuspected diversity among marine aerobic anoxygenic phototrophs. *Nature* **415**: 630–633.
- Bowman, J.S., Rasmussen, S., Blom, N., Deming, J.W., Rysgaard, S., and Sicheritz-Ponten, T. (2011) Microbial community structure of Arctic multiyear sea ice and surface seawater by 454 sequencing of the 16S RNA gene. *ISME J* doi:10.1038/ismej.2011.76.
- Brinkmeyer, R., Knittel, K., Jurgens, J., Weyland, H., Amann, R., and Helmke, E. (2003) Diversity and structure of bacterial communities in Arctic versus Antarctic pack ice. *Appl Environ Microbiol* **69**: 6610–6619.
- Brown, M.V., and Bowman, J.P. (2001) A molecular phylogenetic survey of sea-ice microbial communities. *FEMS Microbiol Ecol* **35**: 267–275.
- Buchan, A., Gonzalez, J.M., and Moran, M.A. (2005) Overview of the Marine Roseobacter Lineage. *Appl Environ Microbiol* **71**: 5665–5677.
- Buckley, R.G., and Trodahl, H.J. (1987) Scattering and absorption of visible light by sea ice. *Nature* **326**: 867–869.
- Cho, J.-C., Stapels, M.D., Morris, R.M., Vergin, K.L., Schwalbach, M.S., Givan, S.A., *et al.* (2007) Polyphyletic photosynthetic reaction centre genes in oligotrophic marine Gammaproteobacteria. *Environ Microbiol* **9**: 1456–1463.
- Collins, R.E., Carpenter, S.D., and Deming, J.W. (2008) Spatial heterogeneity and temporal dynamics of particles, bacteria, and pEPS in Arctic winter sea ice. *J Mar Syst* **74**: 902–917.
- Collins, R.E., Rocap, G., and Deming, J.W. (2010) Persistence of bacterial and archaeal communities in sea ice through an Arctic winter. *Environ Microbiol* **12**: 1828–1841.
- Cottrell, M.T., and Kirchman, D.L. (2009) Photoheterotrophic Microbes in the Arctic Ocean in Summer and Winter. *Appl Environ Microbiol* **75**: 4958–4966.
- Cottrell, M.T., Ras, J., and Kirchman, D.L. (2010) Bacteriochlorophyll and community structure of aerobic anoxygenic phototrophic bacteria in a particle-rich estuary. *ISME J* **4**: 945–954.
- Cowie, R.O.M., Maas, E.W., and Ryan, K.G. (2011) Archaeal diversity revealed in Antarctic sea ice. *Antarct Sci* doi:10.1017/S0954102011000368.
- Csotonyi, J., Swiderski, J., Stackebrandt, E., and Yurkov, V. (2008) Novel halophilic aerobic anoxygenic phototrophs from a Canadian hypersaline spring system. *Extremophiles* **12**: 529–539.
- Delille, D. (2004) Abundance and function of bacteria in the Southern Ocean. *Cell Mol Biol* **50**: 543–551.
- Fuchs, B.M., Spring, S., Teeling, H., Quast, C., Wulf, J., Schattenhofer, M., *et al.* (2007) Characterization of a marine gammaproteobacterium capable of aerobic anoxygenic photosynthesis. *Proc Natl Acad Sci USA* **104**: 2891–2896.
- Giebel, H.A., Brinkhoff, T., Zwisler, W., Selje, N., and Simon, M. (2009) Distribution of *Roseobacter* RCA and SAR11 lineages and distinct bacterial communities from the subtropics to the Southern Ocean. *Environ Microbiol* **11**: 2164–2178.
- Giebel, H.-A., Kalhoefer, D., Lemke, A., Thole, S., Gahl-Janssen, R., Simon, M., and Brinkhoff, T. (2011) Distribution of *Roseobacter* RCA and SAR11 lineages in the North Sea and characteristics of an abundant RCA isolate. *ISME J* **5**: 8–19.
- Harashima, K., Kawazoe, K., Yoshida, I., and Kamata, H. (1987) Light-stimulated aerobic growth of *Erythrobacter* species Och-114. *Plant Cell Physiol* **28**: 365–374.
- Hu, Y.H., Du, H.L., Jiao, N.Z., and Zeng, Y.H. (2006) Abundant presence of the gamma-like Proteobacterial *puM* gene in oxic seawater. *FEMS Microbiol Lett* **263**: 200–206.
- Jiang, H., Dong, H., Yu, B., Lv, G., Deng, S., Wu, Y., *et al.* (2009) Abundance and diversity of aerobic anoxygenic phototrophic bacteria in saline lakes on the Tibetan plateau. *FEMS Microbiol Ecol* **67**: 268–278.

- Jiang, H., Deng, S., Huang, Q., Dong, H., and Yu, B. (2010) Response of aerobic anoxygenic phototrophic bacterial diversity to environment conditions in saline lakes and Daotang River on the Tibetan Plateau, NW China. *Geomicrobiol J* **27**: 400–408.
- Jiao, N.Z., Zhang, Y., Zeng, Y.H., Hong, N., Liu, R.L., Chen, F., and Wang, P.X. (2007) Distinct distribution pattern of abundance and diversity of aerobic anoxygenic phototrophic bacteria in the global ocean. *Environ Microbiol* **9**: 3091–3099.
- Jiao, N.Z., Zhang, F., and Hong, N. (2010) Significant roles of bacteriochlorophylla supplemental to chlorophylla in the ocean. *ISME J* **4**: 595–597.
- Junge, K., Eicken, H., and Deming, J.W. (2004) Bacterial activity at  $-2$  to  $-20^{\circ}\text{C}$  in Arctic wintertime sea ice. *Appl Environ Microbiol* **70**: 550–557.
- Karr, E.A., Sattley, W.M., Jung, D.O., Madigan, M.T., and Achenbach, L.A. (2003) Remarkable diversity of phototrophic purple bacteria in a permanently frozen Antarctic lake. *Appl Environ Microbiol* **69**: 4910–4914.
- Koblizek, M., Masin, M., Ras, J., Poulton, A.J., and Prasil, O. (2007) Rapid growth rates of aerobic anoxygenic phototrophs in the ocean. *Environ Microbiol* **9**: 2401–2406.
- Koh, E.Y., Atamna-Ismaeel, N., Martin, A., Cowie, R.O.M., Béjà, O., Davy, S.K., *et al.* (2010) Proteorhodopsin-bearing bacteria in Antarctic sea ice. *Appl Environ Microbiol* **76**: 5918–5925.
- Kolber, Z.S., van Dover, C.L., Niederman, R.A., and Falkowski, P.G. (2000) Bacterial photosynthesis in surface waters of the open ocean. *Nature* **407**: 177–179.
- Kolber, Z.S., Plumley, F.G., Lang, A.S., Beatty, J.T., Blankenship, R.E., VanDover, C.L., *et al.* (2001) Contribution of aerobic photoheterotrophic bacteria to the carbon cycle in the ocean. *Science* **292**: 2492–2495.
- Krembs, C., Eicken, H., Junge, K., and Deming, J.W. (2002) High concentrations of exopolymeric substances in Arctic winter sea ice: implications for the polar ocean carbon cycle and cryoprotection of diatoms. *Deep Sea Res I* **49**: 2163–2181.
- Krembs, C., Eicken, H., and Deming, J.W. (2011) Exopolymer alteration of physical properties of sea ice and implications for ice habitability and biogeochemistry in a warmer Arctic. *Proc Natl Acad Sci USA* **108**: 3653–3658.
- Lambeck, K., Esat, T.M., and Potter, E.K. (2002) Links between climate and sea levels for the past three million years. *Nature* **419**: 199–206.
- Lami, R., Cottrell, M.T., Ras, J., Ulloa, O., Obernosterer, I., Claustre, H., *et al.* (2007) High abundances of aerobic anoxygenic photosynthetic bacteria in the South Pacific Ocean. *Appl Environ Microbiol* **73**: 4198–4205.
- Lami, R., Cottrell, M.T., Campbell, B.J., and Kirchman, D.L. (2009) Light-dependent growth and proteorhodopsin expression by *Flavobacteria* and SAR11 in experiments with Delaware coastal waters. *Environ Microbiol* **11**: 3201–3209.
- Lazzara, L., Nardello, I., Ermanni, C., Mangoni, O., and Saggiomo, V. (2007) Light environment and seasonal dynamics of microalgae in the annual sea ice at Terra Nova Bay, Ross Sea, Antarctica. *Antarct Sci* **19**: 83–92.
- Lizotte, M.P. (2003) Sea ice – an introduction to its physics, chemistry, biology and geology. In *The Microbiology of Sea Ice*. Thomas, D.N., and Dieckmann, G.S. (eds). Oxford, UK: Blackwell, pp. 184–210.
- McMinn, A., Ashworth, C., and Ryan, K. (1999) Growth and productivity of Antarctic sea ice algae under PAR and UV irradiances. *Bot Mar* **42**: 401–407.
- McMinn, A., Ryan, K.G., and Gademann, R. (2003) Diurnal changes in photosynthesis of Antarctic fast ice algal communities determined by pulse amplitude modulation fluorometry. *Mar Biol* **143**: 359–367.
- McMinn, A., Pankowski, A., Ashworth, C., Bhagooli, R., Ralph, P., and Ryan, K. (2010) *In situ* net primary productivity and photosynthesis of Antarctic sea ice algal, phytoplankton and benthic algal communities. *Mar Biol* **157**: 1345–1356.
- Martin, A., Hall, J.A., O'Toole, R., Davy, S.K., and Ryan, K.G. (2008) High single-cell metabolic activity in Antarctic sea ice bacteria. *Aquat Microb Ecol* **52**: 25–31.
- Martin, A., Hall, J., and Ryan, K. (2009) Low Salinity and High-Level UV-B Radiation Reduce Single-Cell Activity in Antarctic Sea Ice Bacteria. *Appl Environ Microbiol* **75**: 7570–7573.
- Masuda, S., Nagashima, K.V.P., Shimada, K., and Matsuura, K. (2000) Transcriptional control of expression of genes for photosynthetic reaction center and light-harvesting proteins in the purple bacterium *Rhodovulum sulfidophilum*. *J Bacteriol* **182**: 2778–2786.
- Moreau, S., Ferreyra, G.A., Mercier, B., Lemarchand, K., Lionard, M., Roy, S., *et al.* (2010) Variability of the microbial community in the western Antarctic Peninsula from late fall to spring during a low ice cover year. *Polar Biol* **33**: 1599–1614.
- Murray, A.E., and Grzyski, J.J. (2007) Diversity and genomics of Antarctic marine micro-organisms. *Philos Trans R Soc Lond B* **362**: 2259–2271.
- Nishimura, K., Shimada, H., Ohta, H., Masuda, T., Shioi, Y.Z., and Takamiya, K. (1996) Expression of the *puf* operon in an aerobic photosynthetic bacterium, *Roseobacter denitrificans*. *Plant Cell Physiol* **37**: 153–159.
- Oz, A., Sabehi, G., Koblizek, M., Massana, R., and Béjà, O. (2005) *Roseobacter*-like bacteria in Red and Mediterranean Sea aerobic anoxygenic photosynthetic populations. *Appl Environ Microbiol* **71**: 344–353.
- Piquet, A.M.-T., Bolhuis, H., Meredith, M.P., and Buma, A.G.J. (2011) Shifts in coastal Antarctic marine microbial communities during and after melt water-related surface stratification. *FEMS Microbiol Ecol* **76**: 413–427.
- Pradella, S., Allgaier, M., Hoch, C., Pauker, O., Stackebrandt, E., and Wagner-Dobler, I. (2004) Genome organization and localization of the *puLM* genes of the photosynthesis reaction center in phylogenetically diverse marine Alphaproteobacteria. *Appl Environ Microbiol* **70**: 3360–3369.
- Ralph, P.J., Ryan, K.G., Martin, A., and Fenton, G. (2007) Melting out of sea ice causes greater photosynthetic stress in algae than freezing in. *J Phycol* **43**: 948–956.
- Rathgeber, C., Beatty, J.T., and Yurkov, V. (2004) Aerobic phototrophic bacteria: new evidence for the diversity, ecological importance and applied potential of this previously overlooked group. *Photosynth Res* **81**: 113–128.
- Rusch, D.B., Halpern, A.L., Sutton, G., Heidelberg, K.B., Williamson, S., Yooseph, S., *et al.* (2007) *The Sorcerer II*

- Global Ocean Sampling Expedition: Northwest Atlantic through Eastern Tropical Pacific. *PLoS Biol* **5**: e77.
- Ryan, K.G., Tay, M.L., Martin, A., McMin, A., and Davy, S.K. (2011) Chlorophyll fluorescence imaging analysis of the responses of Antarctic bottom-ice algae to light and salinity during melting. *J Exp Mar Bio Ecol* **399**: 156–161.
- Schwalbach, M.S., and Fuhrman, J.A. (2005) Wide-ranging abundances of aerobic anoxygenic phototrophic bacteria in the world ocean revealed by epifluorescence microscopy and quantitative PCR. *Limnol Oceanogr* **50**: 620–628.
- Selje, N., Simon, M., and Brinkhoff, T. (2004) A newly discovered Roseobacter cluster in temperate and polar oceans. *Nature* **427**: 445–448.
- Shiba, T., and Harashima, K. (1986) Aerobic Photosynthetic Bacteria. *Microbiol Sci* **3**: 376–378.
- Shiba, T., Simidu, U., and Taga, N. (1979) Distribution of Aerobic Bacteria Which Contain Bacteriochlorophyll a. *Appl Environ Microbiol* **38**: 43–45.
- Slightom, R.N., and Buchan, A. (2009) Surface Colonization by Marine Roseobacters: Integrating Genotype and Phenotype. *Appl Environ Microbiol* **75**: 6027–6037.
- Spring, S., Lunsdorf, H., Fuchs, B.M., and Tindall, B.J. (2009) The Photosynthetic Apparatus and Its Regulation in the Aerobic Gammaproteobacterium *Congregibacter litoralis* gen. nov., sp nov. *PLoS ONE* **4**: e4866.
- Sullivan, C.W., and Palmisano, A.C. (1984) Sea ice microbial communities: distribution, abundance, and diversity of ice bacteria in McMurdo Sound, Antarctica, in 1980. *Appl Environ Microbiol* **47**: 788–795.
- Tamura, K., Dudley, J., Nei, M., and Kumar, S. (2007) MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol* **24**: 1596–1599.
- Tank, M., Thiel, V., and Imhoff, J.F. (2009) Phylogenetic relationship of phototrophic purple sulfur bacteria according to *pufl* and *puflM* genes. *Int Microbiol* **12**: 175–185.
- Thomas, D.N., and Dieckmann, G.S. (2002) Biogeochemistry of Antarctic sea ice. *Oceanogr Mar Biol* **40**: 143–169.
- Venter, J.C., Remington, K., Heidelberg, J.F., Halpern, A.L., Rusch, D., Eisen, J.A., et al. (2004) Environmental genome shotgun sequencing of the Sargasso Sea. *Science* **304**: 66–74.
- Wagner-Dobler, I., and Biebl, H. (2006) Environmental biology of the marine *Roseobacter* lineage. *Annu Rev Microbiol* **60**: 255–280.
- Waidner, L.A., and Kirchman, D.L. (2008) Diversity and distribution of ecotypes of the aerobic anoxygenic phototrophy gene *puflM* in the Delaware estuary. *Appl Environ Microbiol* **74**: 4012–4021.
- Yurkov, V., and Csotonyi, J.T. (2009) New light on aerobic anoxygenic phototrophs. In *The Purple Phototrophic Bacteria*. Hunter, C.N., Daldal, F., Thurnauer, M.C., and Beatty, J.T. (eds). New York, NY: Springer Science, pp. 31–55.
- Yurkov, W., and Beatty, J.T. (1998) Aerobic anoxygenic phototrophic bacteria. *Microbiol Mol Biol Rev* **62**: 695–724.
- Yutin, N., Suzuki, M.T., and Béjà, O. (2005) Novel primers reveal wider diversity among marine aerobic anoxygenic phototrophs. *Appl Environ Microbiol* **71**: 8958–8962.
- Yutin, N., Suzuki, M.T., Teeling, H., Weber, M., Venter, J.C., Rusch, D.B., and Béjà, O. (2007) Assessing diversity and biogeography of aerobic anoxygenic phototrophic bacteria in surface waters of the Atlantic and Pacific Oceans using the Global Ocean Sampling expedition metagenomes. *Environ Microbiol* **9**: 1464–1475.