

Generated by elijahsheridan on 29 March 2020, 07:03:48

This report has been generated automatically by Madanalysis 5.

Please cite:

E. Conte, B. Fuks and G. Serret, MadAnalysis 5, A User-Friendly Framework for Collider Phenomenology, Comput. Phys. Commun. 184 (2013) 222-256,

To contact us:

 ${\bf http://madanalysis.irmp.ucl.ac.be} \\ {\bf ma5team@iphc.cnrs.fr} \\$

arXiv:1206.1599 [hep-ph].

Contents Setup 2 2 1.1 Command history 1.2 Configuration 3 Datasets 4 2.1signal 4 2.2 $bg_vbf_0_100$ 4 2.3 $bg_vbf_100_200$ 4 2.4 $bg_vbf_200_400$ 5 $bg_vbf_400_600$ 2.55 $2.6 \quad \, \mathrm{bg_vbf_600_800}$ 6 $2.7 ext{ bg_vbf_}800_1200$ 6 bg_vbf_1200_1600 7 2.8 2.9 bg_vbf_1600_inf 7 $2.10 \ bg_dip_0_100$ 7 $2.11 \ \ \mathrm{bg_dip_100_200}$ 8 2.12 bg dip 200 4008 2.13 bg dip 400 600 9 9 $2.14 \ \ bg_dip_600_800$ $2.15 \ \ bg_dip_800_1200$ 9 $2.16 \ \ bg_dip_1200_1600$ 10 $2.17 \hspace{0.1in} bg_dip_1600_inf$ 10 Histos and cuts 11 3.1 Cut 1 11 3.2 Cut 2 12 Summary **13**

13

4.1 Cut-flow charts

1 Setup

1.1 Command history

```
ma5># set directory where running "./bin/ma5"; set lumi; define the signal significance
ma5>set main.currentdir = /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno # need to
change this directory path -> exit and type "pwd" to get the path
ma5>set main.lumi = 40.0
ma5># import samples -> change the path to the LHE file
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/axion_signal/-
axion_signal_gurrola_cuts_1MeV.lhe.gz as signal
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/vbf_diphoton_background_
merged_lhe/vbf_diphoton_background_ht_0_100_merged.lhe.gz as bg_vbf_0_100
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/vbf_diphoton_background_
merged_lhe/vbf_diphoton_background_ht_100_200_merged.lhe.gz as bg_vbf_100_200
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/vbf_diphoton_background_
merged_lhe/vbf_diphoton_background_ht_200_400_merged.lhe.gz as bg_vbf_200_400
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/vbf_diphoton_background_
merged_lhe/vbf_diphoton_background_ht_400_600_merged.lhe.gz as bg_vbf_400_600
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/vbf_diphoton_background_
merged_lhe/vbf_diphoton_background_ht_600_800_merged.lhe.gz as bg_vbf_600_800
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/vbf_diphoton_background_
merged_lhe/vbf_diphoton_background_ht_800_1200_merged.lhe.gz as bg_vbf_800_1200
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/vbf_diphoton_background_
merged_lhe/vbf_diphoton_background_ht_1200_1600_merged.lhe.gz as bg_vbf_1200_1600
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/vbf_diphoton_background_
merged_lhe/vbf_diphoton_background_ht_1600_inf_merged.lhe.gz as bg_vbf_1600_inf
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/diphoton_double_isr_back
merged_lhe/diphoton_double_isr_background_ht_0_100_merged.lhe.gz as bg_dip_0_100
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/diphoton_double_isr_back
merged_lhe/diphoton_double_isr_background_ht_100_200_merged.lhe.gz as bg_dip_100_200
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/diphoton_double_isr_back
merged_lhe/diphoton_double_isr_background_ht_200_400_merged.lhe.gz as bg_dip_200_400
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/diphoton_double_isr_back
merged_lhe/diphoton_double_isr_background_ht_400_600_merged.lhe.gz as bg_dip_400_600
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/diphoton_double_isr_back
merged_lhe/diphoton_double_isr_background_ht_600_800_merged.lhe.gz as bg_dip_600_800
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/diphoton_double_isr_back
merged_lhe/diphoton_double_isr_background_ht_800_1200_merged.lhe.gz as bg_dip_800_1200
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/diphoton_double_isr_back
merged_lhe/diphoton_double_isr_background_ht_1200_1600_merged.lhe.gz as bg_dip_1200_1600
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/diphoton_double_isr_back
merged_lhe/diphoton_double_isr_background_ht_1600_inf_merged.lhe.gz as bg_dip_1600_inf
ma5># define bg and signal samples
ma5>set signal.type = signal
ma5>set bg_vbf_0_100.type = background
ma5>set bg_vbf_100_200.type = background
ma5>set bg_vbf_200_400.type = background
ma5>set bg_vbf_400_600.type = background
ma5>set bg_vbf_600_800.type = background
```

```
ma5>set bg_vbf_800_1200.type = background
ma5>set bg_vbf_1200_1600.type = background
ma5>set bg_vbf_1600_inf.type = background
ma5>set bg_dip_0_100.type = background
ma5>set bg_dip_100_200.type = background
ma5>set bg_dip_200_400.type = background
ma5>set bg_dip_400_600.type = background
ma5>set bg_dip_600_800.type = background
ma5>set bg_dip_800_1200.type = background
ma5>set bg_dip_1200_1600.type = background
ma5>set bg_dip_1600_inf.type = background
ma5># define weights for the samples
ma5>#set sample_1.weight = 1
ma5>#set sample_2.weight = 1
ma5># a jet can be from a light quark or b quark
ma5>define jets = j
ma5>define e = e+ e-
ma5>define mu = mu+ mu-
ma5>define ta = ta+ ta-
ma5>define lept = e mu ta
ma5>define ax = 9000005
ma5># selections
ma5>select (sdETA(jets[1] jets[2]) > 3.6 or sdETA(jets[1] jets[2]) < -3.6) and M(jets[1] jets[2]) < -3.6) and M(jets[1] jets[2]) < -3.6)
jets[2]) > 1250
ma5>select PT(a[1]) > 500 and M(a[1] a[2]) > 400
ma5>submit analysis_tight_pta500_maa400
```

1.2 Configuration

- MadAnalysis version 1.6.33 (2017/11/20).
- Histograms given for an integrated luminosity of 40.0fb⁻¹.

2 Datasets

2.1 signal

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: signal events.

• Generated events: 1000000 events.

• Normalization to the luminosity: 4094+/- 2 events.

• Ratio (event weight): 0.0041.

| Path to the event file | Nr. of events | Cross section (pb) | Negative wgts (%) |
|------------------------------------|---------------|--------------------|-------------------|
| /Users/elijahsheridan/- | | | |
| $MG5_aMC_v2_6_5/-$ | | | |
| axion_pheno/- | 1000000 | 0.102 @ 0.028% | 0.0 |
| madgraph_data/axion_signal/- | | | |
| _axion_signal_gurrola_cuts_1MeV.ll | | | |

$2.2 \quad bg_vbf_0_100$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 1000000 events.

• Normalization to the luminosity: 12150+/- 24 events.

 \bullet Ratio (event weight): 0.012 $\,$.

| Path to the event file | Nr. of events | Cross section (pb) | Negative wgts (%) |
|--|---------------|--------------------|-------------------|
| /Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/- vbf_diphoton_background_data/- merged_lhe/- vbf_diphoton_background_ht 0 10 | 1000000 | 0.304 @ 0.19% | 0.0 |

$2.3 \quad \text{bg vbf } 100 \quad 200$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 965662 events.

 \bullet Normalization to the luminosity: 9695+/- 17 $\,$ events.

• Ratio (event weight): 0.01.

| Path to the event file | Nr. of events | Cross section (pb) | Negative wgts (%) |
|---------------------------------|---------------|--------------------|-------------------|
| /Users/elijahsheridan/- | | | |
| $MG5_aMC_v2_6_5/-$ | | | |
| $axion_pheno/madgraph_data/-$ | 067660 | 0.242 @ 0.17% | 0.0 |
| vbf_diphoton_background_data/- | 965662 | 0.242 @ 0.17% | 0.0 |
| merged_lhe/- | | | |
| vbf_diphoton_background_ht_100_ | | | |

$\mathbf{2.4} \quad \mathbf{bg_vbf_200_400}$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 984165 events.

 \bullet Normalization to the luminosity: 5413+/- 11 events.

• Ratio (event weight): 0.0055.

| Path to the event file | Nr. of events | Cross section (pb) | Negative wgts (%) |
|---------------------------------|---------------|--------------------|-------------------|
| /Users/elijahsheridan/- | | | |
| $MG5_aMC_v2_6_5/-$ | | | |
| $axion_pheno/madgraph_data/-$ | 004165 | 0.127 @ 0.207 | 0.0 |
| vbf_diphoton_background_data/- | 984165 | 0.135 @ 0.2% | 0.0 |
| $\mathrm{merged_lhe/-}$ | | | |
| vbf_diphoton_background_ht_200_ | | | |

$\mathbf{2.5} \quad \mathbf{bg_vbf_400_600}$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 1000000 events.

• Normalization to the luminosity: 986+/-2 events.

 \bullet Ratio (event weight): 0.00099 .

| Path to the event file | Nr. of events | Cross section (pb) | Negative wgts (%) |
|--|---------------|--------------------|-------------------|
| /Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/- vbf_diphoton_background_data/- merged_lhe/- vbf_diphoton_background_ht_400_ | 1000000 | 0.0247 @ 0.14% | 0.0 |

$2.6 \quad \mathrm{bg_vbf_600_800}$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 1000000 events.

• Normalization to the luminosity: 252+/- 1 events.

• Ratio (event weight): 0.00025.

| Path to the event file | Nr. of events | Cross section (pb) | Negative wgts (%) |
|---|---------------|--------------------|-------------------|
| /Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/- vbf_diphoton_background_data/- merged_lhe/- vbf_diphoton_background_ht 600 | 1000000 | 0.0063 @ 0.13% | 0.0 |

$2.7 \quad \mathrm{bg_vbf_800_1200}$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

 \bullet Generated events: 400839 events.

• Normalization to the luminosity: 114+/- 1 events.

 \bullet Ratio (event weight): 0.00028.

| Path to the event file | Nr. of events | Cross section (pb) | Negative wgts (%) |
|---------------------------------|---------------|--------------------|-------------------|
| /Users/elijahsheridan/- | | | |
| $MG5_aMC_v2_6_5/-$ | | | |
| $axion_pheno/madgraph_data/-$ | 400020 | 0.00287 @ 0.16% | 0.0 |
| vbf_diphoton_background_data/- | 400839 | 0.00207 @ 0.10% | 0.0 |
| merged_lhe/- | | | |
| vbf_diphoton_background_ht_800_ | | | |

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 953803 events.

• Normalization to the luminosity: 20+/- 1 events.

• Ratio (event weight): 2.1e-05 .

| Path to the event file | Nr. of events | Cross section (pb) | Negative wgts (%) |
|--|---------------|--------------------|-------------------|
| /Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/- vbf_diphoton_background_data/- merged_lhe/- vbf_diphoton_background_ht_1200 | 953803 | 0.000515 @ 0.16% | 0.0 |

2.9 bg vbf 1600 inf

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

 \bullet Generated events: 270148 $\,$ events.

• Normalization to the luminosity: 7+/-1 events.

• Ratio (event weight): 2.6e-05 .

| Path to the event file | Nr. of events | Cross section (pb) | Negative wgts (%) |
|--|---------------|--------------------|-------------------|
| /Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/- vbf_diphoton_background_data/- merged_lhe/- vbf_diphoton_background_ht 1600 | 270148 | 0.000191 @ 0.11% | 0.0 |

$2.10 \quad \text{bg dip } 0 \quad 100$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 1040000 events.

 \bullet Normalization to the luminosity: 2710847+/- 4614 events.

 \bullet Ratio (event weight): 2.6 - warning: please generate more events (weight larger than 1)!

| Path to the event file | Nr. of events | Cross section (pb) | Negative wgts (%) |
|--|---------------|--------------------|-------------------|
| /Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/- | 1040000 | 67.8 @ 0.17% | 0.0 |
| diphoton_double_isr_background_d merged_lhe/- diphoton_double_isr_background_h | 1040000 | 01.0 & 0.11/0 | 0.0 |

2.11 bg dip 100 200

- \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .
- Sample consisting of: background events.
- Generated events: 1040000 events.
- Normalization to the luminosity: 1095362+/- 1528 events.
- Ratio (event weight): 1.1 warning: please generate more events (weight larger than 1)!

| Path to the event file | Nr. of events | Cross section (pb) | Negative wgts (%) |
|---|---------------|--------------------|-------------------|
| /Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/- diphoton_double_isr_background_d merged_lhe/- diphoton_double_isr_background_l | 1040000 | 27.4 @ 0.14% | 0.0 |

$2.12 \quad \ \, \text{bg_dip_200_400}$

- \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .
- Sample consisting of: background events.
- Generated events: 1040000 events.
- Normalization to the luminosity: 239548+/- 414 events.
- Ratio (event weight): 0.23 .

| Path to the event file | Nr. of events | Cross section (pb) | Negative wgts (%) |
|--|---------------|--------------------|-------------------|
| /Users/elijahsheridan/- | | | |
| $MG5_aMC_v2_6_5/-$ | | | |
| $axion_pheno/madgraph_data/-$ | 1040000 | 5.99 @ 0.17% | 0.0 |
| diphoton_double_isr_background_d | 1040000 | 5.99 @ 0.1770 | 0.0 |
| merged_lhe/- | | | |
| $_diphoton_double_isr_background_l$ | | | |

$2.13 \quad bg_dip_400_600$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 1040000 events.

• Normalization to the luminosity: 28798+/- 53 events.

• Ratio (event weight): 0.028 .

| Path to the event file | Nr. of events | Cross section (pb) | Negative wgts (%) |
|----------------------------------|---------------|--------------------|-------------------|
| /Users/elijahsheridan/- | | | |
| $MG5_aMC_v2_6_5/-$ | | | |
| $axion_pheno/madgraph_data/-$ | 1040000 | 0.72 @ 0.18% | 0.0 |
| diphoton_double_isr_background_o | 1040000 | 0.72 @ 0.1670 | 0.0 |
| $merged_lhe/-$ | | | |
| diphoton_double_isr_background_l | | | |

$2.14 ext{ bg_dip_}600_800$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

 \bullet Generated events: 662009 events.

• Normalization to the luminosity: 6674+/- 28 events.

• Ratio (event weight): 0.01 .

| Path to the event file | Nr. of events | Cross section (pb) | Negative wgts (%) |
|--|---------------|--------------------|-------------------|
| /Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/- | 662009 | 0.167 @ 0.41% | 0.0 |
| diphoton_double_isr_background_d merged_lhe/- diphoton_double_isr_background_h | 002000 | 0.107 @ 0.4170 | 0.0 |

2.15 bg dip 800 1200

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 1040000 events.

 \bullet Normalization to the luminosity: 2942+/- 6 events.

 \bullet Ratio (event weight): 0.0028 % =0.0028 .

| Path to the event file | Nr. of events | Cross section (pb) | Negative wgts (%) |
|---|---------------|--------------------|-------------------|
| /Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/- diphoton_double_isr_background_d merged_lhe/- diphoton_double_isr_background_h | 1040000 | 0.0736 @ 0.17% | 0.0 |

2.16 bg dip 1200 1600

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 337115 events.

• Normalization to the luminosity: 513+/-3 events.

| Path to the event file | Nr. of events | Cross section (pb) | Negative wgts (%) |
|----------------------------------|---------------|--------------------|-------------------|
| /Users/elijahsheridan/- | | | |
| $MG5_aMC_v2_6_5/-$ | | | |
| $axion_pheno/madgraph_data/-$ | 337115 | 0.0128 @ 0.51% | 0.0 |
| diphoton_double_isr_background_o | 337113 | 0.0126 @ 0.5176 | 0.0 |
| $\mathrm{merged_lhe/-}$ | | | |
| diphoton_double_isr_background_l | | | |

$2.17 \quad \ \, \text{bg_dip_1600_inf}$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 1040000 events.

• Normalization to the luminosity: 187+/- 1 events.

 \bullet Ratio (event weight): 0.00018 .

| Path to the event file | Nr. of events | Cross section (pb) | Negative wgts (%) |
|----------------------------------|---------------|--------------------|-------------------|
| /Users/elijahsheridan/- | | | |
| $MG5_aMC_v2_6_5/-$ | | | |
| $axion_pheno/madgraph_data/-$ | 1040000 | 0.00469 @ 0.15% | 0.0 |
| diphoton_double_isr_background_c | 1040000 | 0.00409 @ 0.15% | 0.0 |
| merged_lhe/- | | | |
| diphoton_double_isr_background_h | | | |

3 Histos and cuts

3.1 Cut 1

* Cut: select (sdETA (jets[1] jets[2]) > 3.6 or sdETA (jets[1] jets[2]) < -3.6) and M (jets[1] jets[2]) > 1250.0

| Dataset | Events kept: K | Rejected events: R | Efficiency: K / (K + R) | Cumul. efficiency: K / Initial |
|--------------|---------------------|-----------------------|-------------------------|--|
| signal | 814.6 +/- 25.5 | 3279.5 + /- 25.6 | 0.19896 + / - 0.00624 | 0.19896 + / - 0.00624 |
| bg_vbf_0_10 | 204.2 +/- 14.2 | 11946.1 +/- 26.8 | 0.01681 + / - 0.00117 | 0.01681 + / - 0.00117 |
| bg_vbf_100_ | 950.9 +/- 29.3 | 8744.4 +/- 32.9 | 0.09808 + / - 0.00302 | 0.09808 + / - 0.00302 |
| bg_vbf_200_ | 1147.9 +/- 30.2 | 4265.4 + /- 31.3 | 0.21205 + / - 0.00556 | 0.21205 + / - 0.00556 |
| bg_vbf_400_ | 273.7 +/- 14.1 | 713.1 +/- 14.1 | 0.2774 + / - 0.0143 | 0.2774 + / - 0.0143 |
| bg_vbf_600_ | 47.78 +/- 6.22 | 204.30 +/- 6.23 | 0.1895 + / - 0.0247 | 0.1895 + / - 0.0247 |
| bg_vbf_800_ | 12.06 + / - 3.29 | 102.70 + / - 3.29 | 0.1051 + / - 0.0286 | 0.1051 + / - 0.0286 |
| bg_vbf_1200 | 0.678 +/- 0.810 | 19.92 +/- 0.81 | 0.0329 + / - 0.0393 | 0.0329 + / - 0.0393 |
| bg_vbf_1600 | 0.0483 + / - 0.2191 | 7.610 +/- 0.219 | 0.00631 +/- 0.02860 | 0.00631 + / - 0.02860 |
| bg_dip_0_10 | 229.4 +/- 15.1 | 2710617 +/- 4612 | 8.46e-05 +/- 5.59e-06 | $8.46\text{e-}05 +/- 5.59\text{e-} \ 06$ |
| bg_dip_100_ | 990.1 +/- 31.5 | 1094372 +/- 1526 | 9.04e-04 +/- 2.87e-05 | 9.04e-04 +/- 2.87e- 05 |
| 1 1: 200 | 10410 / 40 5 | 007007 / 410 | 0.006854 +/- | 0.006854 +/- |
| bg_dip_200_ | 1641.8 + / - 40.5 | 237907 + / - 412 | 0.000169 | 0.000169 |
| ha din 400 | F02 2 1 / 24 1 | 202011 / 161 | 0.020599 +/- | 0.020599 +/- |
| bg_dip_400_ | 593.2 +/- 24.1 | 28205.5 + /- 56.5 | 0.000837 | 0.000837 |
| bg_dip_600_ | 88.41 +/- 9.35 | 6585.9 +/- 28.8 | 0.0132 +/- 0.0014 | 0.0132 + / - 0.0014 |
| bg_dip_800_ | 22.00 +/- 4.67 | 2920.34 +/- 6.86 | 0.00748 + / - 0.00159 | 0.00748 + / - 0.00159 |
| bg_dip_1200_ | 1.34 +/- 1.16 | 512.16 +/- 2.87 | 0.00261 + / - 0.00225 | 0.00261 + / - 0.00225 |
| bg_dip_1600 | 0.0921 + / - 0.3034 | 187.691 + / - 0.412 | 0.00049 + / - 0.00162 | 0.00049 + / - 0.00162 |

3.2 Cut 2 $* \mbox{ Cut: select PT (a[1])} > 500.0 \mbox{ and M (a[1] a[2])} > 400.0$

| Dataset | Events kept: K | Rejected events: | Efficiency: K / (K + R) | Cumul. efficiency: K / Initial |
|--------------|----------------------|-------------------|--------------------------|---|
| signal | 400.8 +/- 19.0 | 413.7 +/- 19.3 | 0.4921 +/- 0.0175 | 0.09790 +/- 0.00464 |
| bg_vbf_0_10 | 0.0 +/- 0.0 | 204.2 +/- 14.2 | 0.0 +/- 0.0 | 0.0 +/- 0.0 |
| bg_vbf_100_ | 0.0503 + / - 0.2242 | 950.8 +/- 29.3 | 5.29e-05 +/- 2.36e-04 | 5.18e-06 +/- 2.31e- 05 |
| bg_vbf_200_ | 0.33 +/- 0.57 | 1147.5 +/- 30.2 | 0.000287 +/- 0.000500 | 6.10e-05 +/- 1.06e- 04 |
| bg_vbf_400_ | 1.11 + /- 1.05 | 272.6 +/- 14.1 | 0.00405 + / - 0.00384 | 0.00112 +/- 0.00107 |
| bg_vbf_600_ | 1.10 +/- 1.04 | 46.68 +/- 6.17 | 0.0229 +/- 0.0217 | 0.00435 + / - 0.00414 |
| bg_vbf_800_ | 0.628 + / - 0.790 | 11.43 +/- 3.21 | 0.052 + / - 0.064 | 0.00547 + / - 0.00688 |
| bg_vbf_1200 | 0.0589 + / - 0.2424 | 0.619 + / - 0.775 | 0.087 + / - 0.342 | 0.00286 + / - 0.01177 |
| bg_vbf_1600_ | 0.00667 + /- 0.08166 | 0.0416 +/- 0.2034 | 0.138 +/- 1.571 | 0.000872 +/- 0.010663 |
| bg_dip_0_10 | 0.0 +/- 0.0 | 229.4 +/- 15.1 | 0.0 +/- 0.0 | 0.0 +/- 0.0 |
| bg_dip_100_ | 0.0 +/- 0.0 | 990.1 +/- 31.5 | 0.0 +/- 0.0 | 0.0 +/- 0.0 |
| bg_dip_200_ | 0.691 + / - 0.831 | 1641.1 +/- 40.5 | 0.000421 +/- 0.000506 | $oxed{2.88\text{e-}06} +/\text{-} 3.47\text{e-} \ 06$ |
| bg_dip_400_ | 3.18 +/- 1.78 | 590.0 +/- 24.1 | 0.005 +/- 0.003 | 1.11e-04 +/- 6.20e- 05 |
| bg_dip_600_ | 2.43 +/- 1.56 | 85.98 +/- 9.22 | 0.0275 +/- 0.0174 | 0.000364 +/- 0.000233 |
| bg_dip_800_ | 1.28 +/- 1.13 | 20.71 +/- 4.54 | 0.06 +/- 0.05 | $0.000437 + /- \\ 0.000385$ |
| bg_dip_1200_ | 0.107 +/- 0.326 | 1.23 +/- 1.11 | 0.0794 +/- 0.2335 | 0.000208 +/- 0.000636 |
| bg_dip_1600_ | 0.011 +/- 0.105 | 0.0811 +/- 0.2846 | 0.12 +/- 1.07 | 5.87e-05 +/- 5.59e- 04 |

4 Summary

4.1 Cut-flow charts

- \bullet How to compare signal (S) and background (B): S/sqrt(S+B) .
- \bullet Object definition selections are indicated in cyan.
- Reject and select are indicated by 'REJ' and 'SEL' respectively

| Cuts | Signal (S) | Background (B) | S vs B |
|--------------------------|---------------------|--------------------|-----------------------|
| Initial (no cut) | 4094.08 + /- 1.13 | 4113516 + / - 4877 | 2.01760 + / - 0.00132 |
| SEL: (sdETA (jets[1] | | | |
| jets[2]) > 3.6 or sdETA | 814.6 + /- 25.5 | 6203.5 +/- 76.0 | 9.723 + / - 0.292 |
| (| | | |
| SEL: PT (a[1]) > | | | |
| 500.0 and M (a[1] a[2] | 400.8 + /- 19.0 | 10.98 + /- 3.31 | 19.752 + / - 0.488 |
|) > 400 | | | |