

Generated by elijahsheridan on 27 March 2020, 14:13:28

This report has been generated automatically by Madanalysis 5.

Please cite:

E. Conte, B. Fuks and G. Serret,

MadAnalysis 5, A User-Friendly Framework for Collider Phenomenology, Comput. Phys. Commun. **184** (2013) 222-256, arXiv:1206.1599 [hep-ph].

To contact us:

 ${\bf http://madanalysis.irmp.ucl.ac.be} \\ {\bf ma5team@iphc.cnrs.fr} \\$

Contents Setup 2 2 1.1 Command history 1.2 Configuration 3 Datasets 4 2.1signal 4 2.2 $bg_vbf_0_100$ 4 2.3 $bg_vbf_100_200$ 4 2.4 $bg_vbf_200_400$ 5 $bg_vbf_400_600$ 2.55 $2.6 \quad \, \mathrm{bg_vbf_600_800}$ 6 $2.7 ext{ bg_vbf_}800_1200$ 6 bg_vbf_1200_1600 7 2.8 2.9 bg_vbf_1600_inf 7 $2.10 \ bg_dip_0_100$ 7 $2.11 \ \ \mathrm{bg_dip_100_200}$ 8 2.12 bg dip 200 4008 2.13 bg dip 400 600 9 9 $2.14 \ \ bg_dip_600_800$ $2.15 \ \ bg_dip_800_1200$ 9 $2.16 \ \ bg_dip_1200_1600$ 10 $2.17 \hspace{0.1in} bg_dip_1600_inf$ 10 Histos and cuts 11 3.1 Cut 1 11 3.2 Cut 2 12 Summary **13**

13

4.1 Cut-flow charts

1 Setup

1.1 Command history

```
ma5># set directory where running "./bin/ma5"; set lumi; define the signal significance
ma5>set main.currentdir = /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno # need to
change this directory path -> exit and type "pwd" to get the path
ma5>set main.lumi = 40.0
ma5># import samples -> change the path to the LHE file
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/axion_signal/-
axion_signal_gurrola_cuts_1MeV.lhe.gz as signal
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/vbf_diphoton_background_
merged_lhe/vbf_diphoton_background_ht_0_100_merged.lhe.gz as bg_vbf_0_100
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/vbf_diphoton_background_
merged_lhe/vbf_diphoton_background_ht_100_200_merged.lhe.gz as bg_vbf_100_200
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/vbf_diphoton_background_
merged_lhe/vbf_diphoton_background_ht_200_400_merged.lhe.gz as bg_vbf_200_400
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/vbf_diphoton_background_
merged_lhe/vbf_diphoton_background_ht_400_600_merged.lhe.gz as bg_vbf_400_600
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/vbf_diphoton_background_
merged_lhe/vbf_diphoton_background_ht_600_800_merged.lhe.gz as bg_vbf_600_800
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/vbf_diphoton_background_
merged_lhe/vbf_diphoton_background_ht_800_1200_merged.lhe.gz as bg_vbf_800_1200
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/vbf_diphoton_background_
merged_lhe/vbf_diphoton_background_ht_1200_1600_merged.lhe.gz as bg_vbf_1200_1600
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/vbf_diphoton_background_
merged_lhe/vbf_diphoton_background_ht_1600_inf_merged.lhe.gz as bg_vbf_1600_inf
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/diphoton_double_isr_back
merged_lhe/diphoton_double_isr_background_ht_0_100_merged.lhe.gz as bg_dip_0_100
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/diphoton_double_isr_back
merged_lhe/diphoton_double_isr_background_ht_100_200_merged.lhe.gz as bg_dip_100_200
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/diphoton_double_isr_back
merged_lhe/diphoton_double_isr_background_ht_200_400_merged.lhe.gz as bg_dip_200_400
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/diphoton_double_isr_back
merged_lhe/diphoton_double_isr_background_ht_400_600_merged.lhe.gz as bg_dip_400_600
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/diphoton_double_isr_back
merged_lhe/diphoton_double_isr_background_ht_600_800_merged.lhe.gz as bg_dip_600_800
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/diphoton_double_isr_back
merged_lhe/diphoton_double_isr_background_ht_800_1200_merged.lhe.gz as bg_dip_800_1200
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/diphoton_double_isr_back
merged_lhe/diphoton_double_isr_background_ht_1200_1600_merged.lhe.gz as bg_dip_1200_1600
ma5>import /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/madgraph_data/diphoton_double_isr_back
merged_lhe/diphoton_double_isr_background_ht_1600_inf_merged.lhe.gz as bg_dip_1600_inf
ma5># define bg and signal samples
ma5>set signal.type = signal
ma5>set bg_vbf_0_100.type = background
ma5>set bg_vbf_100_200.type = background
ma5>set bg_vbf_200_400.type = background
ma5>set bg_vbf_400_600.type = background
ma5>set bg_vbf_600_800.type = background
```

```
ma5>set bg_vbf_800_1200.type = background
ma5>set bg_vbf_1200_1600.type = background
ma5>set bg_vbf_1600_inf.type = background
ma5>set bg_dip_0_100.type = background
ma5>set bg_dip_100_200.type = background
ma5>set bg_dip_200_400.type = background
ma5>set bg_dip_400_600.type = background
ma5>set bg_dip_600_800.type = background
ma5>set bg_dip_800_1200.type = background
ma5>set bg_dip_1200_1600.type = background
ma5>set bg_dip_1600_inf.type = background
ma5># define weights for the samples
ma5>#set sample_1.weight = 1
ma5>#set sample_2.weight = 1
ma5># a jet can be from a light quark or b quark
ma5>define jets = j
ma5>define e = e+ e-
ma5>define mu = mu+ mu-
ma5>define ta = ta+ ta-
ma5>define lept = e mu ta
ma5>define ax = 9000005
ma5># selections
ma5>select (sdETA(jets[1] jets[2]) > 2.6 or sdETA(jets[1] jets[2]) < -2.6) and M(jets[1] jets[2]) < -2.6) and M(jets[1] jets[2]) < -2.6)
jets[2]) > 1250
ma5>select PT(a[1]) > 400 and M(a[1] a[2]) > 650
ma5>submit analysis_loose_pta400_maa650
```

1.2 Configuration

- MadAnalysis version 1.6.33 (2017/11/20).
- Histograms given for an integrated luminosity of 40.0fb⁻¹.

2 Datasets

2.1 signal

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: signal events.

• Generated events: 1000000 events.

• Normalization to the luminosity: 4094+/- 2 events.

• Ratio (event weight): 0.0041.

Path to the event file	Nr. of events	Cross section (pb)	Negative wgts (%)
/Users/elijahsheridan/-			
$MG5_aMC_v2_6_5/-$			
axion_pheno/-	1000000	0.102 @ 0.028%	0.0
madgraph_data/axion_signal/-			
_axion_signal_gurrola_cuts_1MeV.ll			

$2.2 \quad bg_vbf_0_100$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 1000000 events.

• Normalization to the luminosity: 12150+/- 24 events.

 \bullet Ratio (event weight): 0.012 $\,$.

Path to the event file	Nr. of events	Cross section (pb)	Negative wgts (%)
/Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/- vbf_diphoton_background_data/- merged_lhe/- vbf_diphoton_background_ht 0 10	1000000	0.304 @ 0.19%	0.0

$2.3 \quad \text{bg vbf } 100 \quad 200$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 965662 events.

 \bullet Normalization to the luminosity: 9695+/- 17 $\,$ events.

• Ratio (event weight): 0.01.

Path to the event file	Nr. of events	Cross section (pb)	Negative wgts (%)
/Users/elijahsheridan/-			
$MG5_aMC_v2_6_5/-$			
$axion_pheno/madgraph_data/-$	067660	0.242 @ 0.17%	0.0
vbf_diphoton_background_data/-	965662	0.242 @ 0.17%	0.0
merged_lhe/-			
vbf_diphoton_background_ht_100_			

$\mathbf{2.4} \quad \mathbf{bg_vbf_200_400}$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 984165 events.

 \bullet Normalization to the luminosity: 5413+/- 11 events.

• Ratio (event weight): 0.0055.

Path to the event file	Nr. of events	Cross section (pb)	Negative wgts (%)
/Users/elijahsheridan/-			
$MG5_aMC_v2_6_5/-$			
$axion_pheno/madgraph_data/-$	004165	0.127 @ 0.207	0.0
vbf_diphoton_background_data/-	984165	0.135 @ 0.2%	0.0
$\mathrm{merged_lhe/-}$			
vbf_diphoton_background_ht_200_			

$\mathbf{2.5} \quad \mathbf{bg_vbf_400_600}$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 1000000 events.

• Normalization to the luminosity: 986+/-2 events.

 \bullet Ratio (event weight): 0.00099 .

Path to the event file	Nr. of events	Cross section (pb)	Negative wgts (%)
/Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/- vbf_diphoton_background_data/- merged_lhe/- vbf_diphoton_background_ht_400_	1000000	0.0247 @ 0.14%	0.0

$2.6 \quad \mathrm{bg_vbf_600_800}$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 1000000 events.

• Normalization to the luminosity: 252+/- 1 events.

• Ratio (event weight): 0.00025.

Path to the event file	Nr. of events	Cross section (pb)	Negative wgts (%)
/Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/- vbf_diphoton_background_data/- merged_lhe/- vbf_diphoton_background_ht 600	1000000	0.0063 @ 0.13%	0.0

$2.7 \quad \mathrm{bg_vbf_800_1200}$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

 \bullet Generated events: 400839 events.

• Normalization to the luminosity: 114+/- 1 events.

 \bullet Ratio (event weight): 0.00028.

Path to the event file	Nr. of events	Cross section (pb)	Negative wgts (%)
/Users/elijahsheridan/-			
$MG5_aMC_v2_6_5/-$			
$axion_pheno/madgraph_data/-$	400020	0.00287 @ 0.16%	0.0
vbf_diphoton_background_data/-	400839	0.00207 @ 0.10%	0.0
merged_lhe/-			
vbf_diphoton_background_ht_800_			

$2.8 \quad \ \, bg_vbf_1200_1600$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 953803 events.

• Normalization to the luminosity: 20+/- 1 events.

• Ratio (event weight): 2.1e-05 .

Path to the event file	Nr. of events	Cross section (pb)	Negative wgts (%)
/Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/- vbf_diphoton_background_data/- merged_lhe/- vbf_diphoton_background_ht_1200	953803	0.000515 @ 0.16%	0.0

2.9 bg vbf 1600 inf

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

 \bullet Generated events: 270148 $\,$ events.

• Normalization to the luminosity: 7+/-1 events.

• Ratio (event weight): 2.6e-05 .

Path to the event file	Nr. of events	Cross section (pb)	Negative wgts (%)
/Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/- vbf_diphoton_background_data/- merged_lhe/- vbf_diphoton_background_ht 1600	270148	0.000191 @ 0.11%	0.0

$2.10 \quad \text{bg dip } 0 \quad 100$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 1040000 events.

 \bullet Normalization to the luminosity: 2710847+/- 4614 events.

 \bullet Ratio (event weight): 2.6 - warning: please generate more events (weight larger than 1)!

Path to the event file	Nr. of events	Cross section (pb)	Negative wgts (%)
/Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/-	1040000	67.8 @ 0.17%	0.0
diphoton_double_isr_background_d merged_lhe/- diphoton_double_isr_background_h	1040000	01.0 & 0.11/0	0.0

2.11 bg dip 100 200

- \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .
- Sample consisting of: background events.
- Generated events: 1040000 events.
- Normalization to the luminosity: 1095362+/- 1528 events.
- Ratio (event weight): 1.1 warning: please generate more events (weight larger than 1)!

Path to the event file	Nr. of events	Cross section (pb)	Negative wgts (%)
/Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/- diphoton_double_isr_background_d merged_lhe/- diphoton_double_isr_background_l	1040000	27.4 @ 0.14%	0.0

$2.12 \quad \ \, \text{bg_dip_200_400}$

- \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .
- Sample consisting of: background events.
- Generated events: 1040000 events.
- Normalization to the luminosity: 239548+/- 414 events.
- Ratio (event weight): 0.23 .

Path to the event file	Nr. of events	Cross section (pb)	Negative wgts (%)
/Users/elijahsheridan/-			
$MG5_aMC_v2_6_5/-$			
$axion_pheno/madgraph_data/-$	1040000	5.99 @ 0.17%	0.0
diphoton_double_isr_background_d	1040000	5.99 @ 0.1770	0.0
merged_lhe/-			
diphoton_double_isr_background_l			

$2.13 \quad bg_dip_400_600$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 1040000 events.

• Normalization to the luminosity: 28798+/- 53 events.

• Ratio (event weight): 0.028 .

Path to the event file	Nr. of events	Cross section (pb)	Negative wgts (%)
/Users/elijahsheridan/-			
$MG5_aMC_v2_6_5/-$			
$axion_pheno/madgraph_data/-$	1040000	0.72 @ 0.18%	0.0
diphoton_double_isr_background_o	1040000	0.72 @ 0.1670	0.0
$merged_lhe/-$			
diphoton_double_isr_background_l			

$2.14 ext{ bg_dip_}600_800$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

 \bullet Generated events: 662009 events.

• Normalization to the luminosity: 6674+/- 28 events.

• Ratio (event weight): 0.01 .

Path to the event file	Nr. of events	Cross section (pb)	Negative wgts (%)
/Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/-	662009	0.167 @ 0.41%	0.0
diphoton_double_isr_background_d merged_lhe/- diphoton_double_isr_background_h	002000	0.107 @ 0.4170	0.0

2.15 bg dip 800 1200

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 1040000 events.

 \bullet Normalization to the luminosity: 2942+/- 6 events.

 \bullet Ratio (event weight): 0.0028 % =0.0028 .

Path to the event file	Nr. of events	Cross section (pb)	Negative wgts (%)
/Users/elijahsheridan/- MG5_aMC_v2_6_5/- axion_pheno/madgraph_data/- diphoton_double_isr_background_d merged_lhe/- diphoton_double_isr_background_h	1040000	0.0736 @ 0.17%	0.0

2.16 bg dip 1200 1600

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 337115 events.

• Normalization to the luminosity: 513+/-3 events.

Path to the event file	Nr. of events	Cross section (pb)	Negative wgts (%)
/Users/elijahsheridan/-			
$MG5_aMC_v2_6_5/-$			
$axion_pheno/madgraph_data/-$	337115	0.0128 @ 0.51%	0.0
diphoton_double_isr_background_o	337113	0.0126 @ 0.5176	0.0
$\mathrm{merged_lhe/-}$			
diphoton_double_isr_background_l			

$2.17 \quad \ \, \text{bg_dip_1600_inf}$

 \bullet Samples stored in the directory: /Users/elijahsheridan/MG5_aMC_v2_6_5/axion_pheno/optimization .

• Sample consisting of: background events.

• Generated events: 1040000 events.

• Normalization to the luminosity: 187+/- 1 events.

 \bullet Ratio (event weight): 0.00018 .

Path to the event file	Nr. of events	Cross section (pb)	Negative wgts (%)
/Users/elijahsheridan/-			
$MG5_aMC_v2_6_5/-$			
$axion_pheno/madgraph_data/-$	1040000	0.00469 @ 0.15%	0.0
diphoton_double_isr_background_c	1040000	0.00409 @ 0.15%	0.0
merged_lhe/-			
diphoton_double_isr_background_h			

3 Histos and cuts

3.1 Cut 1

* Cut: select (sdETA (jets[1] jets[2]) > 2.6 or sdETA (jets[1] jets[2]) < -2.6) and M (jets[1] jets[2]) > 1250.0

Dataset	Events kept: K	Rejected events:	Efficiency: $K / (K + R)$	Cumul. efficiency: K / Initial
signal	1711.8 +/- 31.6	2382.3 +/- 31.6	0.41812 + / - 0.00771	0.41812 + / - 0.00771
bg_vbf_0_10	204.2 + / - 14.2	11946.1 + /- 26.8	0.01681 + / - 0.00117	0.01681 + / - 0.00117
bg_vbf_100_	950.9 + /- 29.3	8744.4 +/- 32.9	0.09808 + / - 0.00302	0.09808 + / - 0.00302
bg_vbf_200_	1147.9 +/- 30.2	4265.4 +/- 31.3	0.21205 + / - 0.00556	0.21205 + / - 0.00556
bg_vbf_400_	349.4 +/- 15.0	637.5 +/- 15.0	0.3540 + / - 0.0152	0.3540 + / - 0.0152
bg_vbf_600_	111.21 +/- 7.88	140.87 +/- 7.89	0.4412 + / - 0.0313	0.4412 + / - 0.0313
bg_vbf_800_	40.31 + /- 5.11	74.45 + /- 5.12	0.3513 + / - 0.0446	0.3513 + / - 0.0446
bg_vbf_1200	4.49 + /- 1.87	16.10 +/- 1.87	0.218 +/- 0.091	0.218 +/- 0.091
bg_vbf_1600	0.784 + / - 0.839	6.874 +/- 0.839	0.102 +/- 0.110	0.102 +/- 0.110
bg_dip_0_10	229.4 +/- 15.1	2710617 +/- 4612	8.46e-05 +/- 5.59e-06	8.46e-05 +/- 5.59e-
bg_dip_100_	990.1 +/- 31.5	1094372 +/- 1526	9.04e-04 +/- 2.87e-05	9.04e-04 +/- 2.87e- 05
bg_dip_200_	1641.8 +/- 40.5	237907 +/- 412	0.006854 +/-	0.006854 +/-
	,	,	0.000169	0.000169
bg_dip_400_	1066.5 +/- 32.1	27732.1 +/- 59.6	0.03703 +/- 0.00111	0.03703 +/- 0.00111
bg_dip_600_	531.3 +/- 22.2	6143.1 +/- 33.7	0.07960 +/- 0.00331	0.07960 +/- 0.00331
bg_dip_800_	193.3 +/- 13.4	2749.1 +/- 14.2	0.06568 + / - 0.00457	0.06568 + / - 0.00457
bg_dip_1200_	21.80 + / - 4.57	491.71 +/- 5.22	0.0424 +/- 0.0089	0.0424 + / - 0.0089
bg_dip_1600_	4.1 + /- 2.0	183.71 + / - 2.01	0.0217 + / - 0.0106	0.0217 + / - 0.0106

3.2 Cut 2 $* \mbox{ Cut: select PT (a[1])} > 400.0 \mbox{ and M (a[1] a[2])} > 650.0$

Dataset	Events kept: K	Rejected events:	Efficiency: $K / (K + R)$	Cumul. efficiency: K / Initial
signal	951.9 + /- 27.0	759.9 +/- 24.9	0.556 + / - 0.012	0.2325 + / - 0.0066
bg_vbf_0_10	0.0 +/- 0.0	204.2 +/- 14.2	0.0 +/- 0.0	0.0 +/- 0.0
bg_vbf_100_	0.151 +/- 0.388	950.7 +/- 29.3	0.000158 +/- 0.000408	1.55e-05 +/- 4.00e- 05
bg_vbf_200_	0.985 +/- 0.992	1146.9 +/- 30.2	0.000858 +/- 0.000864	0.000182 +/- 0.000183
bg_vbf_400_	1.40 +/- 1.18	347.9 +/- 15.0	0.00402 + / - 0.00339	0.0014 +/- 0.0012
bg_vbf_600_	1.03 + / - 1.02	110.17 +/- 7.88	0.0093 + / - 0.0091	0.00411 + / - 0.00403
bg_vbf_800_	0.715 + / - 0.843	39.60 +/- 5.09	0.0177 +/- 0.0208	0.00623 + / - 0.00735
bg_vbf_1200	0.12 + / - 0.34	4.37 + /- 1.86	0.0266 + / - 0.0759	0.0058 + / - 0.0167
bg_vbf_1600	0.0242 + / - 0.1553	0.76 + / - 0.83	$0.0308 + / ext{-} 0.1952$	0.00316 + / - 0.02027
bg_dip_0_10	0.0 +/- 0.0	229.4 +/- 15.1	0.0 +/- 0.0	0.0 +/- 0.0
bg_dip_100_	0.0 + / - 0.0	990.1 + / - 31.5	0.0 +/- 0.0	0.0 +/- 0.0
bg_dip_200_	1.84 +/- 1.36	1640.0 +/- 40.5	0.001122 +/- 0.000826	$egin{array}{cccccccccccccccccccccccccccccccccccc$
bg_dip_400_	2.66 +/- 1.63	1063.9 +/- 32.1	0.00249 +/- 0.00153	9.23e-05 +/- 5.66e- 05
bg_dip_600_	2.6 + /- 1.6	528.7 +/- 22.2	0.005 + / - 0.003	0.00038 + / - 0.00024
bg_dip_800_	2.15 +/- 1.47	191.1 +/- 13.4	0.01113 +/- 0.00754	0.000731 +/- 0.000498
bg_dip_1200	0.344 + / - 0.587	21.45 +/- 4.54	0.0158 + / - 0.0267	0.00067 +/- 0.00114
bg_dip_1600_	0.0608 + / - 0.2466	4.01 +/- 1.98	0.0149 +/- 0.0601	0.000324 +/- 0.001313

4 Summary

4.1 Cut-flow charts

- \bullet How to compare signal (S) and background (B): S/sqrt(S+B) .
- \bullet Object definition selections are indicated in cyan.
- Reject and select are indicated by 'REJ' and 'SEL' respectively

Cuts	Signal (S)	Background (B)	S vs B
Initial (no cut)	4094.08 + /- 1.13	4113516 + / - 4877	2.01760 + / - 0.00132
SEL: (sdETA (jets[1]			
$\mathrm{jets}[2]$) > 2.6 or sdETA	1711.8 + / - 31.6	7487.3 + / -82.9	17.848 + / - 0.309
(
SEL: PT (a[1]) >			
400.0 and M (a[1] a[2]	951.9 + /- 27.0	14.05 + /- 3.75	30.628 + / - 0.445
)>650			