

Problem Set 2

Seth Harrison

02/03/2020

```
# Load Packages/Data
```

```
library(tidyverse)
library(readr)
library(rsample)
library(broom)
library(rcfss)
library(ISLR)
library(yardstick)
```

```
nes <- read_csv("./nes2008.csv")
```

1. (10 points) Estimate the MSE of the model using the traditional approach. That is, fit the linear regression model using the *entire* dataset and calculate the mean squared error for the *entire* dataset. Present and discuss your results at a simple, high level.

```
# Fit Linear Model
```

```
Model1 <- lm(nes$biden~nes$female+nes$age+nes$educ+nes$dem+nes$rep)
summary(Model1)
```

```
##
## Call:
## lm(formula = nes$biden ~ nes$female + nes$age + nes$educ + nes$dem +
##     nes$rep)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -75.546 -11.295   1.018  12.776  53.977
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  58.81126    3.12444  18.823  < 2e-16 ***
## nes$female    4.10323    0.94823   4.327 1.59e-05 ***
## nes$age       0.04826    0.02825   1.708  0.0877 .
## nes$educ     -0.34533    0.19478  -1.773  0.0764 .
## nes$dem      15.42426    1.06803  14.442  < 2e-16 ***
## nes$rep     -15.84951    1.31136 -12.086  < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 19.91 on 1801 degrees of freedom
## Multiple R-squared:  0.2815, Adjusted R-squared:  0.2795
## F-statistic: 141.1 on 5 and 1801 DF, p-value: < 2.2e-16
```

```
# Calculate MSE
```

```
mse1 <- mean(Model1$residuals^2)
mse1
```

```
## [1] 395.2702
```

The mean squared error (MSE) first eliminates negative directionality by squaring the residuals, or the distance between the observed observation and observation expected by the regression line. It then takes arithmetic mean of those products. One easy way to interpret MSE is to convert it back to a mean error by taking the square root of the MSE. In this case, the mean error is about 20, meaning that, on average, the expected thermometer rating was about 20 units off (either higher or lower) from the observed rating.

2. (30 points) Calculate the test MSE of the model using the simple holdout validation approach.

- (5 points) Split the sample set into a training set (50%) and a holdout set (50%).

```
# Split nes into test and train
```

```
set.seed(1)
```

```
nes_split <- initial_split(data = nes,
                           prop = 0.5)
```

```
nes_train <- training(nes_split)
nes_test <- testing(nes_split)
```

- (5 points) Fit the linear regression model using only the training observations.

```
# Fit Linear Model Using nes_train
```

```
Model2 <- lm(biden~female+age+educ+dem+rep, data = nes_train)
mse2 <- mean(Model2$residuals^2)
summary(Model2)
```

```
##
```

```
## Call:
```

```
## lm(formula = biden ~ female + age + educ + dem + rep, data = nes_train)
```

```
##
```

```
## Residuals:
```

```
##      Min       1Q   Median       3Q      Max
## -75.875 -10.974   0.638  13.968  45.989
```

```
##
```

```
## Coefficients:
```

```
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  61.94663    4.52928  13.677 < 2e-16 ***
## female       5.14561    1.38493   3.715 0.000215 ***
## age        -0.02402    0.04197  -0.572 0.567281
## educ        -0.46983    0.28126  -1.670 0.095179 .
## dem         16.27265    1.55652  10.454 < 2e-16 ***
## rep        -16.41671    1.96592  -8.351 2.55e-16 ***
```

```
## ---
```

```
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
##
## Residual standard error: 20.74 on 898 degrees of freedom
## Multiple R-squared:  0.2799, Adjusted R-squared:  0.2759
## F-statistic: 69.8 on 5 and 898 DF,  p-value: < 2.2e-16

- (10 points) Calculate the MSE using only the test set observations.

test_mse <- augment(Model2, newdata = nes_test) %>%
  mse(truth = biden, estimate = .fitted)

test_mse$.estimate

## [1] 370.1792

- (10 points) How does this value compare to the training MSE from question 1?
Present numeric comparison and discuss a bit.
```

The first MSE was 395.27, while the second from the split data set was lower at 370.18. We avoided an optimistically biased, overfit evaluation by estimating the model from the training data and evaluating it on the test data but since the training and test sets were generated through random processes, we can expect high variance between MSE predictions dependent on the composition of the split.

3. (30 points) Repeat the simple validation set approach from the previous question 1000 times, using 1000 different splits of the observations into a training set and a test/validation set. Visualize your results as a sampling distribution (hint: think histogram or density plots). Comment on the results obtained.