## Exercises

- 1. Find **ALL** integer solutions to the following equation if the solution exists or show the solution does not exist.
  - (1) 4x + 6y = 2
  - (2) 4x + 6y = 6
  - (3) 15x + 51y = 41
  - (4) 121x 88y = 572

For Questions (1) and (2), plot the solution points (if exist) on the graph.

The following are constrained linear Diophantine equations.

- 2. How many ways are there to make \$2.00 dollars from only nickels and quarters? (Hints: converts this to a linear Diophantine equation. By its nature, solutions will be non-negative).
- 3. A grocer orders apples and bananas at a total cost of \$8.4. If the apples cost 25 cents each and the bananas 5 cents each, how many of each type of fruit did he order.
- 4. Find how many integer solutions there are to the following equation 3a + 5b = 7 subject to  $a, b \in [-10, 10]$ .
- 5. Find the least common multiples of:
  - (1) lcm(14, 15).
  - (2) lcm(240,610).
  - (3) lcm(n, n + 1).
  - (4) lcm(2n-1, 2n+1).
  - (5)  $lcm(2^5 \cdot 3 \cdot 5^6 \cdot 7^2 \cdot 11, 2^3 \cdot 5^8 \cdot 7^2 \cdot 13).$

(Hints: use the F.T.A and/or the relation between gcd and lcm).

- 6. Show that if a and b are nonzero positive integers then the  $gcd(a,b) \mid lcm(a,b)$ . But  $(gcd(a,b))^2 \nmid lcm(a,b)$  unless a, b are coprime.
- 7. When are the least common multiple and the greatest common divisor equal to each other?