

STATISTICS WORKSHEET-1

Q1 to Q9 have only one correct answer. Choose the correct option to answer your question.

1. Bernoulli random variables take (only) the values 1 and 0.
 - a) **True**
 - b) False
2. Which of the following theorem states that the distribution of averages of iid variables, properly normalized, becomes that of a standard normal as the sample size increases?
 - a) **Central Limit Theorem**
 - b) Central Mean Theorem
 - c) Centroid Limit Theorem
 - d) All of the mentioned
3. Which of the following is incorrect with respect to use of Poisson distribution?
 - a) Modeling event/time data
 - b) **Modeling bounded count data**
 - c) Modeling contingency tables
 - d) All of the mentioned
4. Point out the correct statement.
 - a) The exponent of a normally distributed random variables follows what is called the log- normal distribution
 - b) Sums of normally distributed random variables are again normally distributed even if the variables are dependent
 - c) The square of a standard normal random variable follows what is called chi-squared distribution
 - d) **All of the mentioned**
5. _____ random variables are used to model rates.
 - a) Empirical
 - b) Binomial
 - c) **Poisson**
 - d) All of the mentioned
6. 10. Usually replacing the standard error by its estimated value does change the CLT.
 - a) True
 - b) **False**
7. 1. Which of the following testing is concerned with making decisions using data?
 - a) Probability
 - b) **Hypothesis**
 - c) Causal
 - d) None of the mentioned
8. 4. Normalized data are centered at _____ and have units equal to standard deviations of the original data.
 - a) **0**
 - b) 5
 - c) 1
 - d) 10
9. Which of the following statement is incorrect with respect to outliers?
 - a) Outliers can have varying degrees of influence
 - b) Outliers can be the result of spurious or real processes
 - c) **Outliers cannot conform to the regression relationship**
 - d) None of the mentioned

Q10 and Q15 are subjective answer type questions, Answer them in your own words briefly.

10. What do you understand by the term Normal Distribution?

Normal distribution, also known as the Gaussian distribution, is a probability distribution that is symmetric about the mean, showing that data near the mean are more frequent in occurrence than data far from the mean.

11. How do you handle missing data? What imputation techniques do you recommend?

Use deletion methods to eliminate missing data. The deletion methods only work for certain datasets where participants have missing fields. ...

Use regression analysis to systematically eliminate data. ...

Data scientists can use data imputation techniques.

12. What is A/B testing?

A/B testing, also known as split testing, refers to a randomized experimentation process wherein two or more versions of a variable (web page, page element, etc.) are shown to different segments of website visitors at the same time to determine which version leaves the maximum impact and drive business metrics.



FLIP ROBO

13. Is mean imputation of missing data acceptable practice?

True, imputing the mean preserves the mean of the observed data. So if the data are missing completely at random, the estimate of the mean remains unbiased. ... Since most research studies are interested in the relationship among variables, mean imputation is not a good solution.

14. What is linear regression in statistics?

In statistics, linear regression is a linear approach for modelling the relationship between a scalar response and one or more explanatory variables (also known as dependent and independent variables). Linear regression has many practical uses.

What are the various branches of statistics?

There are three real branches of statistics: data collection, descriptive statistics and inferential statistics.