



**ATLAS PUB Note**  
ATL-PHYS-PUB-2019-034  
16th September 2019



## **HBSM working group hMSSM summary plots**

The ATLAS Collaboration

This note presents an update to the hMSSM summary plots. These plots summarize the interpretation of various HBSM analyses and the Higgs coupling combination in the hMSSM model. This version is an update of the plots released in October 2018.






Figure 1: Regions of the  $[m_A, \tan \beta]$  plane excluded in the hMSSM model via direct searches for heavy Higgs bosons and fits to the measured rates of observed Higgs boson production and decays. Limits are quoted at 95% CL and are indicated for the data (solid lines) and the expectation for the SM Higgs sector (dashed lines). The light shaded or hashed regions indicate the observed exclusions. The cross sections for the Higgs boson production in the hMSSM [1, 2] are calculated using up to NNLO QCD corrections for gluon-gluon fusion and  $b$ -associated production in the five-flavour scheme as implemented in Sush1 [3–6]. For  $b$ -associated production a cross section in the four-flavour scheme is calculated as described in Refs. [7, 8] and the results are combined with the five-flavour scheme calculation following Ref. [9]. The Higgs boson widths and branching ratios have been calculated using HDECAY [10]. The procedure for the calculation of the cross sections and the branching ratios follows Ref. [11]. Discussion of the application of the hMSSM can be found in Ref. [12].

## 1 Plots and results used in them

Figures 1–4 shows the expected and observed limits at 95% CL in the hMSSM of the following analyses:

- Search for additional heavy neutral Higgs and gauge bosons in the ditau final state produced in  $36 \text{ fb}^{-1}$  of  $pp$  collisions at  $\sqrt{s} = 13 \text{ TeV}$  with the ATLAS detector. [13]
- Search for charged Higgs bosons decaying via  $H^\pm \rightarrow \tau^\pm \nu_\tau$  in the  $\tau$ +jets and  $\tau$ +lepton final states with  $36 \text{ fb}^{-1}$  of  $pp$  collision data recorded at  $\sqrt{s} = 13 \text{ TeV}$  with the ATLAS experiment. [14]
- Search for charged Higgs bosons decaying into top and bottom quarks at  $\sqrt{s} = 13 \text{ TeV}$  with the ATLAS detector. [15]
- Search for heavy neutral Higgs bosons produced in association with  $b$ -quarks and decaying to  $b$ -quarks at  $\sqrt{s} = 13 \text{ TeV}$  with the ATLAS detector. [16]




Figure 2: Regions of the  $[m_A, \tan \beta]$  plane excluded in the hMSSM model via direct searches for heavy Higgs bosons and fits to the measured rates of observed Higgs boson production and decays. Limits are quoted at 95% CL and are indicated for the data (solid lines) and the expectation for the SM Higgs sector (dashed lines). The light shaded or hashed regions indicate the observed exclusions. The cross sections for the Higgs boson production in the hMSSM [1, 2] are calculated using up to NNLO QCD corrections for gluon-gluon fusion and  $b$ -associated production in the five-flavour scheme as implemented in Sush1 [3–6]. For  $b$ -associated production a cross section in the four-flavour scheme is calculated as described in Refs. [7, 8] and the results are combined with the five-flavour scheme calculation following Ref. [9]. The Higgs boson widths and branching ratios have been calculated using HDECAY [10]. The procedure for the calculation of the cross sections and the branching ratios follows Ref. [11]. Discussion of the application of the hMSSM can be found in Ref. [12].

- Search for heavy  $ZZ$  resonances in the  $\ell^+\ell^-\ell^+\ell^-$  and  $\ell^+\ell^-\nu\bar{\nu}$  final states using proton–proton collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector. [17]
- Search for heavy resonances decaying into a  $W$  or  $Z$  boson and a Higgs boson in final states with leptons and  $b$ -jets in 36 fb $^{-1}$  of  $\sqrt{s} = 13$  TeV  $pp$  collisions with the ATLAS detector [18]
- Search for heavy resonances decaying into  $WW$  in the  $e\nu\mu\nu$  final state in  $pp$  collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector. [19]
- Combination of searches for Higgs boson pairs in  $pp$  collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector. [20]
- Combined measurements of Higgs boson production and decay using up to 80 fb $^{-1}$  of proton–proton collision data at  $\sqrt{s} = 13$  TeV collected with the ATLAS experiment’ [21]




Figure 3: Regions of the  $[m_A, \tan \beta]$  plane excluded in the hMSSM model via direct searches for heavy Higgs bosons and fits to the measured rates of observed Higgs boson production and decays. Limits are quoted at 95% CL and are indicated for the data (solid lines and shaded or hashed regions). The cross sections for the Higgs boson production in the hMSSM [1, 2] are calculated using up to NNLO QCD corrections for gluon-gluon fusion and  $b$ -associated production in the five-flavour scheme as implemented in *Sushi* [3–6]. For  $b$ -associated production a cross section in the four-flavour scheme is calculated as described in Refs. [7, 8] and the results are combined with the five-flavour scheme calculation following Ref. [9]. The Higgs boson widths and branching ratios have been calculated using *HDECAY* [10]. The procedure for the calculation of the cross sections and the branching ratios follows Ref. [11]. Discussion of the application of the hMSSM can be found in Ref. [12].




Figure 4: Regions of the  $[m_A, \tan \beta]$  plane excluded in the hMSSM model via direct searches for heavy Higgs bosons and fits to the measured rates of observed Higgs boson production and decays. Expected limits are quoted at 95% CL and are indicated for the expectation for the SM Higgs sector (dashed lines and shaded regions). The cross sections for the Higgs boson production in the hMSSM [1, 2] are calculated using up to NNLO QCD corrections for gluon-gluon fusion and  $b$ -associated production in the five-flavour scheme as implemented in *Sushi* [3–6]. For  $b$ -associated production a cross section in the four-flavour scheme is calculated as described in Refs. [7, 8] and the results are combined with the five-flavour scheme calculation following Ref. [9]. The Higgs boson widths and branching ratios have been calculated using *HDECAY* [10]. The procedure for the calculation of the cross sections and the branching ratios follows Ref. [11]. Discussion of the application of the hMSSM can be found in Ref. [12].

## References

- [1] A. Djouadi et al., *The post-Higgs MSSM scenario: Habemus MSSM?*, *Eur. Phys. J. C* **73** (2013) 2650, arXiv: [1307.5205 \[hep-ph\]](#) (cit. on pp. 2–5).
- [2] A. Djouadi, L. Maiani, A. Polosa, J. Quevillon and V. Riquer, *Fully covering the MSSM Higgs sector at the LHC*, *JHEP* **06** (2015) 168, arXiv: [1502.05653 \[hep-ph\]](#) (cit. on pp. 2–5).
- [3] R. V. Harlander, S. Liebler and H. Mantler, *SusHi: A program for the calculation of Higgs production in gluon fusion and bottom-quark annihilation in the Standard Model and the MSSM*, *Comput. Phys. Commun.* **184** (2013) 1605, arXiv: [1212.3249 \[hep-ph\]](#) (cit. on pp. 2–5).
- [4] R. Harlander and P. Kant, *Higgs production and decay: Analytic results at next-to-leading order QCD*, *JHEP* **12** (2005) 015, arXiv: [0509189 \[hep-ph\]](#) (cit. on pp. 2–5).
- [5] R. V. Harlander and W. B. Kilgore, *Higgs boson production in bottom quark fusion at next-to-next-to-leading order*, *Phys. Rev. D* **68** (2003) 013001, arXiv: [0304035 \[hep-ph\]](#) (cit. on pp. 2–5).
- [6] R. V. Harlander and W. B. Kilgore, *Next-to-next-to-leading order Higgs production at hadron colliders*, *Phys. Rev. Lett.* **88** (2002) 201801, arXiv: [0201206 \[hep-ph\]](#) (cit. on pp. 2–5).
- [7] S. Dawson, C. B. Jackson, L. Reina and D. Wackeroth, *Exclusive Higgs boson production with bottom quarks at hadron colliders*, *Phys. Rev. D* **69** (2004) 074027, arXiv: [0311067 \[hep-ph\]](#) (cit. on pp. 2–5).
- [8] S. Dittmaier, M. Krämer and M. Spira, *Higgs radiation off bottom quarks at the Tevatron and the CERN LHC*, *Phys. Rev. D* **70** (2004) 074010, arXiv: [0309204 \[hep-ph\]](#) (cit. on pp. 2–5).
- [9] R. Harlander, M. Kramer and M. Schumacher, *Bottom-quark associated Higgs-boson production: reconciling the four- and five-flavour scheme approach*, 2011, arXiv: [1112.3478 \[hep-ph\]](#) (cit. on pp. 2–5).
- [10] A. Djouadi, J. Kalinowski and M. Spira, *HDECAY: A Program for Higgs boson decays in the standard model and its supersymmetric extension*, *Comput. Phys. Commun.* **108** (1998) 56, arXiv: [hep-ph/9704448 \[hep-ph\]](#) (cit. on pp. 2–5).
- [11] D. de Florian et al., *Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector*, 2016, arXiv: [1610.07922 \[hep-ph\]](#) (cit. on pp. 2–5).
- [12] E. Bagnaschi et al., *Benchmark scenarios for low  $\tan \beta$  in the MSSM*, tech. rep. LHCHXSWG-2015-002, 2015, URL: <http://cds.cern.ch/record/2039911> (cit. on pp. 2–5).
- [13] ATLAS Collaboration, *Search for additional heavy neutral Higgs and gauge bosons in the ditau final state produced in  $36 \text{ fb}^{-1}$  of  $pp$  collisions at  $\sqrt{s} = 13 \text{ TeV}$  with the ATLAS detector*, *JHEP* **01** (2018) 055, arXiv: [1709.07242 \[hep-ex\]](#) (cit. on p. 2).
- [14] ATLAS Collaboration, *Search for charged Higgs bosons decaying via  $H^\pm \rightarrow \tau^\pm \nu_\tau$  in the  $\tau$ +jets and  $\tau$ +lepton final states with  $36 \text{ fb}^{-1}$  of  $pp$  collision data recorded at  $\sqrt{s} = 13 \text{ TeV}$  with the ATLAS experiment*, *JHEP* **09** (2018) 139, arXiv: [1807.07915 \[hep-ex\]](#) (cit. on p. 2).
- [15] ATLAS Collaboration, *Search for charged Higgs bosons decaying into top and bottom quarks at  $\sqrt{s} = 13 \text{ TeV}$  with the ATLAS detector*, *JHEP* **11** (2018) 085, arXiv: [1808.03599 \[hep-ex\]](#) (cit. on p. 2).
- [16] ATLAS Collaboration, *Search for heavy neutral Higgs bosons produced in association with  $b$ -quarks and decaying to  $b$ -quarks at  $\sqrt{s} = 13 \text{ TeV}$  with the ATLAS detector*, (2019), arXiv: [1907.02749 \[hep-ex\]](#) (cit. on p. 2).

- [17] ATLAS Collaboration, *Search for heavy ZZ resonances in the  $\ell^+\ell^-\ell^+\ell^-$  and  $\ell^+\ell^-\nu\bar{\nu}$  final states using proton–proton collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector*, *Eur. Phys. J. C* **78** (2018) 293, arXiv: [1712.06386 \[hep-ex\]](#) (cit. on p. 3).
- [18] ATLAS Collaboration, *Search for heavy resonances decaying into a W or Z boson and a Higgs boson in final states with leptons and b-jets in  $36\text{ fb}^{-1}$  of  $\sqrt{s} = 13$  TeV pp collisions with the ATLAS detector*, *JHEP* **03** (2018) 174, [Erratum: JHEP11,051(2018)], arXiv: [1712.06518 \[hep-ex\]](#) (cit. on p. 3).
- [19] ATLAS Collaboration, *Search for heavy resonances decaying into WW in the  $e\nu\mu\nu$  final state in pp collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector*, *Eur. Phys. J. C* **78** (2018) 24, arXiv: [1710.01123 \[hep-ex\]](#) (cit. on p. 3).
- [20] ATLAS Collaboration, *Combination of searches for Higgs boson pairs in pp collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector*, (2019), arXiv: [1906.02025 \[hep-ex\]](#) (cit. on p. 3).
- [21] ATLAS Collaboration, *Combined measurements of Higgs boson production and decay using up to  $80\text{ fb}^{-1}$  of proton–proton collision data at  $\sqrt{s} = 13$  TeV collected with the ATLAS experiment*, tech. rep. ATLAS-CONF-2019-005, CERN, 2019, URL: <https://cds.cern.ch/record/2668375> (cit. on p. 3).