
Shihab Muhtasim

STUDENT ID: 21301610

MAT 110

ASSIGNMENT 03

SET 7

Ans to the question no 01

Given function,

$$f(x) = \sin x + 2 \cos x$$

By differentiating this function we get,

$$f'(x) = \cos x - 2 \sin x$$

$$f''(x) = -\sin x - 2 \cos x$$

$$f'''(x) = -\cos x + 2 \sin x$$

Now, at $x=0$,

$$f(0) = \sin 0 + 2 \cos 0 = 2$$

$$f'(0) = \cos 0 - 2 \sin 0 = 1$$

$$f''(0) = -\sin 0 - 2 \cos 0 = -2$$

$$f'''(0) = -\cos 0 + 2 \sin 0 = -1$$

We know, $P_n(x) = \frac{f^0(x)}{0!}x^0 + \frac{f'(x)}{1!}x^1 + \frac{f''(x)}{2!}x^2 + \frac{f'''(x)}{3!}x^3 + \dots$

According to Taylor series around $x=0$,

$$P_3(x) = \frac{f^0(0)}{0!}x^0 + \frac{f'(0)}{1!}x^1 + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3$$

$$\Rightarrow P_3(x) = \frac{2}{0!}(x-0)^0 + \frac{1}{1!}(x-0)^1 + \frac{-2}{2!}(x-0)^2 + \frac{-1}{3!}(x-0)^3$$

$$\Rightarrow P_3(x) = 2 + (x-0) - (x-0)^2 - \frac{x^3}{6}$$

$$\Rightarrow P_3(x) = 2 + x - x^2 - \frac{1}{6}x^3$$

$$\therefore P_3(x) = 2 + x - x^2 - \frac{1}{6}x^3$$

Ans to the question no 02

We know,

$$P_n(0) = \frac{f^0(x)}{0!}x^0 + \frac{f'(x)}{1!}x^1 + \frac{f''(x)}{2!}x^2 + \frac{f'''(x)}{3!}x^3 + \dots$$

Given function,

$$f(x) = \ln(e^{2x} + e^{-2x})$$

By differentiating we get,

$$f'(x) = \frac{1}{e^{2x}+e^{-2x}} \cdot (2e^{2x} - 2e^{-2x}) = \frac{2(e^{4x}-1)}{e^{4x}+1}$$

$$f''(x) = \frac{2 \cdot 4e^{4x}(e^{4x}+1) - 4e^{4x}2(e^{4x}-1)}{(e^{4x}+1)^2}$$

$$\Rightarrow f''(x) = \frac{8e^{4x}(e^{4x}+1) - 8e^{4x}(e^{4x}-1)}{(e^{4x}+1)^2}$$

$$\Rightarrow f''(x) = \frac{8e^{4x}(e^{4x}-e^{4x}+2)}{(e^{4x}+1)^2}$$

$$\Rightarrow f''(x) = \frac{16e^{4x}}{(e^{4x}+1)^2}$$

Putting $x=0$ in $f(x)$ and its derivatives we get,

$$f(0) = \ln(e^{2 \cdot 0} + e^{-2 \cdot 0}) = \ln 2$$

$$f'(0) = \frac{2(e^{4 \cdot 0}-1)}{e^{4 \cdot 0}+1} = \frac{0}{2} = 0$$

$$f''(0) = \frac{16e^{4 \cdot 0}}{(e^{4 \cdot 0}+1)^2} = \frac{16}{4} = 4$$

Therefore,

$$P_0 = \frac{f^0(0)x^0}{0!} = \ln 2$$

$$P_1 = \frac{f^0(0)x^0}{0!} + \frac{f'(0)x^1}{1!} = \ln 2 + 0 = \ln 2$$

$$P_2 = \frac{f^0(0)x^0}{0!} + \frac{f'(0)x^1}{1!} + \frac{f''(0)x^2}{2!} = \ln 2 + 2x^2$$

Ans to the question no 03

Given multi-variable function,

$$f(x, y) = (x^2 + 2y^2x + 3x + y^2)^2$$

Derivatives of $f(x, y)$ with respect to x considering other variables constant,

$$f_x = 2(x^2 + 2y^2x + 3x + y^2)(2x + 2y^2 + 3)$$

$$f_{xx} = 2\{2(x^2 + 2y^2x + 3x + y^2) + (2x + 2y^2 + 3)^2\}$$

$$\Rightarrow f_{xx} = 2\{(2x^2 + 4xy^2 + 6x + 2y^2) + (4x^2 + 4y^4 + 9 + 8xy^2 + 12y^2 + 12x)\}$$

$$\Rightarrow f_{xx} = 2(6x^2 + 12xy^2 + 18x + 14y^2 + 4y^4 + 9)$$

$$\therefore f_{xx} = 2(6x^2 + 12xy^2 + 18x + 14y^2 + 4y^2 + 9)$$

Similarly,

$$f_y = 2(x^2 + 2y^2x + 3x + y^2)(4xy + 2y)$$

$$f_{yy} = 2\{(4x + 2)(x^2 + 2y^2x + 3x + y^2) + (4xy + 2y)^2\}$$

$$\Rightarrow f_{yy} = 2\{(4x^3 + 8x^2y^2 + 12x^2 + 4xy^2 + 2x^2 + 4xy^2 + 6x + 2y^2) + (16x^2y^2 + 4y^2 + 16xy^2)\}$$

$$\Rightarrow f_{yy} = 2(4x^3 + 24x^2y^2 + 14x^2 + 24xy^2 + 6x + 6y^2)$$

$$\therefore f_{yy} = 2(4x^3 + 24x^2y^2 + 14x^2 + 24xy^2 + 6x + 6y^2)$$

Ans to the question no 04

Given function,

$$f(x, y) = \ln((x^2 + 3y + xy)^2)$$

$$\text{Where } x(t) = 2t + 3, y(t) = t^2 + 3t$$

Chain rule for partial derivatives,

$$\frac{df}{dt} = \frac{\partial f}{\partial x} \cdot \frac{dx}{dt} + \frac{\partial f}{\partial y} \cdot \frac{dy}{dt}$$

Now,

$$\frac{\partial f}{\partial x} = \frac{1}{(x^2+3y+xy)^2} \cdot 2(x^2 + 3y + xy) \cdot (2x + y) = \frac{2(2x+y)}{(x^2+3y+xy)}$$

$$\frac{dx}{dt} = 2$$

$$\frac{\partial f}{\partial y} = \frac{1}{(x^2+3y+xy)^2} \cdot 2(x^2 + 3y + xy) \cdot (3 + x) = \frac{2(x+3)}{(x^2+3y+xy)}$$

$$\frac{dy}{dt} = 2t + 3$$

Imposing these values on the chain rule formula,

$$\frac{df}{dt} = \frac{2 \cdot 2(2x+y)}{(x^2+3y+xy)} + \frac{2(x+3)(2t+3)}{(x^2+3y+xy)}$$

Since we have to evaluate this expression when $t=0$,

$$\begin{aligned} & \frac{4(2x+y)}{(x^2+3y+xy)} + \frac{2(x+3)(2 \cdot 0+3)}{(x^2+3y+xy)} \\ &= \frac{8x+4y}{(x^2+3y+xy)} + \frac{6x+18}{(x^2+3y+xy)} \\ &= \frac{14x+4y+18}{(x^2+3y+xy)} \\ &\therefore \frac{df}{dt} = \frac{14x+4y+18}{(x^2+3y+xy)} \text{ when } t=0 \end{aligned}$$

Ans to the question no 05

Given function,

$$y^2 \sin(x^3) + ze^{3x} - \cos(z^2) = z^4 - y^2 + x^2$$

Imposing partial differentiation on both sides with respect to x ,

$$\begin{aligned}\frac{\partial}{\partial x}\{y^2 \sin(x^3) + ze^{3x} - \cos(z^2)\} &= \frac{\partial}{\partial x}(z^4 - y^2 + x^2) \\ \Rightarrow y^2 \cos(x^3) \cdot 3x^2 + \frac{\partial z}{\partial x}e^{3x} + z \cdot 3e^{3x} + \sin(z^2)2z\frac{\partial z}{\partial x} &= 4z^3\frac{\partial z}{\partial x} + 2x \\ \Rightarrow \frac{\partial z}{\partial x}(e^{3x} + 2z \sin(z^2) - 4z^3) &= 2x - z \cdot 3e^{3x} - 3x^2y^2 \cos(x^3) \\ \Rightarrow \frac{\partial z}{\partial x} &= \frac{2x - 3ze^{3x} - 3x^2y^2 \cos(x^3)}{(e^{3x} + 2z \sin(z^2) - 4z^3)} \\ \therefore \frac{\partial z}{\partial x} &= \frac{2x - 3ze^{3x} - 3x^2y^2 \cos(x^3)}{(e^{3x} + 2z \sin(z^2) - 4z^3)}\end{aligned}$$

Similarly partial differentiation on both sides with respect to y,

$$\begin{aligned}\frac{\partial}{\partial y}\{y^2 \sin(x^3) + ze^{3x} - \cos(z^2)\} &= \frac{\partial}{\partial y}(z^4 - y^2 + x^2) \\ \Rightarrow 2y \sin(x^3) + \frac{\partial z}{\partial y}e^{3x} + \sin(z^2)2z\frac{\partial z}{\partial y} &= 4z^3\frac{\partial z}{\partial y} - 2y \\ \Rightarrow \frac{\partial z}{\partial y}(e^{3x} + 2z \sin(z^2) - 4z^3) &= -2y - 2y \sin(x^3) \\ \Rightarrow \frac{\partial z}{\partial y} &= \frac{-2y(1 + \sin x^3)}{(e^{3x} + 2z \sin(z^2) - 4z^3)} \\ \therefore \frac{\partial z}{\partial y} &= \frac{-2y(1 + \sin x^3)}{(e^{3x} + 2z \sin(z^2) - 4z^3)}\end{aligned}$$

Ans to the question no 06

Given function,

$$h(x, y, z) = y^3 z^3 \cos(x^4) + x^3 \sin(z^2)$$

We have to use partial differentiation to find h_{yzx} .

Imposing partial differentiation of $h(x, y, x)$ with respect to y,

$$h_y = 3y^2 z^3 \cos(x^4)$$

Again partial differentiation of h_y with respect to z,

$$h_{yz} = 3y^2 3z^2 \cos(x^4) = 9y^2 z^2 \cos(x^4)$$

Finally partial differentiation of h_{yz} with respect to x,

$$h_{yzx} = 9y^2 z^2 \{-\sin(x^4)(4x^3)\} = 36x^3 y^2 z^2 \{-\sin(x^4)\}$$

evaluating the expression of h_{yzx} at point (0,1,1) we get,

$$h_{yzx} = 36 \cdot 0^3 \cdot 1^2 \cdot 1^2 \{-\sin(0^4)\} = 0$$