

# WEAK TYPE ESTIMATES OF RIESZ MEANS ON RIEMANNIAN SYMMETRIC SPACES

SHIJUN ZHENG

ABSTRACT. Let  $G$  be a noncompact semisimple Lie group of rank one,  $X = G/K$  its homogeneous space. We investigate the weak type estimates for the Bochner-Riesz means of functions in certain Hardy spaces on  $X$ . Solution to this problem relies on an explicit formula for the kernel of the Riesz means. The critical index are the same as in the Euclidean case. The Riesz means on  $X$  can also be defined via a weighted Jacobi transform. It is possible we can obtain a weak type estimate for the corresponding maximal operator. We also study the boundedness result for the Hardy spaces on  $G/K$  by an atomic decomposition adapted to  $G$ .

## 1. INTRODUCTION

The subject of  $L^p$  boundedness for Riesz means is one of the major problems in harmonic analysis. The Bochner-Riesz multiplier  $r_\delta = (1 - |x|^2)_+^\delta$ . It is known that  $\delta(p) = n(1/p - 1/2) - 1/2$  is the critical index for  $L^p$  problems (with  $1/p - 1/2 \geq 1/(n+1)$ ). It is showed in [Stein, Taibleson and Weiss] [62] that this is also the critical index for Hardy spaces. Namely, they proved that the maximal function associated to Riesz means maps  $H^p$  continuously to weak  $L^p$  for  $0 < p < 1$ . By interpolation we know this implies the  $L^p$  boundedness.

We are concerned with obtaining the analogous result on symmetric spaces. Previously, Clerc studied the Riesz means of Hardy spaces on compact Lie groups and he obtained the same critical index for the maximal Bochner-Riesz operator by proving that the maximal operator satisfies the weak type estimates for atoms, which serve as “building blocks” for Hardy spaces on  $G$ .

The study of Hardy spaces on a noncompact symmetric space is more difficult. One of the reason is that the definition of  $H^p$  classes on  $G$  is relatively not well understood. There are atomic decompositions

---

*Date:* August, 2003.

*2000 Mathematics Subject Classification.* Primary:43A15; Secondary:22E30.

*Key words and phrases.* Semisimple Lie group, symmetric space, Bochner-Riesz means, Hardy spaces.

for  $H^p$  classes on  $R^n$ , compact Lie groups or compact manifolds as well as Heisenberg groups [Coifman-Weiss][Clerc],[Folland-Stein] and [H.P. Liu]. However so far atomic decomposition is known only for  $K$ -invariant functions on  $G$  as studied in [T.Kawazoe, 1985, 2000]. Also we do not know whether the atomic definition is equivalent to the Poisson function characterization on a symmetric space.

Let  $X = G/K$  be the homogeneous space for a connected semisimple Lie group  $G$ , where  $K$  is a maximal compact subgroup of  $G$ . Let  $G = KAN$  and  $\mathfrak{g} = \mathfrak{k} + \mathfrak{a} + \mathfrak{n}$  be Iwasawa decomposition for  $G$  and its Lie algebra  $\mathfrak{g}$ . Let  $\mathcal{L}$  be the Laplacian-Beltrami operator on  $G$ . In rank one case  $\mathcal{L} = \Delta^{-1} \frac{d}{dx} (\Delta \frac{d}{dx})$ , where  $\Delta(x) = (2 \sinh x)^{2\alpha+1} (2 \cosh x)^{2\beta+1}$  for radial functions.

The sperical functions on  $G$  are  $K$ -invariant and satisfies

$$\mathcal{L}\phi_\lambda = -(\lambda^2 + \rho^2)\phi_\lambda,$$

with  $\rho = \alpha + \beta + 1$ , where  $2\alpha = m_1 + m_2 - 1$  and  $2\beta = m_2 - 1$  and so  $\rho = (m_1 + 2m_2)/2$  is equal to half of the sum of the positive roots with multiplicities with respect to the pair  $(\mathfrak{g}, \mathfrak{a})$ . There are integral formulas for the spectral operator  $m(\mathcal{L})$  in [Giulini-Mauceri], for complex or rank one semisimple Lie groups. We shall study the Bochner-Riesz operators near the critical index.

## 2. PRELIMINARIES

Definition. A function is said to be  $K$  bi-invariant if  $f$  is invariant under left and right translation by  $K$ .

Denote the Fourier (spherical) transform of  $f$  by

$$\hat{f} = \int_G f(g) \bar{\phi}_\lambda(g) dg.$$

The inversion formula is given by

$$f(g) = \int_0^\infty \phi_\lambda(g) \hat{f}(\lambda) d\mu(\lambda),$$

where  $d\mu(\lambda) = |c(\lambda)|^{-2} d\lambda$  is a measure on  $R^+$ . (Gangolli, R. Ann. Math., 93(1971),150-165).

Sperical multiplier. To a function  $m \in L^\infty(R^+)$  we associate a map  $T_m : C_c(G/K) \rightarrow L^2(G/K)$  by

$$T_m f(g) = \int_0^\infty m(\lambda) \phi_\lambda(g) \hat{f}(\lambda) |c(\lambda)|^{-2} d\lambda.$$

( $T_m f := m^\vee * f$  can be viewed as a distribution in  $S'(G/K)$ .)

When  $m_z(x) = (1 - \frac{x^2 + \rho^2}{R})^z$  we have the Bochner-Riesz multiplier of order  $z$ . If  $-\mathcal{L} = \int_{\rho^2}^{\infty} \lambda dE_{\lambda}$  is the spectral resolution of the unique self-adjoint extension of  $-\mathcal{L}$  (using the same notation for its extension), then Bochner-Riesz operator can be written as for  $\lambda \geq \rho^2$

$$S_R^z = \int_{\rho^2}^{\infty} (1 - \frac{\lambda}{R})_+^z dE_{\lambda}.$$

We can represent  $S_R^z$  by convolution on  $G$ :  $S_R^z f = f * r_R^z$  with the kernel  $r_R^z$  being  $C^{\infty}$  smooth. The spherical transform of this kernel is exactly  $m_z$ . In Giulini-Mauceri, it was proved that if  $1 \leq p \leq 2$  and if  $Re z > \delta(p) := (\frac{1}{p} - \frac{1}{2})(n-1)$ , then the maximal operator  $S_*^z$  maps  $L^p(X)$  continuous into  $L^{p+r}(X)$  with  $r$  large enough, hence they obtained the almost everywhere convergence for  $L^p$  functions.

We are concerned with studying the  $H^p$  boundedness of Riesz means for  $G$ . There are two ways to address this issue. First we can consider the Riesz mean on the atomic Hardy spaces introduced in T.Kawazoe[2000] for  $K$ -invariant functions. The second way is to study the Riesz means for Hardy spaces defined by "Poisson" semigroup  $e^{-t\mathcal{L}}$ . In this regard, we may refer to the paper on general affine symmetric spaces (which are non-Riemannian) by Hilgert, Olafsson and Orsted [1991].

We mention that in Riemannian case Kawazoe studied in [38] the properties of Poisson kernels related to the Laplacian-Beltrami operator on  $G/K$ . The Poisson kernel  $P$  is defined as a function on  $G/K \times K/M$ :  $P(gK, kM) = \exp(-2\rho(H(g^{-1}k)))$ , where  $G/k$  is identified with the unit ball in  $\mathbb{F}^n$  ( $\mathbb{F}^n$  is  $\mathbb{R}$ ,  $\mathbb{C}$  or  $\mathbb{H}$ ), and  $K/M$  with its boundary, the unit sphere. This suggests that the Hardy classes introduced in the paper by Olafsson et al. are extensions to the non-Riemannian case as they defined the Hardy spaces by Poisson kernels of the above type.

It seems interesting and desirable if we can construct an atomic decomposition for this type of Hardy spaces (which fit naturally with our symmetric spaces). Therefore we would be able to study in a simpler way the mapping properties of  $B$ - $R$  means, multiplier operators for  $H^p$  spaces on symmetric spaces.

Examples of semisimple Lie groups of real rank one.

$SL(2, \mathbb{R})$ ,  $SU(n, 1)$ ,  $SO(n, 1)$ ,  $Sp(n, 1)$ , the connected Lie group of real type  $F_4$ .

Let  $X = G/K$ . The classification of real rank one noncompact symmetric spaces is known: a.  $G = SO_o(n, 1)$ ,  $K = SO(n)$ ,  $n \in \mathbb{Z}^+$ ,  $n \geq 2$

b.  $G = SU(n, 1)$ ,  $K = S(U(n) \times U(1))$ ,  $n \in \mathbb{Z}^+$ ,  $n \geq 2$

- c.  $G = SP(n, 1)$ ,  $K = SP(n) \times SP(1)$ ,  $n \in \mathbb{Z}^+$ ,  $n \geq 2$
- d.  $G = F_4(-20)$ ,  $K = \text{spin } 9$ .

The first three kinds can be identified as hyperbolic spaces over real, complex or quaternian fields ( $\mathbb{R}$ ,  $\mathbb{C}$  or  $\mathbb{H}$ ).

Let  $X = G/K$  be a Riemannian symmetric space, where  $G$  is a connected semisimple Lie group with finite center and  $K$  a maximal compact subgroup. In the Lie algebra level, we have the Cartan decomposition  $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$  with a Cartan involution  $\theta$ . Let  $\mathfrak{a}$  be a maximal abelian subspace of  $\mathfrak{p}$ . Denote by  $\mathfrak{a}_{\mathbb{C}}^*$  the complexification of  $\mathfrak{a}^*$ .

Let  $P = MAN$  be a minimal parabolic subgroup of  $G$ , where  $M$  is the centralizer of  $A$  in  $K$ . Then the compact  $G/P = K/M$  is the maximal boundary in the compactification of  $G/K$ .

The spherical function can also be defined via Poisson integral of constant function 1 in  $\mathcal{B}(K/M)$  where  $\mathcal{B}(K/M)$  is the space of the hyperfunctions on  $K/M$ :

$$\phi_{\lambda}(gK) = \int_K e^{-\langle \lambda + \rho, \log(g^{-1}k) \rangle} dk.$$

We have  $\lim_{\alpha \rightarrow \infty} \alpha^{-\lambda + \rho} \phi_{\lambda}(\alpha) = c(\lambda)$ , Harish-Chandra's  $c$ -function.

For elementary spherical functions ( $K$ -bi-invariant), there is an explicit formula ( $G$  being complex) ([22]):

$$\phi_{\lambda}(\exp H) = c(i\lambda) \frac{\sum_w (\det w) e^{-w\lambda(H)}}{\prod_{\alpha \in \Sigma_+} \sinh \alpha(H)} \quad H \in \mathfrak{a} \quad \lambda \in \mathfrak{a}_{\mathbb{C}}^*,$$

and

$$c(\lambda) = \pi(\rho) / \pi(i\lambda) \quad \lambda \in \mathfrak{a}^*,$$

where  $\pi(z) = \prod_{\alpha \in \Sigma_+} < \alpha, z >$ .

In the end of this section we refer to the following paper by S. Ben Saïd, T. Oshima and N. Shimeno, related to our project. "Fatou's theorems and Hardy-type spaces for eigenfunctions of the invariant differential operators on symmetric spaces" in *IMRN*, **16** (2003), 915-931.

### 3. RIESZ MEANS OF SPHERICAL EXPANSIONS

For  $z \in \mathbb{C}$ ,  $\Re z \geq 0$ , we define the Bochner-Riesz means of order  $z$  by

$$S_R^z f(x) = \int_0^\infty \left(1 - \frac{\lambda^2 + \rho^2}{R}\right)_+^z \hat{f}(\lambda) \phi_{\lambda}(x) |c(\lambda)|^{-2} d\lambda.$$

The Riesz means can be written as  $S_R^z f = s_R^z * f$ , here  $s_R^z$  is a function whose spherical transform is

$$\hat{s}_R^z(\lambda) = (1 - \frac{\lambda^2 + \rho^2}{R})_+^z.$$

Let  $\mathcal{F}_c$  be the usual Fourier-cosine transform,  $\mathcal{A}$  the Abel transform, then  $s_R^z = \mathcal{A}^{-1} \mathcal{F}_c^{-1}(\hat{s}_R^z(\lambda))$ .

Let  $\alpha$  and  $2\alpha$  be two roots of  $(\mathfrak{g}, \mathfrak{a})$  with multiplicities  $m_\alpha$  and  $m_{2\alpha}$  respectively. Then dimension of  $G/K$  is equal to  $n = m_\alpha + m_{2\alpha} + 1$ . Fix an element  $H_0$  in  $\mathfrak{a}$  with  $\alpha(H_0) = 1$ . Let  $G = K\bar{A}^+K$  be Cartan decomposition, where  $A^+ = \{\exp(tH_0) : t > 0\}$ .

For  $K$ -bi-invariant function  $f$ , if the spherical transform is given by  $\hat{f}(\lambda) = \int_0^\infty f(\exp(tH_0))\phi_{-\lambda}(\exp(tH_0))\Delta(t)dt$  then we have the inversion formula

$$f(t) := f(\exp(tH_0)) = \int_0^\infty \hat{f}(\lambda)\phi_\lambda(t)|c(\lambda)|^{-2}d\lambda,$$

noting that as  $\lambda \rightarrow \infty$ ,  $|c(\lambda)| \sim \lambda^{-(n-1)/2}$  (when  $G$  has rank one).

We need the asymptotic estimates for spherical functions (cf. Stanton-Tomas or Giulini-Mauceri-Meda): Let  $\mathcal{J}_\nu(x) = x^{-\nu}J_\nu(x)$ , where  $J_\nu$  is the Bessel function of the first kind and of order  $\nu$ . We have for  $t \leq T_0$  ( $T_0 > 1$ )

$$\phi_\lambda(\exp(tH_0)) = t^{(n-1)/2}\Delta(t)^{-1/2}(\mathcal{J}_{n/2-1}(\lambda t) + t^2a(t)\mathcal{J}_{n/2}(\lambda t) + E(\lambda, t)),$$

where  $a$  is bounded and smooth, and the error term  $E$  verifies

$$|E(\lambda, t)| \leq C \begin{cases} t^4 & \text{if } |\lambda t| \leq 1, \\ t^4(\lambda t)^{-(n+3)/2} & \text{if } |\lambda t| > 1 \end{cases}$$

We also need the estimates for derivatives of  $|c(\lambda)|^{-2}$ .

$$\frac{d^k}{d\lambda^k}(|c(\lambda)|^{-2}) \leq c_k(1 + |\lambda|)^{n-k-1}.$$

whose estimates are based on the expression

$$c(\lambda) = \Gamma(a/2)\Gamma(b/2)\frac{\Gamma(i\lambda)\Gamma(\frac{a+i\lambda}{2})}{\Gamma(\frac{a}{2} + i\lambda)\Gamma(\frac{\rho+i\lambda}{2})},$$

here  $a = m_\alpha$ ,  $b = m_{2\alpha}$  and  $\rho = (m_\alpha + 2m_{2\alpha})/2$ ;

Similarly, we can define the Bochner-Riesz means of order  $z$  associated with Jacobi expansion by

$$S_R^z f(x) = \int_0^\infty (1 - \frac{\lambda^2 + \rho^2}{R})_+^z \hat{f}(\lambda)\phi_\lambda^{\alpha, \beta}(x)|c(\lambda)|^{-2}d\lambda.$$

We can write  $S_R^z f = s_R^z * f$ , where  $s_R^z$  is a function whose Jacobi transform is

$$\hat{s}_R^z(\lambda) = (1 - \frac{\lambda^2 + \rho^2}{R})_+^z.$$

We also have the relation between spherical transform and Abel transform via Fourier transform:  $\mathcal{F}_c \mathcal{A} s_R^z = \hat{s}_R^z(\lambda)$ .

Like in Koornwinder, Jacobi functions and analysis on noncompact semisimple Lie Groups of rank one, in R.Askey et al (eds.), Special functions, D.Reidel Publ. Com., Dordrecht. (1984) and Giulini.S and Mauceri G. ‘Almost everywhere convergence of Riesz means on certain noncompact symmetric spaces’, Ann. Mat.Pura. Appl.(4)159, (1991). We obtain

**Proposition 3.1.** *If  $\alpha \geq \beta - 1/2$ ,  $\alpha \neq -1/2$ , then*

$$|s_R^z(x)| \leq c(z) R^{\alpha+1} (1 + \sqrt{R}|x|)^{-Re z - \alpha - 3/2} \left(\frac{x}{\sinh x}\right)^{\alpha+1/2} (\cosh x)^{\beta+1/2}.$$

Consider the maximal operator

$$S_*^z f(x) = \sup_{R>0} |S_R^z f(x)|, \quad f \in L^p(\Delta(x)dx), 1 \leq p \leq 2.$$

We will use the maximal function  $Mf$ :

$$Mf(x) = M_{\alpha,\beta} f(x) = \sup_{r>0} \chi_r * |f|(x).$$

Here  $\chi_r = \frac{1}{|B(r)|} \chi_{[-r,r]}$ ,  $\chi_{[-r,r]}$  is the characteristic function of  $[-r, r]$ .

Note. For certain discrete  $\alpha, \beta$ ,  $Mf$  has group-theoretical interpretation as the maximal function on symmetric spaces of rank one.

**Theorem 3.2.** *Let  $\alpha \geq \beta \geq -1/2$  and  $\alpha \neq -1/2$ , the maximal operator  $M$  has weak type  $(1, 1)$ ,*

$$|\{x : Mf(x) > \lambda\}| \leq C \lambda^{-1} \|f\|_{L^1(\Delta dx)}.$$

Define

$$(1) \quad M_\phi f(x) = \sup_{t>0} \phi(t) \chi_{[-t,t]} * f(x)$$

$$(2) \quad = \sup_{t>0} \phi(t) \int_0^t T_x f(y) \Delta(y) dy.$$

**Theorem 3.3.** *Let  $\alpha \geq \beta - 1/2$ ,  $\alpha \neq -1/2$ . i) If  $Re z > 0$ , then for  $f \in L^2(\Delta(x)dx)$ ,*

$$\|S_*^z f\|_2 \leq c(z) \|f\|_2.$$

*ii) If  $Re z > \alpha + \frac{1}{2}$ ,  $S_*^z$  is bounded from  $L^1(\Delta(x)dx)$  to weak  $L^1 + L^q$  for every  $q \in [4, \infty]$ .*

iii) If  $\operatorname{Re} z > (\frac{2}{p} - 1)(\alpha + \frac{1}{2})$  and  $1 < p \leq 2$ , then for every  $q \geq \frac{4p}{3p-2}$ ,

$$\|S_*^z f\|_{p+q} \leq c(z)\|f\|_p, \quad f \in L^p(\mathbb{R}^+, \Delta(x)dx).$$

Proof similar to that of Lemma 4.1 in Giulini and Mauceri, 1991.  
(iii) follows from Stein's complex interpolation.

A direct corollary of the above theorem is the pointwise result.

**Theorem 3.4.** *Let  $1 \leq p \leq 2$  and  $f \in L^p(\mathbb{R}^+, \Delta(x)dx)$ . If  $\operatorname{Re} z > (\frac{2}{p} - 1)(\alpha + \frac{1}{2})$ , then  $\lim_{R \rightarrow \infty} S_R^z f(x) = f(x)$  a.e.*

The proof is based on the estimate of the kernel of  $S_R^z$  given by

$$S_R^z(x, y) = \int_0^\infty (1 - \frac{\lambda^2 + \rho^2}{R})_+^z \phi_\lambda^{\alpha, \beta}(x) \phi_\lambda^{\alpha, \beta}(y) |c(\lambda)|^{-2} d\lambda.$$

and some basic properties in the following lemma concerning measure of geodesic balls in  $X = G/K$ .

**Lemma 3.5.** *Define  $B(x, r)$  by  $B(x, r) = [x - r, x + r]$  if  $x - r > 0$ ;  $[0, x + r] := B(x + r)$  if  $x - r \leq 0$ . Then*

- 1) *If  $0 < r < \infty$ ,  $1 + |B(r)| \sim (\cosh r)^{2\rho}$ .*
- 2)

$$|B(r)| \sim r^{2(\alpha+1)} \quad r \leq C$$

$$|B(r)| \sim (\cosh r)^{2\rho} \quad r \geq C.$$

- 3) *For  $r \leq 1$ ,  $x > 2$ ,  $|B(x, r)| \sim r(\cosh x)^{2\rho} \sim r e^{2\rho x}$ .*
- 4) *For  $r \leq 1$ ,  $x < 2$ ,*

$$|B(x, r)| \sim r x^{2(\alpha+1)} \quad x \geq 2r$$

$$|B(x, r)| \sim |B(r)|, \quad x < 2r.$$

**Lemma 3.6.** *Let  $E$  be a subset of  $\mathbb{R}^+$ .  $\{B(x_k, r_k)\}$  is a covering of  $E$  with  $r_k \leq 1$  for all  $k$ . Then there exists a disjoint  $\{B(x_i, r_i)\}$  such that*

$$|E| \leq C \sum_{i=1}^\infty |B(x_i, r_i)|.$$

Proof of Lemma 3.6 is similar to that of the Vitali covering lemma on space of homogeneous type e.g. [1].

## REFERENCES

- [1] J.-Ph. Anker , Sharp estimates for some functions of the Laplacian on non-compact symmetric spaces. *Duke Math. J.* 65 (1992), pp. 257-297.
- [2] R. A. Askey, T. H. Koornwinder and W. Schempp, *Special functions: group theoretical aspects and applications*, Edited by R. A. Askey, T. H. Koornwinder and W. Schempp. *Mathematics and its Applications*, D. Reidel Publishing Co., Dordrecht, 1984.
- [3] S.Ben Saïd, T.Oshima, N.Shimeno, Fatou's theorems and Hardy-type spaces for eigenfunctions of the invariant differential operators on symmetric spaces, *Int. Math. Res. Not.* (2003), **16**, 915–931.
- [4] M. Christ and C. D. Sogge, The weak type  $L^1$  convergence of eigenfunction expansions for pseudodifferential operators, *Invent. Math.* **94** (1988), 421–453.
- [5] J.-L. Clerc, Bochner-Riesz means of  $H^p$  functions ( $0 < p < 1$ ) on compact Lie groups. Noncommutative harmonic analysis and Lie groups (Marseille-Luminy, 1985), 86–107, *Lecture Notes in Math.*, 1243, Springer, Berlin, 1987. (Reviewer: Fulvio Ricci)
- [6] J. L. Clerc and E. M. Stein ,  $L^p$ -multipliers for non-compact symmetric spaces. *Proc. Nat. Acad. Sci. USA* 71 (1974), pp. 3911-3912.
- [7] Colzani, Leonardo, Riesz means of eigenfunction expansions of elliptic differential operators on compact manifolds, *Rend. Sem. Mat. Fis. Milano* **58** (1988), 149–167 (1990).
- [CoT90] L. Colzani, G. Travaglini, Hardy-Lorentz spaces and expansions in eigenfunctions of the Laplace-Beltrami operator on compact manifolds, *Colloq. Math.* **58** (1990), no. 2, 305–315.
- [8] L.Colzani, et. al, Equiconvergence theorem for Fourier-Bessel expansions with applications to the harmonic analysis of radial functions in  $R^n$  and noneuclidean spaces, *Trans. Amer.Math.Soc.*, **338**(1), (1993), 43–.
- [9] Cowling, Michael G. Harmonic analysis on semigroups, *Ann. of Math.(2)*, **117**, (1983), no. 2, 267–283.
- [10] Cowling, Michael, The Kunze-Stein phenomenon, *Ann. Math.(2)*, **107**(2), (1978), 209–234.
- [11] Cowling, Michael; Giulini, S.; Meda, Stefano Oscillatory multipliers related to the wave equation on noncompact symmetric spaces, *J. London Math. Soc.*, (2) 66 (2002), no. 3, 691–709.
- [12] Cowling, Michael; Giulini, Saverio; Meda, Stefano,  $L^p$ - $L^q$  estimates for functions of the Laplace-Beltrami operator on noncompact symmetric spaces. I. *Duke Math. J.* 72 (1993), no. 1, 109–150.
- [13] Cowling, Michael; Giulini, Saverio; Meda, Stefano  $L^p$ - $L^q$ -estimates for functions of the Laplace-Beltrami operator on noncompact symmetric spaces. II. *J. Lie Theory* 5 (1995), no. 1, 1–14.
- [14] Cowling, M.; Giulini, S.; Meda, S.  $L^p$ - $L^q$  estimates for functions of the Laplace-Beltrami operator on noncompact symmetric spaces. III. *Ann. Inst. Fourier (Grenoble)* 51 (2001), no. 4, 1047–1069.
- [15] Cowling, Michael, Sikora, Adam, A spectral multiplier theorem for a sublaplacian on  $SU(2)$ , *Math. Z.*, **238**(1), (2001), 1–36.



- [16] De Michele, Leonede; Mauceri, Giancarlo  $H^p$  multipliers on stratified groups, *Ann. Mat. Pura Appl.*, **148**(4), (1987), 353–366. (Reviewer: Gerald B. Folland)
- [17] M. Flensted-Jensen, Paley-Wiener type theorems for a differential operator connected with symmetric spaces. *Ark. Mat.* 10 (1972), pp. 143–162. Math-SciNet
- [18] Flensted-Jensen, Mogens; Koornwinder, Tom, The convolution structure for Jacobi function expansions, *Ark. Mat.*, **11**, 245–262. (1973).
- [19] Folland, Gerald B., *Harmonic analysis in phase space*, *Annals of Mathematics Studies*, **122**. Princeton University Press, Princeton, NJ, 1989.
- [20] G. B. Folland and E. M. Stein, *Hardy Spaces on Homogeneous Groups*, **Math. Notes**, **28**, Princeton Univ. Press, Princeton (1982).
- [21] Giulini, Saverio; Mauceri, Giancarlo, Almost everywhere convergence of Riesz means on certain noncompact symmetric spaces, *Ann. Mat. Pura Appl.* (4)**159** (1991), 357–369.
- [22] Giulini, Saverio; Mauceri, Giancarlo; Meda, Stefano  $L^p$  multipliers on noncompact symmetric spaces, *J. Reine Angew. Math.* **482** (1997), 151–175.
- [23] Giulini, Saverio(I-GENOE); Sjgren, Peter, A note on maximal functions on a solvable Lie group. *Arch. Math. (Basel)*, **55**(2), (1990), 156–160.
- [24] S. Helgason, *Differential Geometry and Symmetric Spaces*, Academic Press, San Diego (1962).
- [25] Hilgert, J., Ólafsson, G., Orsted, B., Hardy spaces on affine symmetric spaces. *J. Reine Angew. Math.*, **415** (1991), 189–218.
- [26] A. Hulanicki, J.W. Jenkins, Almost everywhere summability on nilmanifolds, *Trans. Amer. Math. Soc* **278** (1983), no. 2, 703–715.
- [27] ——— Nilpotent Lie groups and summability of eigenfunction expansions of Schrödinger operators. *Studia Math.* 80 (1984), no. 3, 235–244.
- [28] ——— Nilpotent Lie groups and eigenfunction expansions of Schrödinger operators. II. *Studia Math.* 87 (1987), no. 3, 239–252.
- [29] Alexandru D. Ionescu, Fourier integral operators on noncompact symmetric spaces of real rank one. *Journal of Functional Analysis* 174, 274–300 (2000).
- [30] Alexandru D. Ionescu, An endpoint estimate for the Kunze-Stein phenomenon and related maximal operators. *Annals of Mathematics* 152, 259–275 (2000).
- [31] Alexandru D. Ionescu, A maximal operator and a covering lemma on noncompact symmetric spaces. *Mathematical Research Letters* 7, 83–93 (2000).
- [32] Alexandru D. Ionescu, On the Poisson transform on symmetric spaces of real rank one. *Journal of Functional Analysis* 174, 513–523 (2000).
- [33] Alexandru D. Ionescu, Singular integrals on symmetric spaces of real rank one. To appear in *Duke Mathematical Journal*.
- [34] Alexandru D. Ionescu, Singular integrals on symmetric spaces, II. Preprint 2001.
- [35] Alexandru D. Ionescu and David Jerison, On the absence of positive eigenvalues of Schrödinger operators with rough potentials. Preprint 2002.
- [36] Kawazoe, T., Hardy spaces and maximal operators on real rank one semisimple Lie groups, *I, Tohoku Math. J.(2)*, **52**(1), (2000), 1–18.
- [37] Kawazoe, T.,  $L^1$  estimates for maximal functions and Riesz transform on real rank 1 semisimple Lie groups. *J. Funct. Anal.*, **157**(2), (1998), 327–357.
- [38] Kawazoe, T., A characterization of the Poisson kernel on the classical real rank one symmetric spaces, *Tokyo J. Math.*, **15**(2), (1992), 365–379.

- [39] Kawazoe, T., Atomic Hardy spaces on semisimple Lie groups, *Japan. J. Math. (N.S.)*, **11**(2), (1985), 293–343.
- [40] Kawazoe, T., Atomic Hardy spaces on semisimple Lie groups, *Noncommutative harmonic analysis and Lie groups* (Marseille-Luminy, 1985), 189–197, *Lecture Notes in Math.*, **1243**, Springer, Berlin, 1987.
- [41] ——— Maximal functions on non-compact rank one symmetric spaces, (radial maximal functions and atoms), *Group Representations and Systems of Differential Equations, Advanced Studies in Pure Math.*, **4** (1984), 121–138.
- [42] ——— Wavelet transform associated to an induced representation of  $SL(n+2, \mathbf{R})$ , *Ann. Inst. H. Poincaré Phys. Théor.*, **65**(1), (1996), 1–13.
- [43] Kawazoe, T., Wavelet transforms associated to a principal series representation of semisimple Lie groups. I, II. *Proc. Japan Acad. Ser. A Math. Sci.*, **71**(7), (1995), 154–157, 158–160.
- [44] Kawazoe, T.,  $KA$ -wavelets on semisimple Lie groups and quasi-orthogonality of matrix coefficients. *Infinite dimensional harmonic analysis* (Kyoto, 1999), 238–252, Grbner, Altendorf, 2000.
- [45] Kawazoe, T.; Liu, J. On a weak  $L^1$  property of maximal operators on non-compact semisimple Lie groups. *Tokyo J. Math.*, **25**(1), (2002), 165–180.
- [46] Kawazoe, T., Tahani, Taoufiq, A characterization of the Poisson kernel associated with  $SU(1, n)$ , *Tokyo J. Math.*, **11**(1), (1988), 37–55. (Reviewer: Jacques Faraut)
- [47] Kawazoe, T., Tahani, Taoufiq, Correction to: "A characterization of the Poisson kernel associated with  $SU(1, n)$ " [*Tokyo J. Math.* 11 (1988), no. 1, 37–55; MR 89h:22016], *Tokyo J. Math.*, **17**(2), (1994), 495–497. (Reviewer: Jacques Faraut)
- [48] Kenig, C. E.; Stanton, R. J.; Tomas, P. A. Divergence of eigenfunction expansions. *J. Funct. Anal.* 46 (1982), no. 1, 28–44.
- [49] Koornwinder, Tom H. Jacobi functions and analysis on noncompact semisimple Lie groups. *Special functions: group theoretical aspects and applications*, 1–85, *Math. Appl.*, Reidel, Dordrecht, 1984.
- [50] Koornwinder, Tom H. Fast wavelet transforms and Caldern-Zygmund operators. *Wavelets: an elementary treatment of theory and applications*, 161–182, *Ser. Approx. Decompos.*, 1, World Sci. Publishing, River Edge, NJ, 1993.
- [51] Liu, Heping, The  $H^p$  boundedness of Riesz means on the Heisenberg group, *Acta Math. Sinica* (N.S.) **11** (1995), Special Issue, 18–25.
- [52] Liu, He Ping; Lu, Shan Zhen The  $H^p$ -boundedness of the Riesz means for the eigenfunction expansions for elliptic operators with constant coefficients. *Proc. Amer. Math. Soc.* 122 (1994), no. 2, 487–494. (Reviewer: Gnter Berger)
- [53] Liu, Jianming, Maximal functions associated with the Jacobi transform, *Bull. London Math. Soc.*, **32**(5), (2000), 582–588.
- [54] N. Lohou, Transforms de Riesz et fonctions sommables, *Amer. J. Math.*, **114**, (1992), pp. 875–9.
- [55] C.Meaney, E.Prestini, Bochner-Riesz means on symmetric spaces. (English. English summary) *Tohoku Math. J. (2)* **50** (1998), no. 4, 557–570.
- [56] C.Meaney, E.Prestini, Almost everywhere convergence of the inverse spherical transform on  $SL(2, \mathbf{R})$ , *Ark. Mat.*, **32** (1994), no. 1, 195–211.

- [57] Prestini, Elena, Convolution with oscillating kernels: Bochner-Riesz and spherical partial sums operators, *Rend. Sem. Mat. Fis. Milano* **58** (1988), 81–89 (1990).
- [Sog87] Sogge, Christopher D. On the convergence of Riesz means on compact manifolds, *Ann. of Math. (2)* **126** (1987), no. 2, 439–447.
- [58] Sogge, Christopher D., *Eigenfunction and Bochner Riesz estimates on manifolds with boundary*, Math. Res. Lett **9** (2002), no. 2-3, 205–216.
- [59] Stanton, Robert J.; Tomas, Peter A. Expansions for spherical functions on noncompact symmetric spaces. *Acta Math.* 140 (1978), no. 3-4, 251–276.
- [60] Stanton, Robert J.; Tomas, Peter A. Pointwise inversion of the spherical transform on  $L^p(G/K)$ ,  $1 \leq p < 2$ . *Proc. Amer. Math. Soc.*, **73**(3), (1979), 398–404.
- [61] E. M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton Math. Ser. 43, Princeton Univ. Press, Princeton (1993).
- [62] E.M. Stein, M.H. Taibleson, G. Weiss, Weak type estimates for maximal operators on certain  $H^p$  classes, *Proceedings of the Seminar on Harmonic Analysis* (Pisa, 1980). *Rend. Circ. Mat. Palermo* (2) 1981, suppl. 1, 81–97.
- [63] J. Strömberg, Weak type estimates for maximal functions on non-compact symmetric spaces, *Ann. of Math.*, **114**, (1981), pp. 115–126.
- [64] M. H. Taibleson and G. Weiss, The molecule characterization of certain Hardy spaces, *Astisque*, **77**, (1980), pp. 68–149.
- [65] G. Warner, *Harmonic Analysis on Semi-Simple Lie Groups II*, Springer-Verlag, Berlin/New York (1972).

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE,  
LA 70803

URL, Shijun Zheng: <http://www.math.lsu.edu/~szheng>