

• *Algebraic Limits*

46. Find the limits.

(a) $\lim_{x \rightarrow -1} (2x^2 - 4x + 4)$

(b) $\lim_{x \rightarrow -3} \frac{x^2 + 2x + 2}{x - x^3}$

(c) $\lim_{x \rightarrow -3} \sqrt{\frac{x^2 + 2x + 2}{x - x^3}}$

47. Find the limits.

(a) $\lim_{x \rightarrow -1} \frac{x^2 - x - 2}{x^2 - 1}$

(b) $\lim_{x \rightarrow -3} \frac{x + 3}{x^2 + 4x + 3}$

(c) $\lim_{x \rightarrow 4} \frac{x^2 - 3x - 4}{x^2 - 5x + 4}$

• *Continuity*

48. **Definition.** A function f is **continuous** at $x = a$ if: (a) $f(a)$ exists, (b) $\lim_{x \rightarrow a} f(x)$ exists, and (c) $\lim_{x \rightarrow a} f(x) = f(a)$.

A function is **continuous over an interval I** if it is continuous at each point a in I . If f is not continuous at $x = a$, we say that f is **discontinuous**, or has a **discontinuity**, at $x = a$.

49. The graph of the function $y = f(x)$ is given as follows. At what points of x is f **not** continuous?


50. Determine whether the function is continuous at the given point.

(a) $f(x) = x^3 + 2x, \quad \text{at } x = 2$

(b) $f(x) = \begin{cases} 2x - 3, & \text{for } x \geq 3, \\ 12 - x^2, & \text{for } x < 3, \end{cases} \quad \text{at } x = 3$

(c) $f(x) = \begin{cases} \frac{x^2 - 1}{x - 1}, & \text{for } x \neq 1, \\ 2, & \text{for } x = 1, \end{cases} \quad \text{at } x = 1$

• *Difference Quotient and Derivatives*



51. The following table shows total production of suits at a company during one morning work. What was the average number of suits produced per hour from 9 am to 11 am?

Time (number of hours since 8 am)	0	1	2	3	4
Total number of suits produced	0	20	55	64	100

52. **Definition.** The average rate of change of $f(x)$ with respect to x is also called the **difference quotient**. It is given by

$$\frac{\Delta y}{\Delta x} = \frac{\Delta f(x)}{\Delta x} := \frac{f(x+h) - f(x)}{h}, \quad \text{where } h \neq 0.$$

53. For $f(x) = x^2$, find the difference quotient for $x = 1$ and

- (a) $h = 1$, $h = 0.5$, $h = 0.2$, and $h = 0.1$ (h approaches 0 from the right)
 (b) $h = -1$, $h = -0.5$, $h = -0.2$, and $h = -0.1$ (h approaches 0 from the left)

54. **Definition.** For a function $y = f(x)$, its **derivative** at x is the function f' (also written as $\frac{dy}{dx}$ or $\frac{df(x)}{dx}$) defined by

$$f'(x) = \frac{df(x)}{dx} := \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h},$$

provided the limit exists. If $f'(x)$ exists, then we say that f is **differentiable** at x . The process of finding a derivative is called **differentiation**.

Geometrically, the derivative of f at x is the slope of the tangent line at $(x, f(x))$. This limit is also called the **instantaneous rate of change** of f at x .

55. For each function $f(x)$, first find $f'(x)$, then find $f'(-3)$ and $f'(2)$.

- (a) $f(x) = 4x - 2$
 (b) $f(x) = \frac{1}{x}$
 (c) $f(x) = x^2$