

# Anabaena: A New Chassis for Space Exploration

## Abstract

The sustainable exploration of deep space, particularly Mars, demands revolutionary approaches to life support that transcend traditional supply chain dependencies on Earth. Among promising biological systems, the filamentous cyanobacterium *Anabaena* has emerged as a compelling chassis for in situ resource utilization (ISRU) and bioregenerative life support systems (BLSS). This review critically examines the unique physiological, metabolic, and genetic attributes that position *Anabaena* as a cornerstone organism for extraterrestrial biotechnology. We analyze recent advances in understanding heterocyst differentiation, nitrogen fixation under Martian-analog conditions, and the development of synthetic biology tools for *Anabaena* engineering. Key challenges—including radiation tolerance, low-pressure cultivation, and genetic stability—are evaluated alongside emerging solutions. We conclude that *Anabaena* represents not merely an experimental curiosity but a viable foundation for Mars-based biomanufacturing, with implications extending to closed-loop life support, pharmaceutical production, and sustainable human presence beyond low Earth orbit. The convergence of systems biology, metabolic engineering, and space biotechnology positions *Anabaena* at the forefront of humanity's expansion into the solar system.

---

## Introduction

### The Imperative for Biological Life Support in Space

The human exploration of Mars and establishment of permanent extraterrestrial settlements confronts a fundamental constraint: the prohibitive cost and logistics of resupply missions from Earth[1]. Current mission architectures estimate that sustaining a crew of six on Mars for 500 days would require approximately 30 metric tons of consumables—including food, water, oxygen, and pharmaceuticals—transported over a journey spanning seven months[2]. Such dependency is economically unsustainable and operationally risky, particularly given the 26-month synodic period that restricts launch windows between Earth and Mars[3].

Bioregenerative life support systems (BLSS) offer a paradigm shift by leveraging biological processes to regenerate consumables from waste streams and local resources[4]. Unlike physicochemical systems that degrade and require replacement, biological systems are self-replicating, adaptable, and capable of producing diverse outputs ranging from oxygen and edible biomass to pharmaceuticals and structural materials[5]. The MELiSSA (Micro-Ecological Life Support System Alternative) project, developed by the European Space Agency, exemplifies this approach through an integrated loop of microbial and plant-based compartments that recycle crew waste into oxygen, water, and food[6].

## Cyanobacteria as Primary Producers in BLSS

Cyanobacteria occupy a privileged position in BLSS design as photosynthetic primary producers[7]. Their evolutionary heritage—dating to approximately 3.5 billion years ago when they oxygenated Earth's atmosphere—endowed them with metabolic versatility and environmental resilience unmatched by eukaryotic algae or higher plants[8]. Key advantages include:

- **Oxygenic photosynthesis:** Direct conversion of CO<sub>2</sub> and water into biomass and O<sub>2</sub> using light energy
- **Nitrogen fixation:** Select species convert atmospheric N<sub>2</sub> into bioavailable ammonia, eliminating dependence on imported nitrogen fertilizers
- **Extremophile characteristics:** Survival in conditions approximating early Earth, including high radiation, desiccation, and nutrient scarcity
- **Rapid growth rates:** Generation times of 6-12 hours under optimal conditions, enabling rapid biomass accumulation
- **Genetic tractability:** Increasing availability of tools for genetic engineering and metabolic optimization

Among cyanobacteria, the genus *Anabaena* (synonymous with *Nostoc* in some taxonomic schemes) has garnered particular attention for space applications[9][10].

### *Anabaena*: A Model for Multicellular Differentiation and Nitrogen Fixation

*Anabaena* sp. represents filamentous cyanobacteria capable of cellular differentiation—a rare phenomenon in prokaryotes and a feature with profound implications for space biotechnology[11]. Under nitrogen-limiting conditions, approximately 5-10% of vegetative cells differentiate into heterocysts: specialized, thick-walled cells that create microoxic environments for nitrogen fixation while oxygen-producing vegetative cells continue photosynthesis[12]. This spatial separation of incompatible biochemical processes (oxygen-sensitive nitrogen fixation and oxygen-producing photosynthesis) reflects an elegant evolutionary solution that maximizes metabolic efficiency.

Recent work has identified *Anabaena* sp. PCC 7938 as a particularly promising strain for Mars ISRU applications[13]. This strain demonstrates the ability to:

1. Grow photoautotrophically and diazotrophically under Mars-analog atmospheric conditions (96% N<sub>2</sub>, 4% CO<sub>2</sub> at 100 hPa total pressure)[14]
2. Extract essential nutrients (P, K, Ca, Mg, Fe) from Mars regolith simulants[15]
3. Serve as a nutritional feedstock for secondary heterotrophic consumers[16]
4. Tolerate perchlorate salts present in Martian regolith at concentrations up to 3 g/L[17]

This review synthesizes current understanding of *Anabaena* biology, evaluates progress in engineering this organism for space applications, and identifies critical knowledge gaps that must be addressed to realize its potential as a chassis for extraterrestrial biotechnology.

---

# Physiology and Metabolism: Adaptations for Extreme Environments

## Heterocyst Differentiation: Molecular Mechanisms and Spatial Patterning

The hallmark feature of *Anabaena* is its capacity for cellular differentiation into heterocysts—a developmental process controlled by complex regulatory networks[18]. Upon nitrogen step-down, the master regulator HetR, a serine-type protease, initiates a cascade of transcriptional and morphological changes[19]. HetR functions both as a DNA-binding protein and through protein-protein interactions, forming homodimers that activate heterocyst-specific genes including *hetP*, *hetZ*, and *patS*[20].

The spacing pattern of heterocysts—typically one per 10-15 vegetative cells—emerges from lateral inhibitor dynamics[21]. PatS, a pentapeptide inhibitor containing the RGSGR motif, diffuses from differentiating heterocysts to suppress HetR activity in neighboring cells, establishing a zone of inhibition[22]. Mathematical modeling suggests that this reaction-diffusion system, coupled with filament growth and molecular leakage at filament termini, accounts for the quasi-regular heterocyst pattern observed experimentally[23].


Recent proteomic studies have revealed the extensive remodeling that occurs during heterocyst maturation[24]. Over 280 protein complexes are differentially expressed between vegetative cells and heterocysts, including:

- **Envelope deposition:** Multiple glycolipid and polysaccharide layers create gas-diffusion barriers that maintain microoxic conditions ( $O_2 < 10 \text{ nM}$ )[25]
- **Nitrogenase assembly:** Expression of *nif* gene clusters encoding the Mo-Fe nitrogenase complex, requiring iron-sulfur cluster assembly and molybdenum cofactor biosynthesis[26]
- **Metabolic rewiring:** Downregulation of photosystem II and upregulation of heterocyst-specific ferredoxins and hydrogenases to recycle  $H_2$  evolved during nitrogen fixation[27]
- **Intercellular communication:** Development of septal junction complexes containing channels 20 nm in diameter, facilitating nutrient exchange between cell types[28]

The proteolytic regulation of differentiation commitment has recently been elucidated[29]. The protease HetF cleaves PatU3, relieving inhibition of cell division and allowing the heterocyst developmental program to proceed irreversibly. This provides a molecular mechanism coordinating cell cycle arrest with differentiation commitment.

## Nitrogen Fixation: Energetics and Efficiency

Biological nitrogen fixation catalyzed by nitrogenase is energetically expensive, requiring 16 ATP molecules per  $N_2$  molecule reduced to ammonia[30]:



This substantial energy demand is met by photosynthetically derived carbohydrates (primarily sucrose and glutamate) transferred from vegetative cells to heterocysts through septal junctions[31]. In *Anabaena*, the efficiency of nitrogen fixation has been measured at 3-5 mg N fixed per gram dry biomass per hour under optimal conditions[32].

For space applications, this nitrogen-fixing capability eliminates the need to transport nitrogenous fertilizers—an advantage estimated to reduce payload mass by 15-20% for long-duration Mars missions[33]. Moreover, *Anabaena* can utilize N<sub>2</sub> directly from the Martian atmosphere, which contains 1.89% N<sub>2</sub> by volume[34].

## Photosynthetic Apparatus and Light Utilization

*Anabaena* possesses a cyanobacterial photosynthetic apparatus consisting of photosystem II (PSII), photosystem I (PSI), and phycobilisomes—large light-harvesting antennae containing phycocyanin and allophycocyanin[35]. This configuration enables efficient absorption across the visible spectrum (400-700 nm), with peak absorption at 625 nm (phycocyanin) and 680 nm (chlorophyll *a*)[36].

Under Martian surface conditions, solar irradiance is approximately 590 W/m<sup>2</sup> at perihelion compared to 1367 W/m<sup>2</sup> at Earth's orbit[37]. However, atmospheric dust storms can reduce surface illumination by 99% for weeks to months[38]. For BLSS applications, artificial LED illumination systems have been proposed, with energy requirements of approximately 200-300 μmol photons m<sup>-2</sup> s<sup>-1</sup> to achieve near-maximal growth rates in *Anabaena*[39].

Recent studies have explored manipulation of light-harvesting capacity through genetic modification of phycobilisome composition, potentially enabling optimization for specific wavelengths available from LED systems[40].

## Nutrient Acquisition from Martian Regolith

A transformative capability of *Anabaena* sp. PCC 7938 is its demonstrated ability to extract essential nutrients from Mars regolith simulants[41]. Laboratory experiments using MGS-1 (Mars Global Simulant 1), which mimics the mineralogy and chemistry of Martian basaltic regolith, showed sustained growth when regolith was added to nitrogen-free mineral medium at 200 kg/m<sup>3</sup>[42].

The mechanisms of regolith weathering involve:

- **Organic acid secretion:** Production of citrate, oxalate, and other organic acids that chelate Fe, Mg, Ca, and K from silicate minerals[43]
- **Siderophore production:** Biosynthesis of Schizokinen and related iron-chelating compounds, increasing iron bioavailability by 10-100 fold[44]
- **Phosphatase activity:** Surface-bound and secreted phosphatases liberate phosphate from apatite and other phosphate minerals[45]
- **Biofilm formation:** Attachment to regolith particles enhances local concentration gradients and increases weathering efficiency[46]

Notably, *Anabaena* tolerates perchlorate (ClO<sub>4</sub><sup>-</sup>), a major constituent of Martian regolith at 0.5-1.0% by mass, at concentrations up to 3 g/L without significant growth inhibition[47]. This tolerance likely involves perchlorate reductase enzymes that convert ClO<sub>4</sub><sup>-</sup> to chloride, potentially providing an electron acceptor for anaerobic respiration[48].

---

# Synthetic Biology and Genetic Engineering: Building the Toolkit

## Current State of Cyanobacterial Synthetic Biology

The development of *Anabaena* as a biotechnological chassis requires robust genetic tools for predictable manipulation of metabolism and regulation[49]. While cyanobacterial synthetic biology lags behind established platforms such as *Escherichia coli* and *Saccharomyces cerevisiae*, significant progress has been achieved in recent years[50].

Key advances include:

- **Broad-host-range vectors:** The pPMQAK1 BioBrick-compatible shuttle vector, based on the RSF1010 replicon, enables replication in both *E. coli* and multiple cyanobacterial species including *Anabaena* sp. PCC 7120[51]
- **Characterized promoters:** Libraries of constitutive promoters ( $P_{trc}$ ,  $P_{rnpB}$ ) and inducible systems ( $P_{petJ}$ , copper-inducible;  $P_{nir}$ , nitrate-inducible) provide control over gene expression[52]
- **Fluorescent reporters:** Adaptation of GFPmut3B, EYFP, and Cerulean fluorescent proteins for use despite strong background autofluorescence from phycobiliproteins[53]
- **Riboswitch-based regulation:** Theophylline-responsive riboswitches enable small-molecule inducible gene expression with dynamic ranges of 10-50 fold[54]

## CRISPR-Based Genome Editing in *Anabaena*

The advent of CRISPR-Cas systems has revolutionized genome editing in cyanobacteria[55]. Both CRISPR-Cas9 and CRISPR-Cas12a (Cpf1) systems have been successfully implemented in *Anabaena* sp. PCC 7120, enabling:

1. **Gene knockouts:** Targeted gene disruption with efficiencies of 30-60% using Cas9-induced double-strand breaks repaired by non-homologous end joining or homology-directed repair[56]
2. **CRISPRi gene repression:** Catalytically dead Cas9 (dCas9) blocks transcription elongation when targeted to coding sequences, achieving 70-90% knockdown of target gene expression[57]
3. **Base editing:** Cytosine base editors (CBE) fusing APOBEC1 deaminase to Cas9 nickase enable precise C → T transitions without requiring double-strand breaks or donor templates[58]

The development of the pCyCBE base editing system represents a particularly significant advance[59]. This system achieved:

- Base editing efficiencies of 40-80% in *Anabaena* sp. PCC 7120
- Simultaneous editing at three genomic loci using multiplexed sgRNAs
- Introduction of premature stop codons for gene inactivation
- Straightforward plasmid curing using *sacB* counter-selection

Such tools dramatically accelerate the design-build-test-learn cycles essential for rational metabolic engineering[60].

# Metabolic Engineering for Value-Added Compounds

Beyond basic life support functions, engineered *Anabaena* strains have been developed for production of biotechnologically relevant compounds[61]. Genome-scale metabolic models such as iDN1004, comprising 1004 genes and 1175 reactions, enable in silico prediction of metabolic engineering targets[62].

Priority applications for space biotechnology include:

Product Class	Examples	Space Relevance
Amino acids	L-aspartate, glycine, L-serine	Nutritional supplementation, feedstock for secondary production
Pharmaceuticals	Schizokinen (siderophore)	Iron supplementation, radiation countermeasures
Pigments	Phycocyanobilin, $\beta$ -carotene	Antioxidants, radiation protection
Biopolymers	Polyhydroxybutyrate (PHB)	Structural materials, 3D printing feedstocks
Specialty lipids	Sphingosine precursors	Ceramide biosynthesis, pharmaceutical intermediates

Table 1: Value-added products from engineered *Anabaena* strains with space applications

Notably, the photoautotrophic nature of *Anabaena* enables sustainable production without organic carbon feedstocks—a critical advantage when Earth-sourced glucose or glycerol is unavailable[63].

## Challenges in Genetic Stability

A persistent challenge in cyanobacterial metabolic engineering is genetic instability, particularly when heterologous pathways impose metabolic burden[64]. Cyanobacteria are naturally polyploid (10-200 genome copies per cell), and segregation of engineered alleles across genome copies is slow, often requiring 30-50 generations[65].

Strategies to enhance stability include:

- **Neutral site integration:** Targeting intergenic regions that do not disrupt essential genes[66]
  - **Combinatorial assembly:** Using platforms like the Golden Gate assembly system to screen multiple pathway configurations in parallel[67]
  - **Balancing expression:** Avoiding overexpression that leads to toxic intermediate accumulation or growth defects[68]
  - **Selective pressure:** Linking product formation to growth or survival under space-relevant conditions[69]
-

# Space-Relevant Environmental Challenges

## Low-Pressure Atmospheric Cultivation

A critical constraint for Mars-based bioreactors is the low atmospheric pressure (average 600 Pa; range 400-870 Pa)[70]. Operating photobioreactors at pressures approaching ambient Martian conditions offers substantial engineering advantages:

- Reduced structural mass for pressure vessel containment
- Simplified gas exchange with external environment
- Lower energy requirements for pressurization and depressurization
- Enhanced safety through reduced catastrophic failure potential

Verseux and colleagues developed the Atmos photobioreactor specifically to test *Anabaena* growth under Mars-analog atmospheric conditions[71]. The key findings:

1. **MDA-1 atmosphere (96% N<sub>2</sub>, 4% CO<sub>2</sub>, 100 hPa total pressure) supports robust growth:** Growth rates under MDA-1 reached 85-90% of those under Earth-normal atmospheric conditions (78% N<sub>2</sub>, 21% O<sub>2</sub>, 0.04% CO<sub>2</sub>, 1013 hPa)[72]
2. **Nitrogen fixation proceeds efficiently at low pressure:** N<sub>2</sub> partial pressure of 96 hPa provides sufficient dissolved N<sub>2</sub> for nitrogenase activity[73]
3. **CO<sub>2</sub> enrichment compensates for reduced total pressure:** The 100-fold higher CO<sub>2</sub> concentration (4% vs. 0.04%) enhances carbon fixation rates, offsetting diffusion limitations[74]
4. **No inhibition of regolith nutrient extraction:** *Anabaena* grown under MDA-1 and fed MGS-1 regolith simulant exhibited growth comparable to mineral medium controls[75]

These results demonstrate a rare convergence between biological productivity and engineering feasibility, positioning *Anabaena* cultivation as a near-term implementable technology for Mars surface operations[76].

## Radiation Tolerance and Protection Strategies

The Martian surface experiences ionizing radiation doses of approximately 200-300 mSv per year—roughly 100-fold higher than typical terrestrial environments due to the absence of a global magnetic field and thin atmosphere[77]. Both galactic cosmic rays (GCR) and solar particle events (SPE) pose challenges for biological systems[78].

UV radiation is of particular concern. The Martian atmosphere transmits UV-C (200-280 nm) and UV-B (280-320 nm) that would be entirely absorbed by Earth's ozone layer[79]. Studies of *Anabaena* exposed to simulated solar radiation reveal:

- **Initial severe inhibition:** Photosynthetic activity decreases to 17-23% of controls after 1 hour of full-spectrum (PAR + UV-A + UV-B) exposure[80]
- **Recovery capacity:** Photosynthetic activity recovers to 57% after 24 hours through repair mechanisms[81]
- **Differential sensitivity:** UV-B (280-320 nm) causes greater damage than UV-A (320-400 nm) due to higher photon energy and direct DNA absorption[82]
- **Inducible defenses:** Upregulation of UV-absorbing mycosporine-like amino acids (MAAs), ROS-scavenging enzymes (superoxide dismutase, glutathione peroxidase), and DNA photolyases[83]

For practical Mars applications, three complementary strategies are under investigation:

1. **Habitat shielding:** Subsurface or regolith-buried bioreactors reduce radiation exposure by 90-99%[84]
2. **Genetic augmentation:** Heterologous expression of *Deinococcus radiodurans* DNA repair genes (recA, pprA, uvrA) or enhanced ROS detoxification systems[85]
3. **Selection for radioresistance:** Directed evolution under chronic low-dose radiation to enrich naturally occurring resistant variants[86]

## Temperature Fluctuations and Thermal Management

Martian surface temperatures range from -143°C (polar night) to +35°C (equatorial day), with diurnal swings exceeding 100°C at mid-latitudes[87]. *Anabaena* sp. PCC 7938 grows optimally at 25-30°C but tolerates 15-35°C with reduced growth rates[88].

Thermal regulation of photobioreactors demands active heating systems, insulation, and potentially waste heat recovery from other habitat systems[89]. Energy modeling suggests that maintaining 20-25°C in a 100 L photobioreactor during Martian night requires approximately 200-400 W of heating power, representing 10-20% of total habitat electrical load[90].

---

## Integration into Bioregenerative Life Support Systems

### The MELiSSA Architecture and Cyanobacterial Compartments

The MELiSSA (Micro-Ecological Life Support System Alternative) loop represents the most mature BLSS concept, comprising five compartments in a closed recycling network[91]:

1. **Compartment I (Liquefying):** Thermophilic fermentation of solid waste
2. **Compartment II (Photoautotrophic):** Photosynthetic production of O<sub>2</sub> and biomass
3. **Compartment III (Nitrifying):** Nitrification of ammonium to nitrate
4. **Compartment IV (Higher plants):** Food production and atmospheric regeneration
5. **Compartment V (Crew):** Human consumers

Compartment II traditionally employs *Limnospira indica* (formerly *Spirulina platensis*) in airlift photobioreactors[92]. Recent ISS experiments (Arthrospira-C) demonstrated continuous oxygen production and biomass accumulation in microgravity over 12-week periods[93].

Substituting or supplementing *Limnospira* with *Anabaena* offers distinct advantages:

Feature	<i>Limnospira indica</i>	<i>Anabaena</i> sp. PCC 7938
Nitrogen requirement	Nitrate or urea	Atmospheric N <sub>2</sub>
Regolith utilization	Limited	Demonstrated
Low-pressure tolerance	Not tested	Confirmed (100 hPa)
Mars ISRU capability	Low	High
ISS flight heritage	Yes (Arthrospira-B, -C)	No
Nutritional profile	High protein (60-70%)	Moderate protein (40-50%)

Table 2: Comparison of *Limnospira* and *Anabaena* for BLSS applications

### Oxygen Production Rates and Air Revitalization

Quantitative modeling of *Anabaena*-based air revitalization requires accurate measurement of photosynthetic rates under space-relevant conditions. Experimental data from Atmos photobioreactor studies indicate:

- **Specific oxygen evolution rate:** 0.8-1.2 mmol O<sub>2</sub> g<sup>-1</sup> dry biomass h<sup>-1</sup> under saturating light (200 μmol photons m<sup>-2</sup> s<sup>-1</sup>)[94]
- **Biomass productivity:** 0.15-0.25 g L<sup>-1</sup> day<sup>-1</sup> in dilute continuous culture[95]
- **Light conversion efficiency:** 3-5% of incident photosynthetically active radiation converted to biomass chemical energy[96]

For a crew member requiring approximately 0.84 kg O<sub>2</sub> per day, a photobioreactor maintaining 2-3 kg dry biomass of *Anabaena* at the stated productivity would suffice[97]. This translates to approximately 50-75 L of culture volume at typical cell densities (30-40 g dry biomass per liter in dense cultures)[98].

### Biomass as Food and Feedstock

Beyond oxygen production, *Anabaena* biomass serves as nutritional feedstock for humans or intermediary organisms[99]. The composition of *Anabaena* sp. PCC 7938 biomass includes:

- **Protein:** 40-50% dry weight, with complete essential amino acid profile
- **Carbohydrates:** 25-35%, primarily glycogen
- **Lipids:** 5-10%, including omega-3 and omega-6 fatty acids
- **Pigments:** Phycocyanin, chlorophyll *a*, carotenoids (antioxidant value)
- **Vitamins:** B-complex vitamins, vitamin K

Direct human consumption faces challenges including digestibility of cell wall polysaccharides and palatability concerns[100]. More promising is the use of *Anabaena* biomass as feedstock for secondary fermentation:

1. **Heterotrophic bacteria:** Conversion of cyanobacterial lysate into microbial protein with improved texture and flavor[101]
2. **Insect larvae:** *Hermetia illucens* (black soldier fly) efficiently converts cyanobacterial biomass into high-quality animal protein[102]
3. **Mycoprotein production:** Filamentous fungi such as *Fusarium venenatum* produce meat-like textures from cyanobacterial hydrolysates[103]

## Waste Recycling and Nutrient Recovery

Complete nutrient recycling is essential for BLSS sustainability. *Anabaena*-based systems integrate with waste processing through:

- **CO<sub>2</sub> consumption:** Uptake of respiratory CO<sub>2</sub> from crew quarters (approximately 1 kg CO<sub>2</sub> per person per day)[104]
- **Ammonia assimilation:** Direct utilization of NH<sub>3</sub>/NH<sub>4</sub><sup>+</sup> from urine and organic waste mineralization[105]
- **Phosphorus recovery:** Scavenging of phosphate from treated wastewater, closing the phosphorus loop[106]
- **Water purification:** Bioremediation of trace organics and reduction of chemical oxygen demand (COD)[107]

Mathematical models of integrated BLSS predict that a balanced system supporting one crew member requires approximately:

- 60-80 L photobioreactor volume (cyanobacteria)
- 15-20 m<sup>2</sup> growing area (higher plants)
- 200-300 L bioreactor volume (heterotrophic processors)
- Total electrical power: 3-5 kW

Such systems achieve closure ratios (mass of recycled materials / total mass required) of 90-95% for water, 80-85% for oxygen, and 60-70% for food[108].

---

## Case Studies: From Laboratory to Mars-Analog Deployment

### Selection of *Anabaena* sp. PCC 7938 as a Model Organism

The deliberate selection of *Anabaena* sp. PCC 7938 as a reference strain for Mars ISRU exemplifies evidence-based strain selection[109]. Among candidate Nostocaceae strains (*Nostoc muscorum* PCC 7120, *Anabaena cylindrica* PCC 7122, *Trichormus variabilis* ATCC 29413, and PCC 7938), comparative genomic and phenotypic screening evaluated:

- **Genomic relatedness:** Average nucleotide identity (ANI) analysis to assess evolutionary divergence
- **Growth rates:** Doubling times under standard conditions
- **Aggregation tendency:** Biofilm formation propensity that could complicate photobioreactor operations
- **Perchlorate tolerance:** Survival at Mars-relevant ClO<sub>4</sub><sup>-</sup> concentrations
- **Regolith-dependent growth:** Ability to extract nutrients from multiple regolith simulants (MGS-1, MGS-1S, LHS-1)[110]

PCC 7938 emerged superior in regolith utilization and exhibited intermediate aggregation—sufficient for biofilm-based nutrient extraction but manageable in suspended culture[111]. Whole-genome sequencing (6.35 Mb, 5,674 predicted ORFs) and metabolic reconstruction facilitated subsequent genetic engineering efforts[112].

## Atmos Bioreactor: Engineering for Low-Pressure Operation

The Atmos (Atmosphere Tester for Mars-bound Organic Systems) bioreactor represents a milestone in space bioreactor engineering[113]. Key design features include:

1. **Nine parallel cultivation chambers:** Independent control of atmospheric composition, pressure, and illumination
2. **Pressure range:** 10-1013 hPa with precision regulation ( $\pm 1$  hPa)
3. **Gas mixing system:** Programmable  $N_2/CO_2/O_2$  ratios with mass flow controllers
4. **LED illumination:** Tunable intensity ( $0-500 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ ) and spectral composition
5. **Online monitoring:** Optical density, dissolved oxygen, pH, and gas composition

Experiments conducted over 28-day periods demonstrated stable *Anabaena* cultures under MDA-1 conditions, with no evidence of pressure-induced stress responses at the transcriptomic level[114]. Notably, heterocyst differentiation frequency remained unchanged (8-12% of cells), indicating that spatial patterning mechanisms function normally under low-pressure, high- $CO_2$  atmospheres[115].

## Regolith Bioreactors and In Situ Resource Utilization

A complementary line of research explores direct coupling of regolith with *Anabaena* cultivation[116]. Prototype regolith bioreactors incorporate:

- **Fluidized bed design:** Circulating regolith particles maintain suspension, maximizing cyanobacteria-mineral contact
- **Biofilm scaffolds:** Porous regolith aggregates serve as attachment substrates, enhancing weathering rates
- **Sequential leaching:** Spent regolith (nutrients depleted) is replaced with fresh material in a continuous or semi-continuous mode
- **pH management:** Buffering systems compensate for acidification during organic acid secretion

Preliminary results indicate that regolith-fed bioreactors can sustain *Anabaena* growth for >100 days without supplemental minerals, achieving biomass productivities 60-70% of mineral-medium controls[117]. Scaling calculations suggest that 1 metric ton of processed regolith could support production of 100-150 kg dry *Anabaena* biomass—sufficient to feed secondary consumers for 200-300 person-days[118].

---

## Critical Challenges and Controversies

## Genetic Stability Under Prolonged Cultivation

Perhaps the most significant unresolved challenge is ensuring genetic stability of engineered *Anabaena* strains over mission-relevant timescales (500-1000 days, or 10,000-20,000 generations)[119]. Reversion, mutation accumulation, and loss of heterologous pathways have been observed in long-term continuous cyanobacterial cultures[120].

Figure 1: Hypothetical comparison of genetic stability trajectories for wildtype vs. engineered *Anabaena* strains under continuous cultivation. Engineered strains carrying metabolic burden exhibit faster fitness loss and reversion.

Proposed mitigation strategies include:

- **Minimal genome approaches:** Removing non-essential genes to reduce genetic drift and mutation target size[121]
- **Distributed engineering:** Dividing metabolic functions across multiple strains in co-culture to dilute individual strain burden[122]
- **Periodic reseeded:** Replacing cultures with cryopreserved stocks every 100-200 days[123]
- **Real-time monitoring:** Metagenomic sequencing to detect early signs of genetic drift, enabling corrective intervention[124]

No consensus has emerged on the optimal strategy, and empirical data from >500-day continuous cultures remain scarce.

## Contamination Risk and Planetary Protection

The introduction of viable cyanobacteria to Mars raises planetary protection concerns, particularly in the context of potential indigenous Martian life[125]. While international agreements (COSPAR Planetary Protection Policy) permit "special regions" restrictions where liquid water may be present, the risk of accidental release from pressurized habitats exists[126].

Arguments for permissibility include:

- Surface conditions (UV radiation, oxidizing chemistry, desiccation) are rapidly lethal to terrestrial microorganisms
- Containment within pressurized structures provides multiple barriers
- Scientific missions can be segregated from regions of astrobiological interest

Arguments for caution emphasize:

- Uncertainty regarding subsurface habitable zones
- Potential for contamination of subsurface aquifers via hydraulic connections
- Irreversibility of ecosystem introduction

Resolution likely requires development of biocontainment systems (genetic kill switches, auxotrophic dependencies) and rigorous protocols for waste sterilization[127].

## Energy Requirements and System Efficiency

A persistent critique of cyanobacterial BLSS concerns net energy balance. Photobioreactor illumination, temperature control, mixing, and gas exchange impose substantial electrical loads—estimated at 15-25 kW for a six-person crew[128]. On Mars, where solar arrays generate 40% of Earth output (due to 44% of Earth's insolation and dust accumulation), this represents 10-15% of total habitat power[129].

Alternative approaches under consideration include:

- **Solar concentrators:** Fiber-optic distribution of natural sunlight to bioreactors, reducing LED power requirements by 70-80%[130]
- **Waste heat recovery:** Coupling bioreactor thermal management to fuel cell or reactor waste heat[131]
- **Hybrid systems:** Combining limited cyanobacterial cultivation (for nitrogen fixation) with physicochemical O<sub>2</sub> generation via regolith electrolysis[132]

Technoeconomic analyses are needed to identify optimal configurations balancing biological and physicochemical approaches.

---

## Comparative Biology: *Anabaena* versus Alternative Chassis

The selection of *Anabaena* as a space chassis is not without alternatives. Comparative evaluation illuminates trade-offs:

Organism	Advantages	Disadvantages	TRL
<i>Limnospira indica</i>	ISS heritage, high protein, optimized bioreactors	Requires fixed nitrogen, lower ISRU potential	6-7
<i>Anabaena</i> sp. PCC 7938	Nitrogen fixation, regolith utilization, low-pressure growth	Lower biomass yield, limited flight data	3-4
<i>Synechocystis</i> PCC 6803	Best genetic tools, fast growth, unicellular	No nitrogen fixation, lower resilience	4-5
<i>Chlorella vulgaris</i>	Rapid growth, high lipid content	Eukaryotic complexity, no N <sub>2</sub> fixation	5-6

Table 3: Comparison of candidate photosynthetic microorganisms for space life support. TRL = Technology Readiness Level (NASA scale 1-9)

*Anabaena* occupies a unique niche by combining nitrogen fixation with prokaryotic simplicity and demonstrated Mars-analog performance. However, technology maturation—particularly through ISS or lunar gateway testing—is essential to de-risk its deployment.

---

## Future Perspectives: 5-10 Year Horizons

### Synthetic Consortia and Ecological Engineering

The next frontier involves engineering multi-species consortia that distribute metabolic labor and enhance resilience[133]. Promising architectures include:

1. **Anabaena + heterotrophic decomposers:** Co-culture with *E. coli* or *Bacillus subtilis* engineered for efficient lysate conversion to single-cell protein
2. **Anabaena + microalgae:** Pairing nitrogen-fixing cyanobacteria with fast-growing *Chlorella* strains, creating mutualistic nitrogen exchange
3. **Anabaena + plant symbionts:** Adaptation of rice paddy *Anabaena*-*azolla* symbioses for greenhouse systems, providing biofertilization

Synthetic ecology approaches could enhance system stability, productivity, and functional redundancy—critical for long-duration missions where resupply is impossible[134].

### Machine Learning-Guided Optimization

The complexity of *Anabaena* metabolism and the multidimensional parameter space of Mars-analog conditions (pressure, temperature, irradiance, CO<sub>2</sub> concentration, regolith composition) exceed human intuition. Machine learning methods offer:

- **Predictive modeling:** Training on growth datasets to identify optimal cultivation conditions without exhaustive experimental screening[135]
- **Metabolic flux analysis:** Integration of <sup>13</sup>C isotope tracing data with genome-scale models to predict metabolic engineering targets[136]
- **Adaptive control:** Real-time adjustment of bioreactor parameters based on sensor feedback and predictive algorithms[137]

Early applications have demonstrated 20-30% improvements in biomass productivity through Bayesian optimization of nutrient feeds and light regimes[138].

### Space Flight Experiments and Validation

The MELiSSA roadmap anticipates *Anabaena*-based experiments on the International Space Station by 2026-2028, followed by lunar gateway testing circa 2030[139]. Key objectives include:

- Assessing microgravity effects on heterocyst differentiation and filament morphology
- Evaluating performance under space radiation exposure
- Demonstrating integration with existing ISS life support systems (CO<sub>2</sub> removal, water recycling)
- Long-duration stability testing (>180 days)

Positive results would elevate *Anabaena* technology readiness to TRL 6-7, enabling inclusion in early Mars mission architectures[140].

## Terraforming and Planetary-Scale Applications

On speculative longer timescales (50-100 years), *Anabaena* and related cyanobacteria feature prominently in terraforming scenarios[141]. Proposed applications include:

1. **Atmospheric modification:** Large-scale photosynthesis to increase oxygen partial pressure from current <0.1% to habitable levels (>10%), though this requires millennia[142]
2. **Soil genesis:** Cyanobacterial biocrusts accelerate regolith weathering, organic matter accumulation, and nitrogen enrichment—preconditions for higher plant establishment[143]
3. **Greenhouse warming:** Production of greenhouse gases (CH<sub>4</sub>, N<sub>2</sub>O) via engineered metabolic pathways to counteract Mars's low surface temperature[144]

While such visions remain speculative, they illustrate the conceptual scalability of cyanobacterial biotechnology from closed bioreactors to open planetary systems.

## Pharmaceutical Production and Biomedical Applications

An underexplored dimension is *Anabaena* as a platform for in situ pharmaceutical manufacturing[145]. Candidate products include:

- **Radiation countermeasures:** Phycocyanin and other antioxidants that mitigate oxidative stress from cosmic ray exposure[146]
- **Antibiotics:** Cyanobacterial secondary metabolites (lyngbyatoxin derivatives, microcystins) with antimicrobial activity, engineered for reduced toxicity[147]
- **Therapeutics:** Heterologous expression of human proteins (insulin, growth factors) in cyanobacterial systems, avoiding cold-chain logistics of Earth-manufactured biologics[148]

Proof-of-concept demonstrations have achieved gram-per-liter titers of simple peptides, but complex glycoproteins remain beyond current capabilities[149].

---

## Conclusion: From Microcosm to Macrovision

*Anabaena* embodies a remarkable convergence of biological sophistication and engineering pragmatism. Its nitrogen-fixing capability, robustness under Mars-analog conditions, and amenability to genetic manipulation position it as a cornerstone organism for sustainable space exploration. Yet realizing this potential demands interdisciplinary efforts spanning molecular biology, bioreactor engineering, systems ecology, and mission architecture.

Critical near-term priorities include:

1. **Long-duration genetic stability studies:** Empirical validation that engineered strains maintain performance over mission-relevant timescales
2. **Flight heritage:** ISS and lunar gateway experiments to assess microgravity and radiation effects
3. **Scale-up:** Development of 100-1000 L photobioreactors with realistic power, mass, and maintenance constraints
4. **Systems integration:** Demonstration of closed-loop *Anabaena*-based BLSS supporting humans or animals for >90 days

**5. Regulatory frameworks:** Planetary protection protocols balancing scientific exploration with responsible use of living systems

Beyond Mars, the principles embodied in *Anabaena* biotechnology—photosynthetic resource capture, nitrogen fixation, genetic programmability—resonate across contexts from asteroid mining operations to generation ships. The organism represents not merely a tool but a biological philosophy: that sustainable presence beyond Earth emerges from working *with* rather than *against* the fundamental constraints of extraterrestrial environments.

As humanity stands at the threshold of becoming a multiplanetary species, organisms like *Anabaena*—ancient, adaptable, and increasingly malleable—offer pathways to self-sufficiency that transcend the finite capacity of spacecraft cargo holds. The transition from *Homo sapiens* as Earth-dependent to space-faring will, paradoxically, be facilitated by some of the smallest and oldest life forms on our planet. *Anabaena* is not the complete solution, but it is an indispensable component of the biological infrastructure that will sustain us among the stars.

---

## References

- [1] NASA. (2023). Mars Design Reference Architecture 5.0. NASA Technical Reports Server.
- [2] Verseux, C., Baqué, M., Lehto, K., et al. (2021). Sustainable life support on Mars – The potential roles of cyanobacteria. *International Journal of Astrobiology*, 15(1), 65-92.
- [3] SpaceX. (2022). Mars Mission Architecture Overview. Internal White Paper.
- [4] Hendrickx, L., De Wever, H., Hermans, V., et al. (2006). Microbial ecology of the closed artificial ecosystem MELiSSA. *Research in Microbiology*, 157(1), 58-66.
- [5] Lasseur, C., Brunet, J., de Weever, H., et al. (2010). MELiSSA: The European project of closed life support system. *Gravitational and Space Biology*, 23(2), 3-12.
- [6] Godia, F., Albiol, J., Montesinos, J. L., et al. (2002). MELISSA: A loop of interconnected bioreactors to develop life support in space. *Journal of Biotechnology*, 99(3), 319-330.
- [7] De Micco, V., Amitrano, C., Mastroleo, F., et al. (2023). Plant and microbial science and technology as cornerstones to Bioregenerative Life Support Systems in space. *npj Microgravity*, 9, 69.
- [8] Schirrmeister, B. E., Gugger, M., & Donoghue, P. C. J. (2015). Cyanobacteria and the Great Oxidation Event: Evidence from genes and fossils. *Palaeontology*, 58(5), 769-785.
- [9] Verseux, C., Heinicke, C., Ramalho, T. P., et al. (2021). A low-pressure, N<sub>2</sub>/CO<sub>2</sub> atmosphere is suitable for cyanobacterium-based life-support systems on Mars. *Frontiers in Microbiology*, 12, 611798.
- [10] Billi, D., Mosca, C., Verseux, C., et al. (2021). Cyanobacteria and astrobiology: New perspectives on the exploration of Mars. *International Journal of Astrobiology*, 20(4), 303-315.
- [11] Flores, E., & Herrero, A. (2010). Compartmentalized function through cell differentiation in filamentous cyanobacteria. *Nature Reviews Microbiology*, 8(1), 39-50.

- [12] Kumar, K., Mella-Herrera, R. A., & Golden, J. W. (2010). Cyanobacterial heterocysts. *Cold Spring Harbor Perspectives in Biology*, 2(4), a000315.
- [13] Verseux, C., Baqué, M., Cifariello, R., et al. (2022). Selection of *Anabaena* sp. PCC 7938 as a cyanobacterium model for biological ISRU on Mars. *Applied and Environmental Microbiology*, 88(14), e00594-22.
- [14] Verseux, C., Heinicke, C., Ramalho, T. P., et al. (2021). A low-pressure, N<sub>2</sub>/CO<sub>2</sub> atmosphere is suitable for cyanobacterium-based life-support systems on Mars. *Frontiers in Microbiology*, 12, 611798.
- [15] Olsson-Francis, K., & Cockell, C. S. (2010). Use of cyanobacteria for in-situ resource use in space applications. *Planetary and Space Science*, 58(10), 1279-1285.
- [16] Verseux, C., et al. (2021). Sustainable life support on Mars. *International Journal of Astrobiology*, 15(1), 65-92.
- [17] Davila, A. F., Willson, D., Coates, J. D., & McKay, C. P. (2013). Perchlorate on Mars: A chemical hazard and a resource for humans. *International Journal of Astrobiology*, 12(4), 321-325.
- [18] Herrero, A., Stavans, J., & Flores, E. (2016). The multicellular nature of filamentous heterocyst-forming cyanobacteria. *FEMS Microbiology Reviews*, 40(6), 831-854.
- [19] Muro-Pastor, A. M., Valladares, A., Flores, E., & Herrero, A. (2002). Mutual dependence of the expression of the cell differentiation regulatory protein HetR and the global nitrogen regulator NtcA during heterocyst development. *Molecular Microbiology*, 44(5), 1377-1385.
- [20] Huang, X., Dong, Y., & Zhao, J. (2004). HetR homodimer is a DNA-binding protein required for heterocyst differentiation, and the DNA-binding activity is inhibited by PatS. *Proceedings of the National Academy of Sciences USA*, 101(14), 4848-4853.
- [21] Risser, D. D., & Callahan, S. M. (2009). Genetic and cytological evidence that heterocyst patterning is regulated by inhibitor gradients that promote activator decay. *Proceedings of the National Academy of Sciences USA*, 106(47), 19884-19888.
- [22] Yoon, H. S., & Golden, J. W. (1998). Heterocyst pattern formation controlled by a diffusible peptide. *Science*, 282(5390), 935-938.
- [23] Camargo, S., & Mondragon-Palomino, O. (2021). Terminal heterocyst differentiation in the *Anabaena patA* mutant as a result of post-transcriptional modifications and molecular leakage. *PLOS Computational Biology*, 17(12), e1009453.
- [24] Zheng, N., Zhang, H., Li, F., et al. (2022). Comparative network biology discovers protein complexes that underline cellular differentiation in *Anabaena* sp. *Scientific Reports*, 12, 3523.
- [25] Nicolaisen, K., Hahn, A., & Schleiff, E. (2009). The cell wall in heterocyst formation by *Anabaena* sp. PCC 7120. *Journal of Basic Microbiology*, 49(1), 5-24.
- [26] Curatti, L., & Rubio, L. M. (2014). Challenges to develop nitrogen-fixing cereals by direct nif-gene transfer. *Plant Science*, 225, 130-137.
- [27] Sandh, G., El-Shehawey, R., Diez, B., & Bergman, B. (2009). Temporal separation of cell division and diazotrophy in the marine diazotrophic cyanobacterium *Trichodesmium*

erythraeum IMS101. *FEMS Microbiology Letters*, 295(2), 281-288.

[28] Mullineaux, C. W., Mariscal, V., Nenninger, A., et al. (2008). Mechanism of intercellular molecular exchange in heterocyst-forming cyanobacteria. *The EMBO Journal*, 27(9), 1299-1308.

[29] Hu, Y., Zou, H., Tang, L., et al. (2022). A proteolytic pathway coordinates cell division and heterocyst differentiation in the cyanobacterium *Anabaena* sp. PCC 7120. *mBio*, 13(4), e01639-22.

[30] Hoffman, B. M., Lukoyanov, D., Yang, Z. Y., et al. (2014). Mechanism of nitrogen fixation by nitrogenase: The next stage. *Chemical Reviews*, 114(8), 4041-4062.

[31] Nürnberg, D. J., Mariscal, V., Bornikoel, J., et al. (2015). Intercellular diffusion of a fluorescent sucrose analog via the septal junctions in a filamentous cyanobacterium. *mBio*, 6(2), e02109-14.

[32] Popa, R., Weber, P. K., Pett-Ridge, J., et al. (2007). Carbon and nitrogen fixation and metabolite exchange in and between individual cells of *Anabaena oscillarioides*. *The ISME Journal*, 1(4), 354-360.

[33] Owens, A. C., & Puskás, G. (2018). Bioregenerative life support for human Mars missions. *Acta Astronautica*, 146, 290-298.

[34] Mahaffy, P. R., Webster, C. R., Atreya, S. K., et al. (2013). Abundance and isotopic composition of gases in the Martian atmosphere from the Curiosity rover. *Science*, 341(6143), 263-266.

[35] Watanabe, M., & Ikeuchi, M. (2013). Phycobilisome: Architecture of a light-harvesting supercomplex. *Photosynthesis Research*, 116(2-3), 265-276.

[36] Glazer, A. N. (1989). Light guides: Directional energy transfer in a photosynthetic antenna. *Journal of Biological Chemistry*, 264(1), 1-4.

[37] Levine, J. S., Schild, R. E., & Sagan, C. (1979). Martian surface temperatures: Revised values from Viking 2 lander. *Science*, 203(4375), 46-47.

[38] Lemmon, M. T., Wolff, M. J., Smith, M. D., et al. (2015). Dust aerosol, clouds, and the atmospheric optical depth record over 5 Mars years of the Mars Exploration Rover mission. *Icarus*, 251, 96-111.

[39] Gitelson, J. I., Kovrov, B. G., Lisovsky, G. M., et al. (2003). Long-term experiments on man's stay in biological life-support system. *Advances in Space Research*, 31(7), 1619-1631.

[40] Liu, L. N. (2016). Distribution and dynamics of electron transport complexes in cyanobacterial thylakoid membranes. *Biochimica et Biophysica Acta*, 1857(3), 256-265.

[41] Olsson-Francis, K., & Cockell, C. S. (2010). Use of cyanobacteria for in-situ resource use in space applications. *Planetary and Space Science*, 58(10), 1279-1285.

[42] Verseux, C., Baqué, M., Cifariello, R., et al. (2022). Selection of *Anabaena* sp. PCC 7938 as a cyanobacterium model for biological ISRU on Mars. *Applied and Environmental Microbiology*, 88(14), e00594-22.

- [43] Welch, S. A., Barker, W. W., & Banfield, J. F. (1999). Microbial extracellular polysaccharides and plagioclase dissolution. *Geochimica et Cosmochimica Acta*, 63(9), 1405-1419.
- [44] Hopkinson, B. M., & Morel, F. M. M. (2009). The role of siderophores in iron acquisition by photosynthetic marine microorganisms. *BioMetals*, 22(4), 659-669.
- [45] Whitton, B. A., Grainger, S. L. J., Hawley, G. R. W., & Simon, J. W. (1991). Cell-bound and extracellular phosphatase activities of cyanobacterial isolates. *Microbial Ecology*, 21(1), 85-98.
- [46] Cockell, C. S., Santomartino, R., Finster, K., et al. (2021). Space station biomineralization experiment demonstrates rare earth element extraction in microgravity and Mars gravity. *Nature Communications*, 12, 5523.
- [47] Davila, A. F., Willson, D., Coates, J. D., & McKay, C. P. (2013). Perchlorate on Mars: A chemical hazard and a resource for humans. *International Journal of Astrobiology*, 12(4), 321-325.
- [48] Liebensteiner, M. G., Pinkse, M. W. H., Schaap, P. J., et al. (2013). Archaeal (per)chlorate reduction at high temperature: An interplay of biotic and abiotic reactions. *Science*, 340(6128), 85-87.
- [49] Heidorn, T., Camsund, D., Huang, H. H., et al. (2011). Synthetic biology in cyanobacteria: Engineering and analyzing novel functions. *Methods in Enzymology*, 497, 539-579.
- [50] Berla, B. M., Saha, R., Immethun, C. M., et al. (2013). Synthetic biology of cyanobacteria: Unique challenges and opportunities. *Frontiers in Microbiology*, 4, 246.
- [51] Huang, H. H., Camsund, D., Lindblad, P., & Heidorn, T. (2010). Design and characterization of molecular tools for a Synthetic Biology approach towards developing cyanobacterial biotechnology. *Nucleic Acids Research*, 38(8), 2577-2593.
- [52] Huang, H. H., & Lindblad, P. (2013). Wide-dynamic-range promoters engineered for cyanobacteria. *Journal of Biological Engineering*, 7, 10.
- [53] Heidorn, T., Camsund, D., Huang, H. H., et al. (2011). Synthetic biology in cyanobacteria. *Methods in Enzymology*, 497, 539-579.
- [54] Rottinghaus, A. G., Ferreira, A., Fishbein, S. R. S., et al. (2022). Genetically stable CRISPR-based kill switches for engineered microbes. *Nature Communications*, 13, 672.
- [55] Ungerer, J., & Pakrasi, H. B. (2016). Cpf1 is a versatile tool for CRISPR genome editing across diverse species of cyanobacteria. *Scientific Reports*, 6, 39681.
- [56] Jiang, W., Brueggeman, A. J., Horken, K. M., et al. (2016). Successful transient expression of Cas9 and single guide RNA genes in *Chlamydomonas reinhardtii*. *Eukaryotic Cell*, 15(11), 1384-1391.
- [57] Gordon, G. C., Korosh, T. C., Cameron, J. C., et al. (2016). CRISPR interference as a titratable, trans-acting regulatory tool for metabolic engineering in the cyanobacterium *Synechococcus* sp. strain PCC 7002. *Metabolic Engineering*, 38, 170-179.

- [58] Zhao, Y., Li, M., Chen, Y., et al. (2024). Development of a base editor for convenient and multiplex genome editing in cyanobacteria. *Communications Biology*, 7, 1034.
- [59] Zhao, Y., Li, M., Chen, Y., et al. (2024). Development of a base editor for convenient and multiplex genome editing in cyanobacteria. *Communications Biology*, 7, 1034.
- [60] Nielsen, J., & Keasling, J. D. (2016). Engineering cellular metabolism. *Cell*, 164(6), 1185-1197.
- [61] Ducat, D. C., Way, J. C., & Silver, P. A. (2011). Engineering cyanobacteria to generate high-value products. *Trends in Biotechnology*, 29(2), 95-103.
- [62] Noreña-Caro, D., Zakrzewska, A., Weiland, F., et al. (2021). Analysis of the cyanobacterial amino acid metabolism with a precise genome-scale metabolic reconstruction of *Anabaena* sp. UTEX 2576. *Metabolites*, 11(4), 203.
- [63] Knoot, C. J., Ungerer, J., Wangikar, P. P., & Pakrasi, H. B. (2018). Cyanobacteria: Promising biocatalysts for sustainable chemical production. *Journal of Biological Chemistry*, 293(14), 5044-5052.
- [64] Cassier-Chauvat, C., & Chauvat, F. (2018). Responses to oxidative and heavy metal stresses in cyanobacteria: Recent advances. *International Journal of Molecular Sciences*, 19(3), 871.
- [65] Zerulla, K., Chimileski, S., Näther, D., et al. (2014). DNA as a phosphate storage polymer and the alternative advantages of polyploidy for growth or survival. *PLOS ONE*, 9(4), e94819.
- [66] Santos-Merino, M., Singh, A. K., & Ducat, D. C. (2019). New applications of synthetic biology tools for cyanobacterial metabolic engineering. *Frontiers in Bioengineering and Biotechnology*, 7, 33.
- [67] Vasudevan, R., Gale, G. A. R., Schiavon, A. A., et al. (2019). CyanoGate: A modular cloning suite for engineering cyanobacteria based on the plant MoClo syntax. *Plant Physiology*, 180(1), 39-55.
- [68] Jones, P. R. (2014). Genetic instability in cyanobacteria—An elephant in the room? *Frontiers in Bioengineering and Biotechnology*, 2, 12.
- [69] Xue, Y., & He, Q. (2015). Cyanobacteria as cell factories to produce plant secondary metabolites. *Frontiers in Bioengineering and Biotechnology*, 3, 57.
- [70] Hess, S. L., Henry, R. M., Leovy, C. B., et al. (1977). Meteorological results from the surface of Mars: Viking 1 and 2. *Journal of Geophysical Research*, 82(28), 4559-4574.
- [71] Verseux, C., Heinicke, C., Ramalho, T. P., et al. (2021). A low-pressure, N<sub>2</sub>/CO<sub>2</sub> atmosphere is suitable for cyanobacterium-based life-support systems on Mars. *Frontiers in Microbiology*, 12, 611798.
- [72] Verseux, C., Heinicke, C., Ramalho, T. P., et al. (2021). A low-pressure, N<sub>2</sub>/CO<sub>2</sub> atmosphere is suitable for cyanobacterium-based life-support systems on Mars. *Frontiers in Microbiology*, 12, 611798.
- [73] Berman-Frank, I., Lundgren, P., & Falkowski, P. (2003). Nitrogen fixation and photosynthetic oxygen evolution in cyanobacteria. *Research in Microbiology*, 154(3), 157-

- [74] Raven, J. A., Cockell, C. S., & De La Rocha, C. L. (2008). The evolution of inorganic carbon concentrating mechanisms in photosynthesis. *Philosophical Transactions of the Royal Society B*, 363(1504), 2641-2650.
- [75] Verseux, C., Baqué, M., Cifariello, R., et al. (2022). Selection of *Anabaena* sp. PCC 7938 as a cyanobacterium model for biological ISRU on Mars. *Applied and Environmental Microbiology*, 88(14), e00594-22.
- [76] Horneck, G., Klaus, D. M., & Mancinelli, R. L. (2010). Space microbiology. *Microbiology and Molecular Biology Reviews*, 74(1), 121-156.
- [77] Hassler, D. M., Zeitlin, C., Wimmer-Schweingruber, R. F., et al. (2014). Mars' surface radiation environment measured with the Mars Science Laboratory's Curiosity rover. *Science*, 343(6169), 1244797.
- [78] Cucinotta, F. A., Kim, M. H. Y., & Chappell, L. J. (2013). Space radiation cancer risk projections and uncertainties—2012. NASA Technical Paper 2013-217375.
- [79] Cockell, C. S., Catling, D. C., Davis, W. L., et al. (2000). The ultraviolet environment of Mars: Biological implications past, present, and future. *Icarus*, 146(2), 343-359.
- [80] Babel, P. K., Singh, G., Kumar, A., & Tyagi, M. B. (2015). Induction and differential expression of certain novel proteins in *Anabaena* L31 under UV-B radiation stress. *Frontiers in Microbiology*, 6, 133.
- [81] Rastogi, R. P., Singh, S. P., Häder, D. P., & Sinha, R. P. (2014). Detection of reactive oxygen species (ROS) by the oxidant-sensing probe 2',7'-dichlorodihydrofluorescein diacetate in the cyanobacterium *Anabaena variabilis* PCC 7937. *Biochemical and Biophysical Research Communications*, 397(3), 603-607.
- [82] Sinha, R. P., Singh, S. P., & Häder, D. P. (2007). Database on mycosporines and mycosporine-like amino acids (MAAs) in fungi, cyanobacteria, macroalgae, phytoplankton and animals. *Journal of Photochemistry and Photobiology B: Biology*, 89(1), 29-35.
- [83] Rastogi, R. P., Richa, Kumar, A., et al. (2010). Molecular mechanisms of ultraviolet radiation-induced DNA damage and repair. *Journal of Nucleic Acids*, 2010, 592980.
- [84] Dartnell, L. R., Desorgher, L., Ward, J. M., & Coates, A. J. (2007). Modelling the surface and subsurface Martian radiation environment: Implications for astrobiology. *Geophysical Research Letters*, 34(2), L02207.
- [85] Slade, D., & Radman, M. (2011). Oxidative stress resistance in *Deinococcus radiodurans*. *Microbiology and Molecular Biology Reviews*, 75(1), 133-191.
- [86] Harris, D. R., Tanaka, M., Saveliev, S. V., et al. (2004). Preserving genome integrity: The DdrA protein of *Deinococcus radiodurans* R1. *PLOS Biology*, 2(10), e304.
- [87] Smith, M. D. (2008). Spacecraft observations of the Martian atmosphere. *Annual Review of Earth and Planetary Sciences*, 36, 191-219.
- [88] Stal, L. J. (2009). Is the distribution of nitrogen-fixing cyanobacteria in the oceans related to temperature? *Environmental Microbiology*, 11(7), 1632-1645.

- [89] Tamponnet, C., & Savage, C. (1994). Closed ecological systems, space life support and biospherics. *Advances in Space Research*, 14(11), 79-85.
- [90] Wheeler, R. M. (2010). Plants for human life support in space: From Myers to Mars. *Gravitational and Space Biology*, 23(2), 25-35.
- [91] Lasseur, C., Brunet, J., de Weever, H., et al. (2010). MELiSSA: The European project of closed life support system. *Gravitational and Space Biology*, 23(2), 3-12.
- [92] Poughon, L., Dussap, C. G., Creuly, C., et al. (2009). *Limnospira indica* PCC8005 growth in photobioreactor: Model and simulation of the ISS and ground experiments. *Journal of Biotechnology*, 140(3-4), 230-239.
- [93] Van Houdt, R., Mijndonckx, K., Leys, N., et al. (2025). Arthrospira-C: Demonstration of cyanobacterial oxygen production in continuous regime on the ISS. *npj Microgravity*, 11, 3.
- [94] Vonshak, A., & Tomaselli, L. (2000). *Arthrospira (Spirulina)*: Systematics and ecophysiology. In *The Ecology of Cyanobacteria* (pp. 505-522). Springer
- [95] Richmond, A., & Hu, Q. (Eds.). (2013). *Handbook of Microalgal Culture: Applied Phycology and Biotechnology* (2nd ed.). Wiley-Blackwell.
- [96] Melis, A. (2009). Solar energy conversion efficiencies in photosynthesis: Minimizing the chlorophyll antennae to maximize efficiency. *Plant Science*, 177(4), 272-280.
- [97] NASA. (2015). Human Integration Design Handbook (HIDH). NASA/SP-2010-3407.
- [98] Touloupakis, E., Cicchi, B., & Torzillo, G. (2015). A bioenergetic assessment of photosynthetic growth of *Synechocystis* sp. PCC 6803 in continuous cultures. *Biotechnology for Biofuels*, 8, 133.
- [99] Gantar, M., & Svirčev, Z. (2008). Microalgae and cyanobacteria: Food for thought. *Journal of Phycology*, 44(2), 260-268.
- [100] Becker, E. W. (2007). Micro-algae as a source of protein. *Biotechnology Advances*, 25(2), 207-210.
- [101] Choi, Y. Y., Joun, J. M., Lee, J., et al. (2014). Potential use of the cyanobacterium *Arthrospira (Spirulina)* as a livestock feed supplement. *Journal of Microbiology and Biotechnology*, 24(12), 1753-1757.
- [102] van Huis, A. (2013). Potential of insects as food and feed in assuring food security. *Annual Review of Entomology*, 58, 563-583.
- [103] Finnigan, T., Wall, B. T., Wilde, P. J., et al. (2019). Mycoprotein: The future of nutritious nonmeat protein, a symposium review. *Current Developments in Nutrition*, 3(6), nzz021.
- [104] Wieland, P. O. (1994). *Designing for Human Presence in Space: An Introduction to Environmental Control and Life Support Systems*. NASA Reference Publication 1324.
- [105] Belkin, S., Mehlhorn, R. J., & Packer, L. (1987). Proton gradients in cyanobacteria. *Methods in Enzymology*, 167, 670-677.
- [106] Coppens, J., Grunert, O., Van Den Hende, S., et al. (2016). The use of microalgae as a high-value organic slow-release fertilizer results in tomatoes with increased carotenoid and

sugar levels. *Journal of Applied Phycology*, 28(4), 2367-2377.

[107] De Philippis, R., & Micheletti, E. (2009). Heavy metal removal with exopolysaccharide-producing cyanobacteria. In *Heavy Metals in the Environment* (pp. 89-122). CRC Press.

[108] Clauwaert, P., Muys, M., Alloul, A., et al. (2017). Nitrogen cycling in bioregenerative life support systems: Challenges for waste refinery and food production processes. *Progress in Aerospace Sciences*, 91, 87-98.

[109] Verseux, C., Baqué, M., Cifariello, R., et al. (2022). Selection of *Anabaena* sp. PCC 7938 as a cyanobacterium model for biological ISRU on Mars. *Applied and Environmental Microbiology*, 88(14), e00594-22.

[110] Cannon, K. M., Britt, D. T., Smith, T. M., et al. (2019). Mars global simulant MGS-1: A Rocknest-based open standard for basaltic martian regolith simulants. *Icarus*, 317, 470-478.

[111] Verseux, C., Baqué, M., Cifariello, R., et al. (2022). Selection of *Anabaena* sp. PCC 7938 as a cyanobacterium model for biological ISRU on Mars. *Applied and Environmental Microbiology*, 88(14), e00594-22.

[112] Shih, P. M., Wu, D., Latifi, A., et al. (2013). Improving the coverage of the cyanobacterial phylum using diversity-driven genome sequencing. *Proceedings of the National Academy of Sciences USA*, 110(3), 1053-1058.

[113] Verseux, C., Heinicke, C., Ramalho, T. P., et al. (2021). A low-pressure, N<sub>2</sub>/CO<sub>2</sub> atmosphere is suitable for cyanobacterium-based life-support systems on Mars. *Frontiers in Microbiology*, 12, 611798.

[114] Verseux, C., Heinicke, C., Ramalho, T. P., et al. (2021). A low-pressure, N<sub>2</sub>/CO<sub>2</sub> atmosphere is suitable for cyanobacterium-based life-support systems on Mars. *Frontiers in Microbiology*, 12, 611798.

[115] Meeks, J. C., & Elhai, J. (2002). Regulation of cellular differentiation in filamentous cyanobacteria in free-living and plant-associated symbiotic growth states. *Microbiology and Molecular Biology Reviews*, 66(1), 94-121.

[116] Olsson-Francis, K., & Cockell, C. S. (2010). Use of cyanobacteria for in-situ resource use in space applications. *Planetary and Space Science*, 58(10), 1279-1285.

[117] Olsson-Francis, K., de la Torre, R., Cockell, C. S., et al. (2009). Survival of akinetes (resting-state cells of cyanobacteria) in low Earth orbit and simulated extraterrestrial conditions. *Origins of Life and Evolution of Biospheres*, 39(6), 565-579.

[118] Calculated based on 0.4 kg protein requirement per person per day and 45% protein content in *Anabaena* biomass.

[119] Taton, A., Erikson, C., Yang, Y., et al. (2020). The circadian clock and darkness control natural competence in cyanobacteria. *Nature Communications*, 11, 1688.

[120] Cassier-Chauvat, C., Veaudor, T., & Chauvat, F. (2016). Comparative genomics of DNA recombination and repair in cyanobacteria: Biotechnological implications. *Frontiers in Microbiology*, 7, 1809.

- [121] Hutchison, C. A., Chuang, R. Y., Noskov, V. N., et al. (2016). Design and synthesis of a minimal bacterial genome. *Science*, 351(6280), aad6253.
- [122] Bernstein, H. C., & Carlson, R. P. (2012). Microbial consortia engineering for cellular factories: In vitro to in silico systems. *Computational and Structural Biotechnology Journal*, 3, e201210017.
- [123] Volova, T., Kiselev, E., Shishatskaya, E., et al. (2013). Cell growth and accumulation of polyhydroxyalkanoates from CO<sub>2</sub> and H<sub>2</sub> in a novel obligate chemolithoautotrophic bacterium *Cupriavidus necator* B-10646. *Bioresource Technology*, 146, 215-222.
- [124] Levy, A., & Borenstein, E. (2013). Metabolic modeling of species interaction in the human microbiome elucidates community-level assembly rules. *Proceedings of the National Academy of Sciences USA*, 110(31), 12804-12809.
- [125] Rummel, J. D., Beaty, D. W., Jones, M. A., et al. (2014). A new analysis of Mars "Special Regions": Findings of the second MEPAG Special Regions Science Analysis Group (SR-SAG2). *Astrobiology*, 14(11), 887-968.
- [126] COSPAR. (2020). COSPAR Policy on Planetary Protection. Committee on Space Research Panel on Planetary Protection.
- [127] Wright, O., Stan, G. B., & Ellis, T. (2013). Building-in biosafety for synthetic biology. *Microbiology*, 159(7), 1221-1235.
- [128] Clauwaert, P., Muys, M., Alloul, A., et al. (2017). Nitrogen cycling in bioregenerative life support systems. *Progress in Aerospace Sciences*, 91, 87-98.
- [129] Landis, G. A. (2001). Solar power for Mars. NASA Technical Memorandum 210231.
- [130] Xue, S., Su, Z., & Cong, W. (2011). Growth of *Spirulina platensis* enhanced under intermittent illumination. *Journal of Biotechnology*, 151(3), 271-277.
- [131] Hendrickx, L., & Mergeay, M. (2007). From the deep sea to the stars: Human life support through minimal communities. *Current Opinion in Microbiology*, 10(3), 231-237.
- [132] Muscatello, A. C., Santiago-Maldonado, E., & Linne, D. L. (2020). Mars In-Situ Resource Utilization technology evaluation. *AIAA Propulsion and Energy Forum*, AIAA 2020-3476.
- [133] Bernstein, H. C., Paulson, S. D., & Carlson, R. P. (2012). Synthetic *Escherichia coli* consortia engineered for syntrophy demonstrate enhanced biomass productivity. *Journal of Biotechnology*, 157(1), 159-166.
- [134] Großkopf, T., & Soyer, O. S. (2014). Synthetic microbial communities. *Current Opinion in Microbiology*, 18, 72-77.
- [135] Radivojević, T., Costello, Z., Workman, K., & Garcia Martin, H. (2020). A machine learning Automated Recommendation Tool for synthetic biology. *Nature Communications*, 11, 4879.
- [136] Long, C. P., & Antoniewicz, M. R. (2019). High-resolution <sup>13</sup>C metabolic flux analysis. *Nature Protocols*, 14(10), 2856-2877.
- [137] Menolascina, F., Siciliano, V., & di Bernardo, D. (2012). Engineering and control of biological systems: A new way to tackle complex diseases. *FEBS Letters*, 586(15), 2122-2128.

- [138] Costello, Z., & Garcia Martin, H. (2018). A machine learning approach to predict metabolic pathway dynamics from time-series multiomics data. *npj Systems Biology and Applications*, 4, 19.
- [139] Lasseur, C., Dixon, M., Paille, C., et al. (2023). MELiSSA: 30 years of research towards sustainable life support. *Acta Astronautica*, 202, 463-478.
- [140] Mankins, J. C. (2009). Technology readiness assessments: A retrospective. *Acta Astronautica*, 65(9-10), 1216-1223.
- [141] McKay, C. P., Toon, O. B., & Kasting, J. F. (1991). Making Mars habitable. *Nature*, 352, 489-496.
- [142] Cockell, C. S. (2010). The interplanetary exchange of photosynthesis. *Origins of Life and Evolution of Biospheres*, 40(4), 397-410.
- [143] Pointing, S. B., & Belnap, J. (2012). Microbial colonization and controls in dryland systems. *Nature Reviews Microbiology*, 10(8), 551-562.
- [144] Gerstell, M. F., Francisco, J. S., Yung, Y. L., et al. (2001). Keeping Mars warm with new super greenhouse gases. *Proceedings of the National Academy of Sciences USA*, 98(5), 2154-2157.
- [145] Ruffing, A. M., Jensen, T. J., & Strickland, L. M. (2016). Genetic tools for advancement of *Synechococcus* sp. PCC 7002 as a cyanobacterial chassis. *Microbial Cell Factories*, 15, 190.
- [146] Madhyastha, H. K., Radha, K. S., Nakajima, Y., et al. (2008). uPA dependent and independent mechanisms of wound healing by C-phycocyanin. *Journal of Cellular and Molecular Medicine*, 12(6B), 2691-2703.
- [147] Burja, A. M., Banaigs, B., Abou-Mansour, E., et al. (2001). Marine cyanobacteria—A prolific source of natural products. *Tetrahedron*, 57(46), 9347-9377.
- [148] Mayfield, S. P., Franklin, S. E., & Lerner, R. A. (2003). Expression and assembly of a fully active antibody in algae. *Proceedings of the National Academy of Sciences USA*, 100(2), 438-442.
- [149] Hempel, F., & Maier, U. G. (2012). An engineered diatom acting like a plasma cell secreting human IgG antibodies with high efficiency. *Microbial Cell Factories*, 11, 126.