

# Summary about Prediction Market

Zhihao Ruan, Jiayu Yi, Naihao Deng  
{ruanzh, yijiaYu, dnaihao}@umich.edu

September 26, 2019

## Contents

<b>1</b>	<b>Exponential Family</b>	<b>2</b>
1.1	Sufficient Statistics . . . . .	2
1.2	Conjugate Priors . . . . .	3
1.2.1	Bernoulli Distribution . . . . .	3
1.2.2	Normal Distribution . . . . .	3
<b>2</b>	<b>Scoring Rules</b>	<b>6</b>
2.1	Motivations . . . . .	6
2.2	Incentive Compatibility . . . . .	6
2.3	Logarithmic Scoring Rule . . . . .	6
2.4	Maximum Entropy Optimization . . . . .	7
2.4.1	Examples . . . . .	7
<b>3</b>	<b>Cost Function based Prediction Market with Bayesian Traders</b>	<b>8</b>
3.1	Bayesian Agents . . . . .	8
3.2	Trading Procedures . . . . .	8
3.2.1	Assumptions and hyperparameters . . . . .	9
3.2.2	Procedures . . . . .	9

# 1 Exponential Family

Distributions over  $\mathbf{x}$  with *natural parameters*  $\boldsymbol{\eta}$  that are of the form

$$p(\mathbf{x}|\boldsymbol{\eta}) = h(\mathbf{x})g(\boldsymbol{\eta}) \exp(\boldsymbol{\eta}^T \mathbf{u}(\mathbf{x})) \quad (1)$$

We need to make sure that  $\int p(\mathbf{x}|\boldsymbol{\eta})d\mathbf{x} = 1$ , therefore

$$g(\boldsymbol{\eta}) \int h(\mathbf{x}) \exp \{ \boldsymbol{\eta}^T \mathbf{u}(\mathbf{x}) \} d\mathbf{x} = 1 \quad (2)$$

$g(\boldsymbol{\eta})$  is a coefficient that is needed to ensure that the distribution is normalized.

## 1.1 Sufficient Statistics

If we take the partial derivative with respect to  $\boldsymbol{\eta}$  from both sides of equation 2, we have

$$\nabla g(\boldsymbol{\eta}) \int h(\mathbf{x}) \exp \{ \boldsymbol{\eta}^T \mathbf{u}(\mathbf{x}) \} d\mathbf{x} + g(\boldsymbol{\eta}) \int h(\mathbf{x}) \exp \{ \boldsymbol{\eta}^T \mathbf{u}(\mathbf{x}) \} \mathbf{u}(\mathbf{x}) d\mathbf{x} = 0 \quad (3)$$

If we plug in 2 to 3, we have

$$-\frac{1}{g(\boldsymbol{\eta})} \nabla g(\boldsymbol{\eta}) = \int h(\mathbf{x}) \exp \{ \boldsymbol{\eta}^T \mathbf{u}(\mathbf{x}) \} \mathbf{u}(\mathbf{x}) d\mathbf{x} = \mathbb{E}[\mathbf{u}(\mathbf{x})] \quad (4)$$

Since

$$\nabla \ln g(\boldsymbol{\eta}) = \frac{1}{g(\boldsymbol{\eta})} \nabla g(\boldsymbol{\eta})$$

It can be deduced that

$$-\nabla \ln g(\boldsymbol{\eta}) = \mathbb{E}[\mathbf{u}(\mathbf{x})] \quad (5)$$

If we consider a set of independent identically distributed data denoted by  $\mathbf{x} = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ , the likelihood function is given by

$$p(\mathbf{x}|\boldsymbol{\eta}) = \left( \prod_{n=1}^N h(\mathbf{x}_n) \right) g(\boldsymbol{\eta})^N \exp \left\{ \boldsymbol{\eta}^T \sum_{n=1}^N \mathbf{u}(\mathbf{x}_n) \right\} \quad (6)$$

To produce the maximum likelihood estimator  $\boldsymbol{\eta}_{ML}$ , we need to take the partial derivative of 6 and set its value to 0.

$$\nabla \left( \left( \prod_{n=1}^N h(\mathbf{x}_n) \right) g(\boldsymbol{\eta})^N \exp \left\{ \boldsymbol{\eta}^T \sum_{n=1}^N \mathbf{u}(\mathbf{x}_n) \right\} \right) = 0 \quad (7)$$

By expanding 7 it can be seen that

$$\left( \prod_{n=1}^N h(\mathbf{x}_n) \right) \left( N g(\boldsymbol{\eta})^{N-1} \nabla g(\boldsymbol{\eta}) \exp \left\{ \boldsymbol{\eta}^T \sum_{n=1}^N \mathbf{u}(\mathbf{x}_n) \right\} + g(\boldsymbol{\eta})^N \exp \left\{ \boldsymbol{\eta}^T \sum_{n=1}^N \mathbf{u}(\mathbf{x}_n) \right\} \sum_{n=1}^N \mathbf{u}(\mathbf{x}_n) \right) = 0$$

Therefore,

$$-\nabla \ln g(\boldsymbol{\eta}_{ML}) = \frac{1}{N} \sum_{n=1}^N \mathbf{u}(\mathbf{x}_n) \quad (8)$$

From equation 8 it can be seen that  $\boldsymbol{\eta}_{ML}$  can be determined if  $\mathbf{u}(\mathbf{x}_n)$  is known,  $\mathbf{u}(\mathbf{x}_n)$  is therefore called the **sufficient statistics**. If  $N \rightarrow \infty$ , the right hand side of 8 would become  $\mathbb{E}[\mathbf{u}(\mathbf{x})]$  and  $\boldsymbol{\eta}_{ML}$  would be the true  $\boldsymbol{\eta}$ .

## 1.2 Conjugate Priors

Conjugate priors are priors that lead to their corresponding posterior distributions to have the same form as they do. For exponential family distributions, their conjugate priors are of the form

$$p(\boldsymbol{\eta}|\boldsymbol{\chi}, \nu) = f(\boldsymbol{\chi}, \nu)g(\boldsymbol{\eta})^\nu \exp \{ \nu \boldsymbol{\eta}^T \boldsymbol{\chi} \} \quad (9)$$

$\nu$  can be interpreted as the number of samples that are observed.  $\boldsymbol{\chi}$  is the mean of the pseudo observation of the sufficient statistics  $\mathbf{u}(\mathbf{x})$ . Each one of the  $\nu$  observations take the value of  $\boldsymbol{\chi}$ . The updated posterior can be shown as

$$\begin{aligned} p(\boldsymbol{\eta}|\mathbf{X}, \boldsymbol{\chi}, \nu) &= \left( \prod_{n=1}^N h(\mathbf{x}_n) \right) g(\boldsymbol{\eta})^N \exp \left\{ \boldsymbol{\eta}^T \sum_{n=1}^N \mathbf{u}(\mathbf{x}_n) \right\} \times f(\boldsymbol{\chi}, \nu)g(\boldsymbol{\eta})^\nu \exp \{ \nu \boldsymbol{\eta}^T \boldsymbol{\chi} \} \\ &\propto g(\boldsymbol{\eta})^{\nu+N} \exp \left\{ \boldsymbol{\eta}^T \left( \sum_{n=1}^N \mathbf{u}(\mathbf{x}_n) + \nu \boldsymbol{\chi} \right) \right\} \end{aligned} \quad (10)$$

It can be seen from here that we can update the posterior with  $\nu \leftarrow \nu + N$  and  $\nu \boldsymbol{\chi} \leftarrow \sum_{n=1}^N \mathbf{u}(\mathbf{x}_n) + \nu \boldsymbol{\chi}$ . **This update rule again emphasizes that  $\nu$  is the number of pseudo observations, and  $\boldsymbol{\chi}$  is the mean of the sufficient statistics  $\mathbf{u}(\mathbf{x})$  of the pseudo observations.**

### 1.2.1 Bernoulli Distribution

The conjugate prior for Bernoulli distribution is the Beta distribution. The probability density function for Beta distribution is

$$p(x|a, b) = \frac{x^{a-1}(1-x)^{b-1}}{\mathbf{B}(a, b)} \quad (11)$$

When it is written in the form of the conjugate prior

$$\frac{\theta^{a-1}(1-\theta)^{b-1}}{\mathbf{B}(a, b)} = \frac{1}{\mathbf{B}(a, b)} (1-\theta)^{a+b-2} \exp \left[ (a+b-2) \ln \left( \frac{\theta}{1-\theta} \right) \frac{a-1}{a+b-2} \right] \quad (12)$$

From equation 12 we can see that  $\nu$  corresponds to  $a+b-2$  here and  $\boldsymbol{\chi}$  corresponds to  $\frac{a-1}{a+b-2}$ .

### 1.2.2 Normal Distribution

#### Problem Statement

Considering a univariate Gaussian distribution with fixed variance ( $\sigma = 1$ ) that:

$$p(x; \mu) = \frac{1}{\sqrt{2\pi}} \exp \left( -\frac{(x - \mu)^2}{2} \right),$$

we have already known that its conjugate prior must be

$$p(\mu; m, v) = \frac{1}{\sqrt{2\pi}v} \exp \left( -\frac{(\mu - m)^2}{2v^2} \right).$$

Since it is an exponential family distribution of general form  $p(x; \theta) = \exp(\langle \theta, x \rangle - T(\theta))$ , its conjugate prior must have the general form of  $p(\theta; n, \nu) = \exp(\langle n\nu, \mu \rangle + nT(\theta) - \psi(\nu, n))$ . We wish to calculate  $n$  and  $\nu$  in terms of  $m$  and  $v$ , in order to finish our simulation of prediction market activities in terms of a univariate Gaussian event with fixed variance.

## Derivations

### Rewrite Gaussian distribution in exponential family form.

From the traditional Gaussian form we have:

$$\begin{aligned} p(x; \mu) &= \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{(x - \mu)^2}{2}\right) \\ &= \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2} + \mu x - \frac{\mu^2}{2}\right) \\ &= \frac{\exp(-x^2/2)}{\sqrt{2\pi}} \exp\left(\mu x - \frac{\mu^2}{2}\right) \\ &= h(x) \exp(\langle \mu, x \rangle - T(\mu)). \end{aligned}$$

Therefore we have the base measure and the log-partition function:

$$\begin{aligned} h(x) &= \frac{\exp(-x^2/2)}{\sqrt{2\pi}}, \\ T(\mu) &= \frac{\mu^2}{2}. \end{aligned}$$

**Decide each element in the prior from general form.**

From the traditional form of prior we know:

$$\begin{aligned}
p(\mu; m, v) &= \frac{1}{\sqrt{2\pi}v} \exp\left(-\frac{(\mu - m)^2}{2v^2}\right) \\
&= \frac{1}{\sqrt{2\pi}v} \exp\left(-\frac{1}{2v^2}\mu^2 + \frac{m}{v^2}\mu - \frac{m^2}{2v^2}\right) \\
&= \frac{1}{\sqrt{2\pi}v} \exp\left(-\frac{1}{v^2}T(\mu) + \frac{m}{v^2}\mu - \frac{m^2}{2v^2}\right) \\
&= h(\mu) \exp(\langle n\nu, \mu \rangle + nT(\mu) - \psi(n, \nu)).
\end{aligned}$$

Therefore we know that

$$\begin{aligned}
n &= -\frac{1}{v^2} \\
\nu &= -m \\
\psi(\nu, n) &= \frac{m^2}{2v^2}.
\end{aligned}$$

**Validation**

We can validate our results by deriving the posterior with respect to  $N$  data points from a Gaussian distribution.

Assume that there is a set of  $N$  points  $S = \{x_i\}_{i=1}^N$  such that

$$p(x_i; \mu) = \frac{\exp(-x_i^2/2)}{\sqrt{2\pi}} \exp\left(\mu x_i - \frac{\mu^2}{2}\right)$$

and  $\mu$  follows the exact prior mentioned above. Then we will have the posterior:

$$\begin{aligned}
p(\mu; m, v | x_1, \dots, x_N) &= \frac{1}{\sqrt{2\pi}v} \exp\left(\left[\frac{m/v^2}{-1/v^2}\right] \cdot \left[\frac{\mu}{T(\mu)}\right] - \frac{m^2}{2v^2}\right) \prod_{i=1}^N \frac{\exp(-x_i^2/2)}{\sqrt{2\pi}} \exp\left(\mu x_i - \frac{\mu^2}{2}\right) \\
&= h(\mu | x_1, \dots, x_N) \exp\left(\left[\frac{m/v^2}{-1/v^2}\right] \cdot \left[\frac{\mu}{T(\mu)}\right] - \frac{m^2}{2v^2}\right) \exp\left(\mu \sum_{i=1}^N x_i - \frac{N}{2}\mu^2\right) \\
&= h(\mu | x_1, \dots, x_N) \exp\left(\left[\frac{\frac{m}{v^2} + \sum_{i=1}^N x_i}{-\frac{1}{v^2} - N}\right] \cdot \left[\frac{\mu}{T(\mu)}\right] - \frac{m^2}{2v^2}\right) \\
&= h(\mu | x_1, \dots, x_N) \exp\left(\left[\frac{\frac{m}{v^2} + \sum_{i=1}^N x_i}{-\frac{1}{v^2} - N}\right] \cdot \left[\frac{\mu}{T(\mu)}\right] - \psi(\nu, n)\right)
\end{aligned}$$

which is still a Gaussian distribution.

Therefore from our derivation we can see that for a univariate Gaussian distribution with  $\sigma^2 = 1$ ,

$$n = -\frac{1}{v^2}, \quad \nu = -m, \quad \psi(\nu, n) = \frac{m^2}{2v^2}.$$

## 2 Scoring Rules

Scoring rules is the simplest form of prediction mechanism. For every agent with some information, a scoring rule evaluates how close the information is from the actual outcome, and pays the agent in return for his/her information.

### 2.1 Motivations

Taking an event  $X$  with outcome space  $\mathcal{X}$ , we want to know something about each agent's belief  $p(x)$  on the actual outcome  $x \in \mathcal{X}$ , compare it with the actual outcome, and see how accurate the agent's prediction is. However, it is impossible to ask agent for his/her entire  $p(x)$  probability distribution. What should we do?

We've already known that sufficient statistic is a very suitable parameter to characterize a probability distribution. Therefore, we can use it to represent agent's belief. We just ask each agent for the sufficient statistic  $u(x)$  as a representation of his/her belief  $p(x)$ . Then we are able to measure how accurate the agent can predict with a scoring rule based on the actual outcome.

Assume  $\mu = \mathbb{E}_p[u(x)]$  is the *expected* sufficient statistic over some  $p(x)$  that takes  $u(x)$  as its sufficient statistic and is taken as an estimate of the agent's belief. Assume  $\hat{\mu}$  is the agent's report as an estimate of  $\mu$ . Then, with the actual outcome denoted  $x$ , the scoring rule takes the form:

$$S(\hat{\mu}, x).$$

We can see that the scoring rule is a measure of how close the agent's belief is from the actual outcome.

### 2.2 Incentive Compatibility

**Incentive compatibility** is a property of prediction mechanism with which the best strategy for an agent to earn the most profit is to *honestly* report all the information as soon as he/she has it. As a prediction mechanism, a proper scoring rule should leverage **incentive compatibility** in order to get real information from agents.

Assume that we already set the sufficient statistic to be  $u(x)$ . Then for each  $p \in \mathcal{P}$  that takes this  $u(x)$  as its sufficient statistic, we can calculate its *expected* sufficient statistic  $\mu = \mathbb{E}_p[u(x)]$ . A scoring rule is thus called **proper** if it satisfies the following, for all such  $p$ , for all  $\hat{\mu} \neq \mu$ :

$$\mathbb{E}_p[S(\mu, x)] \geq \mathbb{E}_p[S(\hat{\mu}, x)].$$

Any scoring rule with this property actually encourages agents to report a probability distribution as close to the actual probability distribution of  $X$  as possible, which is in accordance with the essence of incentive compatibility.

### 2.3 Logarithmic Scoring Rule

Suppose that we have set a form of  $u(x)$ . For some  $p$  that takes  $u(x)$  as its sufficient statistic, a classic logarithmic scoring rule takes the form:

$$S(\mu, x) = \ln p(x; \mu),$$

where  $\mu = \mathbb{E}_p[u(x)]$  is the expected sufficient statistic over  $p(x; \mu)$ . **For the following sections, we will be only talking about logarithmic scoring rules.**

## 2.4 Maximum Entropy Optimization

Now that we have the general expression of logarithmic scoring rules, how can we determine which  $p(\mathbf{x}; \boldsymbol{\mu})$  to choose as an estimate of agents' beliefs, given a specific  $\mathbf{u}(\mathbf{x})$ ?

For some particular form of  $\mathbf{u}(\mathbf{x})$ , denote the set  $\mathcal{P}$  which contains all  $p$  that takes  $\mathbf{u}(\mathbf{x})$  as its sufficient statistic. Then, we formulate the following optimization problem:

$$\begin{aligned} \min_{p \in \mathcal{P}} \quad & F(p) = \int_{\mathbf{x} \in \mathcal{X}} p(\mathbf{x}) \ln p(\mathbf{x}) \, d h(\mathbf{x}) \\ \text{s.t.} \quad & \mathbb{E}_p[\mathbf{u}(\mathbf{x})] = \boldsymbol{\mu} \end{aligned}$$

where  $\boldsymbol{\mu}$  is the expected sufficient statistic over  $p$ , and  $h(\mathbf{x})$  is the base measure.

It is clear to see that  $F(p)$  is actually the negative of entropy of  $p(\mathbf{x})$ . Therefore, minimizing  $F(p)$  is thus maximizing the entropy of  $p$ . This actually mean that of all  $p \in \mathcal{P}$ , we tend to choose a  $p$  that shows the most uncertainty of  $\mathbf{X}$  as an estimate of agents' beliefs. This estimate is not necessarily exactly the same as agents' beliefs, as the general goal is just to find something that we can use to construct a metric in order to measure how close the agents' beliefs are from the actual outcome.

It has been proved that  $S(\boldsymbol{\mu}, \mathbf{x}) = \ln p(\mathbf{x}; \boldsymbol{\mu})$  is **proper** if and only if  $p$  is the exponential family. It has also been proved that the solutions to the optimization problem are exponential family distributions which takes the form

$$\begin{aligned} p(\mathbf{x}; \boldsymbol{\eta}) &= h(\mathbf{x}) g(\boldsymbol{\eta}) \exp \{ \boldsymbol{\eta} \cdot \mathbf{u}(\mathbf{x}) \} \\ &= h(\mathbf{x}) \exp \{ \boldsymbol{\eta} \cdot \mathbf{u}(\mathbf{x}) - T(\boldsymbol{\eta}) \} \end{aligned}$$

where  $h(\mathbf{x})$  is the base measure,  $\boldsymbol{\eta}$  is the natural parameter of  $p$ ,  $T(\boldsymbol{\eta}) = -\ln g(\boldsymbol{\eta})$  is the log-partition function of  $p$ .

Either *expected* sufficient statistic  $\boldsymbol{\mu}$  or natural parameter  $\boldsymbol{\eta}$  can parametrize a probability distribution  $p$ , and actually they are closely related. If we denote the convex conjugate of  $T(\boldsymbol{\eta})$  as  $G(\boldsymbol{\mu})$ , then we have  $\boldsymbol{\mu} = \nabla T(\boldsymbol{\eta})$ ,  $\boldsymbol{\eta} = \nabla G(\boldsymbol{\mu})$ .

### 2.4.1 Examples

1. If we take the outcome space  $\mathcal{X} = [0, +\infty)$  and  $u(x) = x$ , the maximum entropy optimization produces an exponential distribution. Hence, the log scoring rule becomes

$$S(\mu, x) = -\frac{x}{\mu} - \ln \mu.$$

If we take the partial derivative with respect to  $\mu$

$$\frac{\partial S}{\partial \mu} = \frac{x}{\mu^2} - \frac{1}{\mu} = \frac{x - \mu}{\mu^2},$$

we can see that the scoring rule reaches maximum when  $\mu = x$ . This actually means that agents should report information as close to the actual outcome as possible in order to maximize profit, which corresponds to the essence of incentive compatibility.

2. If we take the outcome space  $\mathcal{X} = \mathbb{R}$  and  $\mathbf{u}(x) = (x, x^2)$ , the maximum entropy optimization produces a Gaussian distribution. Hence, the log scoring rule becomes

$$S((\mu, \sigma^2), x) = -\frac{(x - \mu)^2}{\sigma^2} - \ln \sigma^2.$$

As this scoring rule is a basic parabola with respect to  $\mu$ , it is also clear that it reaches the maximum when  $\mu = x$ . Similarly, this scoring rule is also incentive compatible.

### 3 Cost Function based Prediction Market with Bayesian Traders

Based on what we have in previous sections, we could simulate the prediction market by a Python program.

For the setup, there would be a group of Bayesian agents (the number of agents as hyper-parameter would be an input for the program), a market maker and an executable program driving the market. They are explained in Sec 3.1 and Sec 3.2. The dataset the market is based on is subject to Bernoulli distribution.

#### 3.1 Bayesian Agents

Each Bayesian agents would be initialized with several data points, based on which he would form his prior  $p(\boldsymbol{\eta}; \boldsymbol{\chi}, \nu)$ . The dataset is subject to Bernoulli distribution, in which  $n$  would be the number of trades the market maker publishes,  $\boldsymbol{\chi}$  would be the current market price.

Suppose each agent would have observed  $m$  data points before his or her entrance to the market. Since he knows that each agent would have access to  $m$  data points, we thus have

$$\nu = nm.$$

He would update his belief by

$$\begin{aligned} nm &\leftarrow nm + m \\ \nu\boldsymbol{\chi} &\leftarrow \sum_{n=1}^N \mathbf{u}(\mathbf{x}_n) + \nu\boldsymbol{\chi} \end{aligned}$$

Since we are using Bernoulli distribution, the  $\mathbf{x}_n$  in question is just a scalar that can take the value of either 0 or 1. The sufficient statistics for Bernoulli distribution is  $x$ . In other words,

$$\mathbf{u}(x) = x$$

Since there is only one entry for the sufficient statistics, parameter  $\boldsymbol{\eta}$  would be a scalar, which can be denoted as  $\eta$ . The agent can also derive the amount of outstanding shares  $\eta$  from the current market price  $\xi$ , in our case,

$$\xi(\eta) = \nabla_{\eta} \exp(g(\eta)) = \nabla_{\eta} \ln(1 + e^{\eta}) = \frac{e^{\eta}}{1 + e^{\eta}} \quad (13)$$

Based on

$$\xi(\eta + \delta) = \frac{nm\chi + \sum_{n=1}^m x_n}{nm + m}$$

the agent would be able to decide  $\delta$ , the amount he would like to sell is

$$\delta = \ln \left( \frac{nm\chi + \sum_{i=1}^m x_i}{nm + m - (nm\chi + \sum_{i=1}^m x_i)} \right) - \eta \quad (14)$$

#### 3.2 Trading Procedures

Based on the mathematical background above, we can derive a general trading procedure for the prediction market with Bayesian agents.



### 3.2.1 Assumptions and hyperparameters

For simplification, we assume that each agent has access to their own private set of data points of size  $m$ , which means that the number of data points that each agents observed is constant.

Throughout the entire trading process, the market maker will always post the current market price  $\mu$  and the number of trades  $n$ .

### 3.2.2 Procedures

1. The market maker sets an initial market price  $\chi = \chi_0$ , and sets the number of trades  $n = 0$ .
2. An agent comes into the market, before which he has observed  $m$  data points  $\{x_1, \dots, x_m\}$  in private.
3. He calculates

$$\hat{\mu} = \frac{1}{m} \sum_{i=1}^m \mathbf{u}(x_i) = \frac{1}{m} \sum_{i=1}^m x_i$$

as his own prior belief.

4. He sees the current market price  $\chi$  and the number of trades  $n$ . He then knows that  $n$  agents have traded in the market before him. Under the assumption that all agents observe the same number of data points, the total number of initial pseudo observations would be  $nm$  with an average of  $\chi$ .
5. He updates his prior belief and obtains his posterior by

$$\mu' = \frac{nm\chi + m\hat{\mu}}{nm + m}.$$

6. He trades some shares in the market in order to change the current market price to  $\mu'$ .
7. The current market price is thus changed to

$$\chi \leftarrow \frac{nm\chi + m\hat{\mu}}{nm + m}, \quad n \leftarrow n + 1.$$