

Kähler structure on reduction via complex manifolds

Shing Tak Lam

July 19, 2023

Throughout, let M be a compact Kähler manifold, G a compact Lie group acting on X preserving the Kähler structure. Let $\mu : M \rightarrow \mathfrak{g}^*$ be the moment map for the action. Suppose in addition that G acts freely on $\mu^{-1}(0)$. Then we have that

$$M//G := \mu^{-1}(0)/G$$

has a Kähler structure. In this note, we show another way of thinking about this, which is that

$$\mu^{-1}(0)/G \cong M^{\min}/G_{\mathbb{C}}$$

where M^s is an open submanifold of M , and $G_{\mathbb{C}}$ is the complexification of G . In this case, the fact that $M//G$ has a complex structure becomes clear.

Fix a G -invariant Riemannian metric on M , and define $f : M \rightarrow \mathbb{R}$ by

$$f(x) = \|\mu(x)\|^2$$

We want to consider f as a Morse function on M . For $x \in M$, define the curve $(x_t)_{t \geq 0}$ by

$$\begin{aligned} x_0 &= x \\ \frac{dx_t}{dt} &= -\text{grad } f(x_t) \end{aligned}$$

The set of limit points of the trajectory is

$$\omega(x) = \{y \in M \mid \text{every neighbourhood of } y \text{ contains } x_t \text{ for } t \text{ arbitrarily large}\}$$

With this, we define

$$M^{\min} = \{x \in X \mid \omega(x) \subseteq \mu^{-1}(0)\}$$

Then M^{\min} is a $G_{\mathbb{C}}$ invariant open subset of M , with $\mu^{-1}(0) \subseteq M^{\min}$. The inclusion map induces a natural continuous map

$$\mu^{-1}(0)/G \rightarrow M^{\min}/G_{\mathbb{C}}$$

which is a homeomorphism. The proof of this relies on the fact that $G \cdot \mu^{-1}(0) = M^{\min}$, hence the map is surjective. Another argument shows the map is bijective. Therefore it is a continuous bijection from a compact space to a Hausdorff space, hence a homeomorphism.