Algebraic Topology

Shing Tak Lam

Michaelmas 2023*

Lecture 1

1 Introduction

Let X, Y be topological spaces, $f, g : X \to Y$ be continuous functions. Note we will use the terminology 'map := continuous function'. We say that f *is homotopic to* g, written $f \simeq g$, if there exists a map $F : X \times I \to Y$, such that

$$F|_{X\times\{0\}} = f \quad \textit{and} \quad F|_{X\times\{1\}} = g$$

Here $I = [0, 1]$ carries its Euclidean topology. An exercise (sheet 1) is that this defines an equivalence relation on the set of maps $X \to Y$.

Definition 1.1 (homotopy equivalence)
We say that $f : X \to Y$ is a *homotopy equivalence* if there exists a function $g : Y \to X$ such that $f \circ g \simeq \mathrm{id}_Y$ and $g \circ f \simeq \mathrm{id}_X$. We say that g is a *homotopy inverse* to f.

The same exercise on sheet 1 shows that homotopy equivalence is an equivalence relation on topological spaces.

Example 1.2
1. If $f : X \to Y$ is a homeomorphism, then f is also a homotopy equivalence.
2. $i : \{0\} \hookrightarrow \mathbb{R}^n$ is a homotopy equivance.
3. The inclusion $i : S^{n-1} \hookrightarrow \mathbb{R}^n \setminus 0$ is a homotopy equivalence.

Definition 1.3
A space X is *contractible* if it is homotopy equivalent to a point.

Idea of Algebraic Topology: Study spaces up to homotopy equivalence. Broadly, we are interested in "connectivity properties" of topological spaces.

Example 1.4
1. We say a space X is *path-connected* if any two maps $\{*\} \to X$ are homotopic. For example, $\mathbb{R}$ is path connected, $\mathbb{R} \setminus \{0\}$ is not. A corollary of this is the intermediate value theorem.
2. We say a path connected space X is *simply connected* if every map $f : S^1 \to X$ is homotopic to a constant map. Equivalently, every two maps $S^1 \to X$ are homotopic. Or equivalently (again) every continuous map $S^1 \to X$ extends to a continuous map $D^2 \to X$. For example, $\mathbb{R}^2$ is simply connected, by $\mathbb{R}^2 \setminus \{0\}$ is not. If $\gamma : S^1 \to \mathbb{R}^2 \setminus \{0\}$ is continuous, it has a *winding number*, or *degree* $\deg(\gamma) \in \mathbb{Z}$, such that
   - It is invariant under homotopy,
   - If $\gamma_n(t) = \exp(2n\pi it)$, then $\deg(\gamma_n) = n$

*Based on lectures by Ivan Smith. Last updated October 6, 2023.

In particular, taking $n = 0$ gives the constant map. Therefore, a constant map has degree 0. A corollary of this is the fundamental theorem of algebra.

3. We say a path connected space X is *k-connected*, if for all $i \leq k$, every map $S^i \to X$ is homotopic to a constant map. In this case, $\mathbb{R}^n$ is $(n-1)$-connected, but $\mathbb{R}^n \setminus 0$ is not. More precisely, maps from $S^{n-1} \to \mathbb{R}^n \setminus 0$ have a degree $\deg(f) \in Z$, such that degree is a homotopy invariant, and the degree of the inclusion map is 1, and the degree of the constant map is 0. A corollary of this is the Brouwer fixed point theorem.

Index

k-connected, 2

contractible, 1

homotopic, 1
homotopy equivalence, 1

path connected, 1

simply connected, 1