

# **Math Camp: Lesson 1**

**Basics, Notation, Pre-Calculus**

**UW–Madison Political Science**

**August 17, 2020**

# Welcome

First...

Numbers

# Math: not just numbers

Math is a general framework for manipulating various concepts, *one of which* is "numbers"

Types of numbers:

- Integers (whole numbers, including negative):  $\mathbb{Z}$
- Real numbers (the continuous number line):  $\mathbb{R}$
- Positive and negative real numbers:  $\mathbb{R}^+$  and  $\mathbb{R}^-$
- Real numbers in  $n$  dimensions:  $\mathbb{R}^n$
- Complex numbers:  $\mathbb{C}$ 
  - $1 + 2i$
  - where  $i = \sqrt{-1}$

# Special numbers

0 and 1:

# Special numbers

0 and 1:

- Useful for identities in addition and multiplication

- $4 + 0 = 4$

- $1 * x = x$

# Special numbers

0 and 1:

- Useful for identities in addition and multiplication
  - $4 + 0 = 4$
  - $1 * x = x$
- Useful for turning "off" and "on" pieces of an equation
  - We have an equation:  $4 + z(5x) + (1 - z)3x$

# Special numbers

0 and 1:

- Useful for identities in addition and multiplication
  - $4 + 0 = 4$
  - $1 * x = x$
- Useful for turning "off" and "on" pieces of an equation
  - We have an equation:  $4 + z(5x) + (1 - z)3x$
  - What happens if  $z = 0$ ?

# Special numbers

0 and 1:

- Useful for identities in addition and multiplication
  - $4 + 0 = 4$
  - $1 * x = x$
- Useful for turning "off" and "on" pieces of an equation
  - We have an equation:  $4 + z(5x) + (1 - z)3x$
  - What happens if  $z = 0$ ?
  - If  $z = 1$ ?

# Special numbers

$\pi$ : Approximately 3.1415926 ...

- Ratio of a circle's circumference to its diameter
- Useful for angles, circles, polar coordinates (which you may use!)

# Special numbers

$\pi$ : Approximately 3.1415926 ...

- Ratio of a circle's circumference to its diameter
- Useful for angles, circles, polar coordinates (which you may use!)

$e$ : Euler's number

- Approximately 2.718281828 ...
- Derivative of  $e^x$  is equal to  $e^x$
- Very important for probability and statistics

# Special numbers

$\pi$ : Approximately 3.1415926 ...

- Ratio of a circle's circumference to its diameter
- Useful for angles, circles, polar coordinates (which you may use!)

$e$ : Euler's number

- Approximately 2.718281828 ...
- Derivative of  $e^x$  is equal to  $e^x$
- Very important for probability and statistics

$i$ : The imaginary number,  $\sqrt{-1}$ . For advanced algebra

# Special numbers

$\pi$ : Approximately 3.1415926 ...

- Ratio of a circle's circumference to its diameter
- Useful for angles, circles, polar coordinates (which you may use!)

$e$ : Euler's number

- Approximately 2.718281828 ...
- Derivative of  $e^x$  is equal to  $e^x$
- Very important for probability and statistics

$i$ : The imaginary number,  $\sqrt{-1}$ . For advanced algebra

$\infty$ : Infinity. Not really a number but a boundless quantity

# **Numbers and variables**

We can manipulate numbers with simple operations

# Numbers and variables

We can manipulate numbers with simple operations

- addition:  $1 + 2$

# Numbers and variables

We can manipulate numbers with simple operations

- addition:  $1 + 2$
- subtraction:  $3 - 4$

# Numbers and variables

We can manipulate numbers with simple operations

- addition:  $1 + 2$
- subtraction:  $3 - 4$
- multiplication:  $5 \times 6$

# Numbers and variables

We can manipulate numbers with simple operations

- addition:  $1 + 2$
- subtraction:  $3 - 4$
- multiplication:  $5 \times 6$
- division:  $7 \div 8$

# Numbers and variables

We can manipulate numbers with simple operations

- addition:  $1 + 2$
- subtraction:  $3 - 4$
- multiplication:  $5 \times 6$
- division:  $7 \div 8$

We can *generalize* these statements by using *variables*

- $1 + x$
- $3 - y$
- $5 \times a$
- $7 \div m$

*Variable*: a symbol to represent an entity that could take different values

**With me so far?**

# Equations

Statements about equality and inequality

# Equations

Equality: left-hand and right-hand sides (LHS and RHS) are substitutable.

If  $a = 7x$ , then  $8 * a$  and  $8 * 7x$  are equivalent statements

# Equations

Equality: left-hand and right-hand sides (LHS and RHS) are substitutable.

If  $a = 7x$ , then  $8 * a$  and  $8 * 7x$  are equivalent statements

- Usually used to find *new ways to express a known idea* or to *derive implications from known ideas*

# Equations

Equality: left-hand and right-hand sides (LHS and RHS) are substitutable.

If  $a = 7x$ , then  $8 * a$  and  $8 * 7x$  are equivalent statements

- Usually used to find *new ways to express a known idea* or to *derive implications from known ideas*

*Inequalities* show whether LHS or RHS is greater

If  $x + 3 > 2$ , then the sum of  $x + 3$  will be less than 2

- Usually we use them to find the *conditions under which a statement holds*

# Solving equalities and inequalities

Isolate an unknown quantity to express it in terms of known quantities

# Solving equalities and inequalities

Isolate an unknown quantity to express it in terms of known quantities

Solve for  $x$ :

$$-7x + 3 = 12$$

# Solving equalities and inequalities

Isolate an unknown quantity to express it in terms of known quantities

Solve for  $x$ :

$$-7x + 3 = 12$$

Solve for  $x$ :

$$-7x + 3 > 12$$

# Solving equalities and inequalities

Isolate an unknown quantity to express it in terms of known quantities

Solve for  $x$ :

$$-7x + 3 = 12$$

Solve for  $x$ :

$$-7x + 3 > 12$$

Remember to flip an inequality if you multiply or divide by a negative number!

# Data

# Data

The information we record about what we study

- **Cases:** The units being studied (rows)
- **Variables:** Characteristics that describe units (columns)
- **Values:** Specific realization of a variable (cells)

Establishment	Location	Coffee	Vibe	Notable Flaw
Aldo's	Campus	7	good	no plugs
Ancora	Capitol	8	great	hours
Barriques	Capitol	6	good	bathroom key
Colectivo	State St.	7	fair	spotty wifi
Fair Trade	State St.	8	good	expensive, tables
Michelangelo's	Capitol	5	meh	bathroom key
Steep & Brew	Bascom Hill	0	fair	bad coffee (espresso OK)

# Classifying data

Quantitative vs. Qualitative *analysis*

- Broad, sometimes contentious, arguably artificial divides in the study of politics
- Quantitative: larger  $n$ , statistical description and inference
- Qualitative: smaller  $n$ , rich description, non-statistical inference

Quantitative vs. Qualitative *variables*

- Quantitative: countable, numeric (age, number of toes)
- Qualitative: not countable but descriptive (gender, party preference)
- Unlike qualitative *research*, qualitative variables can still be organized in a data table

I advise not getting too hung up these classification systems. Use them only until you can lose them

# Discrete vs. Continuous

## Discrete

- Variables take specific values from a finite set of possible values
- Could be categories, could be discrete numbers
- Layer of the atmosphere, number of parties, country of origin

## Dichotomous

- Special type of discrete variable, two possible values
- 0 or 1, yes or no, war or peace, win or lose, voted or not

## Continuous

- Variables take values from a continuous number line
- Could be a bounded number line
- GDP, vote share, percentage of turnout, unemployment rate

# Related: the "levels of measurement"

## Nominal / Categorical

- *Unordered* categories
- e.g. party affiliation, gender, country of origin, occupation

## Ordinal

- *Ordered or ranked* outcomes
- Could be categories, but numbers are possible (e.g. rankings)
- No fixed "distance" between levels
- e.g. highest level of educational attainment, ranking of countries by the level of corruption, issue prioritization

# Related: the "levels of measurement"

## Interval

- Ordered values with *fixed distance between levels*
- But *no true zero point*
- Issue scales, day of the year, Likert scales (debatable)

## Ratio:

- Ordered, fixed intervals, *and true zero*
- Vote percentage, turnout, minutes in line to vote

# Why do these categorizations matter?

Different types of variables lend themselves to different types of analysis

# Why do these categorizations matter?

Different types of variables lend themselves to different types of analysis

- Types of statistics you calculate

# Why do these categorizations matter?

Different types of variables lend themselves to different types of analysis

- Types of statistics you calculate
- Possibilities for data visualization

# Why do these categorizations matter?

Different types of variables lend themselves to different types of analysis

- Types of statistics you calculate
- Possibilities for data visualization
- Statistical models (you will learn if you take 812, 813, MLE, etc.)

# Why do these categorizations matter?

Different types of variables lend themselves to different types of analysis

- Types of statistics you calculate
- Possibilities for data visualization
- Statistical models (you will learn if you take 812, 813, MLE, etc.)
  - **Discrete/Nominal**: logistic regression (logit), multinomial logit,  $\chi^2$  (chi-square)

# Why do these categorizations matter?

Different types of variables lend themselves to different types of analysis

- Types of statistics you calculate
- Possibilities for data visualization
- Statistical models (you will learn if you take 812, 813, MLE, etc.)
  - **Discrete/Nominal**: logistic regression (logit), multinomial logit,  $\chi^2$  (chi-square)
  - **Ordinal**: ordinal logit, also  $\chi^2$

# Why do these categorizations matter?

Different types of variables lend themselves to different types of analysis

- Types of statistics you calculate
- Possibilities for data visualization
- Statistical models (you will learn if you take 812, 813, MLE, etc.)
  - **Discrete/Nominal**: logistic regression (logit), multinomial logit,  $\chi^2$  (chi-square)
  - **Ordinal**: ordinal logit, also  $\chi^2$
  - **Interval**: least-squares regression

# Why do these categorizations matter?

Different types of variables lend themselves to different types of analysis

- Types of statistics you calculate
- Possibilities for data visualization
- Statistical models (you will learn if you take 812, 813, MLE, etc.)
  - **Discrete/Nominal**: logistic regression (logit), multinomial logit,  $\chi^2$  (chi-square)
  - **Ordinal**: ordinal logit, also  $\chi^2$
  - **Interval**: least-squares regression
  - **Ratio**: least-squares, count/rate models, duration/survival models

**Everyone on board?**

# Sets

(useful for "speaking math" about data)

# Sets

Collections of objects or entities

# Sets

Collections of objects or entities

Sets contain "elements" or "members"

- if  $o$  is in set  $A$ , then we write  $o \in A$
- $o$  "is an element of"  $A$

# Sets

Collections of objects or entities

Sets contain "elements" or "members"

- if  $o$  is in set  $A$ , then we write  $o \in A$
- $o$  "is an element of"  $A$

Sets could contain individual numbers, but they could contain other sorts of entities

- vectors, matrices
- functions, probability distributions

We need *only some* set notation to help us work with data

# What's in a set?

A set could contain individual elements, written as

$$A = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$$

# What's in a set?

A set could contain individual elements, written as

$$A = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$$

A set could contain intervals

- $B = [1, 10]$  (square brackets imply that endpoints are included)

# What's in a set?

A set could contain individual elements, written as

$$A = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$$

A set could contain intervals

- $B = [1, 10]$  (square brackets imply that endpoints are included)
- Is it the case that  $A = B$ ?

# What's in a set?

A set could contain individual elements, written as

$$A = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$$

A set could contain intervals

- $B = [1, 10]$  (square brackets imply that endpoints are included)
- Is it the case that  $A = B$ ?
- $C = (0, 11)$  (parentheses indicate that endpoints are not included)
- Does  $B = C$ ?

# Set notation

$\cup$ : the *union* of two sets

- elements that are members of either set
- if  $A = \{1, 2\}$  and  $B = \{2, 3\}$ , then  $A \cup B = \{1, 2, 3\}$

# Set notation

$\cup$ : the *union* of two sets

- elements that are members of either set
- if  $A = \{1, 2\}$  and  $B = \{2, 3\}$ , then  $A \cup B = \{1, 2, 3\}$

$\cap$ : the *intersection* of two sets

- elements that are members of both sets
- if  $A = \{1, 2\}$  and  $B = \{2, 3\}$ , then  $A \cap B = \{2\}$

# Set notation

$\cup$ : the *union* of two sets

- elements that are members of either set
- if  $A = \{1, 2\}$  and  $B = \{2, 3\}$ , then  $A \cup B = \{1, 2, 3\}$

$\cap$ : the *intersection* of two sets

- elements that are members of both sets
- if  $A = \{1, 2\}$  and  $B = \{2, 3\}$ , then  $A \cap B = \{2\}$

(these concepts are helpful for probabilities)

# Set notation

$\cup$ : the *union* of two sets


- elements that are members of either set
- if  $A = \{1, 2\}$  and  $B = \{2, 3\}$ , then  $A \cup B = \{1, 2, 3\}$

$\cap$ : the *intersection* of two sets

- elements that are members of both sets
- if  $A = \{1, 2\}$  and  $B = \{2, 3\}$ , then  $A \cap B = \{2\}$

(these concepts are helpful for probabilities)

$\emptyset$ : the empty set (null set)



<https://bosker.wordpress.com/2013/07/10/venn-diagram-partitioning/>

# Symbols and Set Notation

A handful of symbols are commonly used when we represent data mathematically.

Symbol	Meaning
$>$	greater than
$\geq$	greater than or equal to
$<$	less than
$\leq$	less than or equal to
$\approx$	approximately equal to ( $x \approx y$ )
$\equiv$	equivalent to (for establishing identities)
$\propto$	proportional to ( $4x \propto x$ )

# Symbols and Set Notation

Symbol	Meaning
$\in$	is an element of a set ( $x_i \in \mathbf{X}$ )
$\neg$	not
/	not (if through a symbol: $a_i \notin \mathbf{B}$ )
	given that ( $A   B$ )
$\rightarrow$	implies ( $A \rightarrow B$ )
$\leftrightarrow$	if and only if ( $[x = y] \leftrightarrow [y = x]$ )

# Subsets

if  $A = \{1, 2\}$  and  $B = \{1, 2, 3\}$ :

# Subsets

if  $A = \{1, 2\}$  and  $B = \{1, 2, 3\}$ :

- $A \subseteq B$  means "A is a subset of B"

# Subsets

if  $A = \{1, 2\}$  and  $B = \{1, 2, 3\}$ :

- $A \subseteq B$  means "A is a subset of B"
- Can  $A$  be a subset of  $A$ ?

# Subsets

if  $A = \{1, 2\}$  and  $B = \{1, 2, 3\}$ :

- $A \subseteq B$  means "A is a subset of B"
- Can  $A$  be a subset of  $A$ ?
- using  $\subseteq$ , yes it can

# Subsets

if  $A = \{1, 2\}$  and  $B = \{1, 2, 3\}$ :

- $A \subseteq B$  means "A is a subset of B"
- Can  $A$  be a subset of  $A$ ?
- using  $\subseteq$ , yes it can

"Proper subset"

# Subsets

if  $A = \{1, 2\}$  and  $B = \{1, 2, 3\}$ :

- $A \subseteq B$  means "A is a subset of B"
- Can  $A$  be a subset of  $A$ ?
- using  $\subseteq$ , yes it can

"Proper subset"

- $A \subset B$
- $A$  isn't a proper subset of  $A$
- proper subsets can't be equivalent to their superset

# Indexing

Variables can be sets, and they can take different values for different individuals in a dataset. It is convenient to *index* individual observations using a subscript (typically  $i$ ).

Student	Math Courses in College
1	3
2	0
3	1
4	4

If  $x$  represents the number of math courses,  $x_i$  refers to the  $i$ th observation in  $x$

- $x_1 = 3$
- $x_2 = 0$
- $x_3 = ?$
- $x_4 = ?$

**Set practice**

# Consider the following sets

$$A = \{10, 20, 30\}$$

$$B = [1, 10]$$

$$C = (0, 40)$$

# Consider the following sets

$$A = \{10, 20, 30\}$$

$$B = [1, 10]$$

$$C = (0, 40)$$

- What is  $A \cap B$ ?

# Consider the following sets

$$A = \{10, 20, 30\}$$

$$B = [1, 10]$$

$$C = (0, 40)$$

- What is  $A \cap B$ ?
- Is  $B \subseteq C$ ?

# Consider the following sets

$$A = \{10, 20, 30\}$$

$$B = [1, 10]$$

$$C = (0, 40)$$

- What is  $A \cap B$ ?
- Is  $B \subseteq C$ ?
- Is  $A_2 \in B$ ?

# Consider the following sets

$$A = \{10, 20, 30\}$$

$$B = [1, 10]$$

$$C = (0, 40)$$

- What is  $A \cap B$ ?
- Is  $B \subseteq C$ ?
- Is  $A_2 \in B$ ?
- Which elements in  $A$  are subsets of  $B$ ? Subsets of  $C$ ?

# Consider the following sets

$$A = \{10, 20, 30\}$$

$$B = [1, 10]$$

$$C = (0, 40)$$

- What is  $A \cap B$ ?
- Is  $B \subseteq C$ ?
- Is  $A_2 \in B$ ?
- Which elements in  $A$  are subsets of  $B$ ? Subsets of  $C$ ?
- Is  $(A \cup C) \subset C$ ?

# Set notation in actual research

Almost verbatim from a paper about congressional votes ("roll call votes")<sup>\*</sup>

- The data consist of  $n$  legislators voting on  $m$  different roll calls [bills].
- Each roll call  $j = 1, \dots, m$  presents legislators  $i = 1, \dots, n$  with a choice between a 'Yea' position  $\zeta_j$  and a 'Nay' position  $\Psi_j$ , locations in  $\mathbb{R}$
- Let  $y_{ij} = 1$  if legislator  $i$  votes Yea on the  $j$ th roll call and  $y_{ij} = 0$  otherwise.

Legislator ( $i$ )	Bill ( $j$ )	Vote ( $y$ )
1	1	0
1	2	1
$\vdots$	$\vdots$	$\vdots$
$n$	$m$	$y_{nm}$

\* - Clinton, Jackman, and Rivers. "The Statistical Analysis of Roll Call Data." *APSR* 2004

# Functions

# Functions

Operations or rules of assignment that map an input to a *unique* (that is, exactly one) output

# Functions

Operations or rules of assignment that map an input to a *unique* (that is, exactly one) output

- inputs: also called arguments
- outputs: also called values
- mapping: also called the *definition*

# Functions

Operations or rules of assignment that map an input to a *unique* (that is, exactly one) output

- inputs: also called arguments
- outputs: also called values
- mapping: also called the *definition*
- value =  $f(\text{argument})$
- $f = [\text{insert definition}]$

# Functions

Operations or rules of assignment that map an input to a *unique* (that is, exactly one) output

- inputs: also called arguments
- outputs: also called values
- mapping: also called the *definition*
- value =  $f(\text{argument})$
- $f = [\text{insert definition}]$

Analogies include:

- algorithms, machines, black box, routinized process

# Functions

We can generically refer to functions using  $f(x)$  or  $g(x)$ , just as we might use variables to stand in for values.

# Functions

We can generically refer to functions using  $f(x)$  or  $g(x)$ , just as we might use variables to stand in for values.

- e.g. let  $f(p) = \log\left(\frac{p}{1-p}\right)$

# Functions

We can generically refer to functions using  $f(x)$  or  $g(x)$ , just as we might use variables to stand in for values.

- e.g. let  $f(p) = \log\left(\frac{p}{1-p}\right)$
- $f(0.5) = ?$

# Functions

We can generically refer to functions using  $f(x)$  or  $g(x)$ , just as we might use variables to stand in for values.

- e.g. let  $f(p) = \log\left(\frac{p}{1-p}\right)$
- $f(0.5) = ?$

Other symbols also are fine, e.g.

- $\Phi(\cdot)$
- $\Gamma(\cdot)$
- $B(\cdot)$
- $\Lambda(\cdot)$

# Operators

*Operators* are components of a function that tell what to do with the inputs to produce the outputs

Several common operators serve as the building blocks of mathematical functions

- +
- -
- ×
- /

# Operators

*Operators* are components of a function that tell what to do with the inputs to produce the outputs

Several common operators serve as the building blocks of mathematical functions

- +
- -
- ×
- /

Order of operations:

- Operations within parentheses
- Exponents
- Multiplication and division (left to right)
- Addition and subtraction (left to right)

# Function examples

---

$$x \ y \ z \quad f(x, y) = x - y \quad g(z) = 2z - 1 \quad h(x, y, z) = \frac{x + y}{z}$$

---

5 0 5

2 5 8

0 3 9

3 2 0

8 4 2

1 2 4

---

# Function examples

$x$	$y$	$z$	$f(x, y) = x - y$	$g(z) = 2z - 1$	$h(x, y, z) = \frac{x + y}{z}$
5	0	5	5	9	1
2	5	8	-3	15	.875
0	3	9	-3	17	.333
3	2	0	1	-1	<i>undefined</i>
8	4	2	4	3	6
1	2	4	-1	7	.75

# Nested functions

Given that all functions do is map an input to an output, we can nest functions

- Imagine we perform one function  $f(\cdot)$
- Output from  $f(\cdot)$  serves as the input for  $g(\cdot)$
- $g(f(\cdot))$

# Nested functions

Given that all functions do is map an input to an output, we can nest functions

- Imagine we perform one function  $f(\cdot)$
- Output from  $f(\cdot)$  serves as the input for  $g(\cdot)$
- $g(f(\cdot))$

Functions are evaluated "inside out" following the order of operations

# Nested functions

Given that all functions do is map an input to an output, we can nest functions

- Imagine we perform one function  $f(\cdot)$
- Output from  $f(\cdot)$  serves as the input for  $g(\cdot)$
- $g(f(\cdot))$

Functions are evaluated "inside out" following the order of operations

Suppose  $x = 3, f(x) = 2x + 5, g(x) = x - 7$

# Nested functions

Given that all functions do is map an input to an output, we can nest functions

- Imagine we perform one function  $f(\cdot)$
- Output from  $f(\cdot)$  serves as the input for  $g(\cdot)$
- $g(f(\cdot))$

Functions are evaluated "inside out" following the order of operations

Suppose  $x = 3, f(x) = 2x + 5, g(x) = x - 7$

$$f(g(x)) = ?$$

# Nested functions

Given that all functions do is map an input to an output, we can nest functions

- Imagine we perform one function  $f(\cdot)$
- Output from  $f(\cdot)$  serves as the input for  $g(\cdot)$
- $g(f(\cdot))$

Functions are evaluated "inside out" following the order of operations

Suppose  $x = 3, f(x) = 2x + 5, g(x) = x - 7$

$$f(g(x)) = ?$$

$$g(f(x)) = ?$$


# Piecewise-defined functions

A *piecewise-defined function* has different definitions for different regions of the domain (domain = function inputs)

# Piecewise-defined functions

A *piecewise-defined function* has different definitions for different regions of the domain (domain = function inputs)


$$f(x) = \begin{cases} 0 & \text{if } x < 0 \\ x & \text{if } x \in [0, 1] \\ x^2 & \text{if } x > 1 \end{cases}$$




# Piecewise-defined functions

A *piecewise-defined function* has different definitions for different regions of the domain (domain = function inputs)

$$f(x) = \begin{cases} 0 & \text{if } x < 0 \\ x & \text{if } x \in [0, 1] \\ x^2 & \text{if } x > 1 \end{cases}$$



$$g(x) = \begin{cases} x + 3 & \text{if } x \in (-\infty, 0) \\ 4 & \text{if } x = 0 \\ x - 3 & \text{if } x \in (0, \infty) \end{cases}$$




## Function practice

# Practice


Sketch graphs of the following functions:

- $p(x) = 2x - 1$ , on the interval  $[-2, 2]$
- $q(x) = x * -1^x$ , for integers  $\{0, 1, 2, 3, 4, 5\}$
- $r(x) = 2x^2 - 3x + 4$ , on the interval  $(0, 4)$


$p(x) = 2x - 1$ , on the interval  $[-2, 2]$



$$q(x) = x * -1^x, \text{ for integers } \{0, 1, 2, 3, 4, 5\}$$



$$r(x) = 2x^2 - 3x + 4, \text{ on the interval } (0, 4)$$




## Important functions, routines, and properties

# Limits

*Limits* will help us formally define concepts in this and other lectures.

A *limit* describes a function's behavior at a given input:  $\lim_{x \rightarrow 0} (2 - x^2) = 2$

...or as the input value changes:  $\lim_{x \rightarrow \infty} (2 - x^2) = -\infty$




# Continuity

A function is *continuous* if it has no gaps or jumps.

$$\lim_{x \rightarrow a} f(x) = f(a)$$

Meaning... small changes in input produce small changes in output



# Continuity and discontinuity in political science research

Why we care about continuity

# Continuity and discontinuity in political science research

Why we care about continuity

- Differentiation (derivatives), which is good for...

# Continuity and discontinuity in political science research

Why we care about continuity

- Differentiation (derivatives), which is good for...
- Utility functions (game theory)

# Continuity and discontinuity in political science research

Why we care about continuity

- Differentiation (derivatives), which is good for...
- Utility functions (game theory)
- How some variable affects another variable (statistical models)

# Continuity and discontinuity in political science research

Why we care about continuity

- Differentiation (derivatives), which is good for...
- Utility functions (game theory)
- How some variable affects another variable (statistical models)
- More on Wednesday

# Continuity and Discontinuity

Why we care about discontinuity: *regression discontinuity*

# Continuity and Discontinuity

Why we care about discontinuity: *regression discontinuity*

- Causal inference method: units are "treated" if above/below a threshold

# Continuity and Discontinuity


Why we care about discontinuity: *regression discontinuity*

- Causal inference method: units are "treated" if above/below a threshold
- Popular example: vote share to wins and losses, wins and losses affect future fundraising

# Continuity and Discontinuity

Why we care about discontinuity: *regression discontinuity*



- Causal inference method: units are "treated" if above/below a threshold
- Popular example: vote share to wins and losses, wins and losses affect future fundraising



# Continuity and Discontinuity

Why we care about discontinuity: *regression discontinuity*


- Causal inference method: units are "treated" if above/below a threshold
- Popular example: vote share to wins and losses, wins and losses affect future fundraising



# Monotonicity

A function is *monotonic* if it always increases (monotonically increasing) or always decreases (monotonically decreasing)


- increasing: for any  $x_1 > x_2$ ,  $f(x_1) > f(x_2)$
- decreasing: for any  $x_1 < x_2$ ,  $f(x_1) < f(x_2)$



# Concavity and Convexity

Imagine you draw a line between two points along a function. A function (or segment of a function) is *concave* if this line is below the function, and *convex* if the line is above the function.

- concave:  $\frac{f(x_1)+f(x_2)}{2} < f\left(\frac{x_1+x_2}{2}\right)$
- convex:  $\frac{f(x_1)+f(x_2)}{2} > f\left(\frac{x_1+x_2}{2}\right)$




# Invertible functions

A function maps an input to an output. A function is *invertible* if there exists a reverse function that maps the output back to the input.

Formally: if  $y = f(x)$ , then  $f^{-1}(y) = x$


Also:  $f^{-1}(f(x)) = x$



# Exponents

The *exponent* operator multiplies a number by itself the number of times indicated in the exponent


$$x^n = x * x * \dots * x \quad (n \text{ times})$$



# Roots

*Root* operators return the number that, when multiplied by itself the number of times indicated in the root, is equal to the input. When no number is given, that indicates the *square root*.

$$x = \sqrt[n]{x} * \sqrt[n]{x} \dots \sqrt[n]{x} \quad (n \text{ times})$$



# Exponents and Roots

All roots can be expressed as exponents (with the same properties)

$$\sqrt[n]{x} \equiv x^{\frac{1}{n}}$$

Important properties for exponents and roots

---

Zeroth power

$$x^0 = 1$$

Negative powers

$$x^{-n} = \frac{1}{x^n}$$

Inversion using exponents

$$x^{-1} = \frac{1}{x}$$

Distribution of powers (multiplication)  $(x * y)^n = x^n * y^n$

Distribution of powers (division)

$$\left(\frac{x}{y}\right)^n = \frac{x^n}{y^n}$$

Product of powers

$$x^n * x^m = x^{n+m}$$

Nested powers

$$(x^n)^m = x^{n*m}$$

**Two important continuous, monotonic, invertible functions:  
Exponentials and Logarithms**

# Exponentials and Logarithms

The *logarithm* (log) of some value  $y$  (with base  $b$ ) is...


...the *power* to which that base would need to be raised to equal  $y$

$$\text{If } b^x = y, \text{ then } \log_b(y) = x$$

Exponentials and logarithms are inverse functions. Logs "undo" exponentials, and exponentials "undo" logs.

# Exponentials and Logarithms



We can see this because exponentials and logs are *reflections of each other over  $y = x$*  (one way to identify inverse functions)



# Exponents $\neq$ Exponentials

Exponent:  $x^2$  ( $x$  is the base)

Exponentials:  $2^x$  ( $x$  is the power)



# Better yet...


What happens when you exponentiate a parabola?

$$f(x) = e^{-x^2}$$

# Better yet...

What happens when you exponentiate a parabola?

$$f(x) = e^{-x^2}$$



# Rules for logarithms

The following apply to all logs, regardless of base

---

Log 1

$$\log(1) = 0$$

Log 0

undefined, approaches  $-\infty$

Multiplication

$$\log(x * y) = \log(x) + \log(y)$$

Division

$$\log\left(\frac{x}{y}\right) = \log(x) - \log(y)$$

Exponentiation

$$\log(x^a) = a * \log(x)$$

Basis

$$\log_b(b^x) = x$$


---

(potentially) helpful video for understanding logs [here](#)


You *WILL* use logs and exponents

They are important for *probability distributions*

$$f(x \mid \mu, \sigma) = \frac{1}{2\pi\sigma^2} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$



## Common transformations



House Candidates, 2012.  
Data: DIME (Bonica 2018)

**They are helpful for analytic manipulation of equations**

e.g. maximum likelihood estimation

## They are helpful for analytic manipulation of equations

e.g. maximum likelihood estimation

We flip a coin  $n = 5$  times and get  $y = 4$  heads. What's the probability of a heads ( $\pi$ )?

## They are helpful for analytic manipulation of equations

e.g. maximum likelihood estimation

We flip a coin  $n = 5$  times and get  $y = 4$  heads. What's the probability of a heads ( $\pi$ )?

Probability mass function for a binomial outcome (  $n$  independent trials,  $y$  successes, success probability  $\pi$ ):  $\Pr(y = k \mid \pi) = \binom{n}{k} \pi^k (1 - \pi)^{n-k}$

Plug in our data:  $\Pr(y = 4 \mid \pi) = \binom{5}{4} \pi^4 (1 - \pi)^{5-4}$

## They are helpful for analytic manipulation of equations

e.g. maximum likelihood estimation

We flip a coin  $n = 5$  times and get  $y = 4$  heads. What's the probability of  $\pi$ ?

Probability mass function for a binomial outcome ( $n$  independent trials,  $y$  successes, success probability  $\pi$ ):  $\Pr(y = k \mid \pi) = \binom{n}{k} \pi^k (1 - \pi)^{n-k}$

Plug in our data:  $\Pr(y = 4 \mid \pi) = \binom{5}{4} \pi^4 (1 - \pi)^{5-4}$

Log probability:  $\log(\Pr(y = 4 \mid \pi)) = \log \binom{5}{4} + 4 \log(\pi) + (5 - 4) \log(1 - \pi)$

## They are helpful for analytic manipulation of equations

e.g. maximum likelihood estimation

We flip a coin  $n = 5$  times and get  $y = 4$  heads. What's the probability of a heads ( $\pi$ )?

Probability mass function for a binomial outcome (  $n$  independent trials,  $y$  successes, success probability  $\pi$ ):  $\Pr(y = k \mid \pi) = \binom{n}{k} \pi^k (1 - \pi)^{n-k}$

Plug in our data:  $\Pr(y = 4 \mid \pi) = \binom{5}{4} \pi^4 (1 - \pi)^{5-4}$

Log probability:  $\log(\Pr(y = 4 \mid \pi)) = \log \binom{5}{4} + 4 \log(\pi) + (5 - 4) \log(1 - \pi)$

This function defines the (log) probability of our data, as a function of an unknown  $\pi$ .

## They are helpful for analytic manipulation of equations

e.g. maximum likelihood estimation

We flip a coin  $n = 5$  times and get  $y = 4$  heads. What's the probability of a heads ( $\pi$ )?

Probability mass function for a binomial outcome (  $n$  independent trials,  $y$  successes, success probability  $\pi$ ):  $\Pr(y = k \mid \pi) = \binom{n}{k} \pi^k (1 - \pi)^{n-k}$

Plug in our data:  $\Pr(y = 4 \mid \pi) = \binom{5}{4} \pi^4 (1 - \pi)^{5-4}$


Log probability:  $\log(\Pr(y = 4 \mid \pi)) = \log \binom{5}{4} + 4 \log(\pi) + (5 - 4) \log(1 - \pi)$

This function defines the (log) probability of our data, as a function of an unknown  $\pi$ .

If we maximize this function with respect to  $\pi$ , we find the  $\pi$  value that gives us the greatest probability. That is, the *most likely value of  $\pi$*  that could give us these data.

# This is how maximum likelihood works


Maximizing the log likelihood




(potentially) helpful video for  
understanding MLE [here](#)

# This is how maximum likelihood works

Maximizing the log likelihood




On the unlogged scale




(potentially) helpful video for  
understanding MLE [here](#)

# This is how maximum likelihood works

Maximizing the log likelihood



On the unlogged scale




(potentially) helpful video for  
understanding MLE [here](#)


Point being: we use logs in MLE

# This is how maximum likelihood works

Maximizing the log likelihood



On the unlogged scale



(potentially) helpful video for  
understanding MLE [here](#)

Point being: we use logs in MLE

(Note what the log transformation does to the  $y$ -axis)

**Logs get easier with experience, which you will have**

# Base $e$

Although many early examples with logs use some arbitrary base (like base 10, the "common log"), most applications use base  $e$  ( $\log_e$ , the "natural log",  $\ln$ )

- $e$  = Euler's number, approximately 2.718281828 ...

# Base $e$

Although many early examples with logs use some arbitrary base (like base 10, the "common log"), most applications use base  $e$  ( $\log_e$ , the "natural log",  $\ln$ )

- $e$  = Euler's number, approximately 2.718281828 ...
- $e^1 = e$
- $\ln(e) = 1$

# Base $e$

Although many early examples with logs use some arbitrary base (like base 10, the "common log"), most applications use base  $e$  ( $\log_e$ , the "natural log",  $\ln$ )

- $e$  = Euler's number, approximately 2.718281828 ...
- $e^1 = e$
- $\ln(e) = 1$
- $e^0 = 1$
- $\ln(1) = 0$

# Why base $e$ ?

It turns out that  $e$  is very important for functions that have constant and continuous growth rates (e.g. compounding interest)

# Why base $e$ ?

It turns out that  $e$  is very important for functions that have constant and continuous growth rates (e.g. compounding interest)

Similar properties in probability

# Why base $e$ ?

It turns out that  $e$  is very important for functions that have constant and continuous growth rates (e.g. compounding interest)

Similar properties in probability

If you roll an  $n$ -sided die  $n$  times, the probability of getting side  $n$  exactly zero times approaches  $\frac{1}{e}$  (as  $n$  increases)

# Why base $e$ ?

It turns out that  $e$  is very important for functions that have constant and continuous growth rates (e.g. compounding interest)

Similar properties in probability

If you roll an  $n$ -sided die  $n$  times, the probability of getting side  $n$  exactly zero times approaches  $\frac{1}{e}$  (as  $n$  increases)

We will emphasize the value of base  $e$  when we talk about probability (and the concept of *odds*)

# Some practice with logs and exponents

# Logs and Exponents

If  $f(x) = \log_{10} (x)$ , what is  $f(1000)$ ?

# Logs and Exponents

If  $f(x) = \log_{10} (x)$ , what is  $f(1000)$ ?

- $\log_{10} (1000) = 3$ , because  $10^3 = 1000$

# Logs and Exponents

If  $f(x) = \log_{10}(x)$ , what is  $f(1000)$ ?

- $\log_{10}(1000) = 3$ , because  $10^3 = 1000$

If  $g(x) = \ln\left(\frac{x}{e}\right)$ , what is  $g(4)$ ?

# Logs and Exponents

If  $f(x) = \log_{10}(x)$ , what is  $f(1000)$ ?

- $\log_{10}(1000) = 3$ , because  $10^3 = 1000$

If  $g(x) = \ln\left(\frac{x}{e}\right)$ , what is  $g(4)$ ?

$$\begin{aligned} \ln\left(\frac{4}{e}\right) &= \ln(4) - \ln(e) \\ \bullet & \\ &\approx 1.386 - 1 \\ &\approx 0.386 \end{aligned}$$

# Logs and Exponents

If  $f(x) = \log_{10}(x)$ , what is  $f(1000)$ ?

- $\log_{10}(1000) = 3$ , because  $10^3 = 1000$

If  $g(x) = \ln\left(\frac{x}{e}\right)$ , what is  $g(4)$ ?

$$\begin{aligned} \ln\left(\frac{4}{e}\right) &= \ln(4) - \ln(e) \\ \bullet & \\ &\approx 1.386 - 1 \\ &\approx 0.386 \end{aligned}$$

If  $h(x) = \log_2(x^5)$ , what is  $h(4)$ ?

# Logs and Exponents

If  $f(x) = \log_{10}(x)$ , what is  $f(1000)$ ?

- $\log_{10}(1000) = 3$ , because  $10^3 = 1000$

If  $g(x) = \ln\left(\frac{x}{e}\right)$ , what is  $g(4)$ ?

$$\begin{aligned} \ln\left(\frac{4}{e}\right) &= \ln(4) - \ln(e) \\ \bullet & \\ &\approx 1.386 - 1 \\ &\approx 0.386 \end{aligned}$$

If  $h(x) = \log_2(x^5)$ , what is  $h(4)$ ?

$$\begin{aligned} \log_2(4^5) &= 5 \log_2(4) \\ \bullet & \\ &= 5 * 2 \\ &= 10 \end{aligned}$$

# Solve:


- $\log_2(4^3)$
- $\ln\left(\frac{x}{y} * q^4 * e\right)$

**Now for something slightly easier**

# Absolute value

The absolute value operator returns the positive representation of a number

$$|x| = \begin{cases} x & \text{if } x \text{ is positive} \\ -x & \text{if } x \text{ is negative} \end{cases}$$



# Multiplying polynomials

Polynomial: an expression with variables and coefficients, using only addition, subtraction, multiplication, and non-negative integer exponents

$$x^3 + 5x^2 + 4x^2 + 7$$

Which are variables, which are coefficients?

# Multiplying polynomials

Polynomial: an expression with variables and coefficients, using only addition, subtraction, multiplication, and non-negative integer exponents

$$x^3 + 5x^2 + 4x^2 + 7$$

Which are variables, which are coefficients?

Multiply them by *distributing*: every element of each polynomial must be multiplied by every element of the other polynomial, then group terms by the powers of the variables.

Assume  $a$ ,  $b$ , and  $c$  are coefficients, what is  $ax(bx^2 + c)$  ?

# Multiplying polynomials

Polynomial: an expression with variables and coefficients, using only addition, subtraction, multiplication, and non-negative integer exponents

$$x^3 + 5x^2 + 4x^2 + 7$$

Which are variables, which are coefficients?

Multiply them by *distributing*: every element of each polynomial must be multiplied by every element of the other polynomial, then group terms by the powers of the variables.

Assume  $a$ ,  $b$ , and  $c$  are coefficients, what is  $ax(bx^2 + c)$  ?

$$ax(bx^2 + c) = (ax \cdot bx^2) + (ax \cdot c)$$

# Multiplying polynomials

Polynomial: an expression with variables and coefficients, using only addition, subtraction, multiplication, and non-negative integer exponents

$$x^3 + 5x^2 + 4x^2 + 7$$

Which are variables, which are coefficients?

Multiply them by *distributing*: every element of each polynomial must be multiplied by every element of the other polynomial, then group terms by the powers of the variables.

Assume  $a$ ,  $b$ , and  $c$  are coefficients, what is  $ax(bx^2 + c)$  ?

$$ax(bx^2 + c) = (ax \cdot bx^2) + (ax \cdot c)$$

$$ax(bx^2 + c) = abx^3 + acx$$

# Multiplying polynomials: FOIL

First, Outside, Inside, Last: for polynomials that each have two terms

$$(ax + b) \cdot (cx + d)$$

How do we FOIL?

# Multiplying polynomials: FOIL

First, Outside, Inside, Last: for polynomials that each have two terms

$$(ax + b) \cdot (cx + d)$$

How do we FOIL?

- First:  $ax \cdot cx$

# Multiplying polynomials: FOIL

First, Outside, Inside, Last: for polynomials that each have two terms

$$(ax + b) \cdot (cx + d)$$

How do we FOIL?

- First:  $ax \cdot cx$
- Outside:  $ax \cdot d$

# Multiplying polynomials: FOIL

First, Outside, Inside, Last: for polynomials that each have two terms

$$(ax + b) \cdot (cx + d)$$

How do we FOIL?

- First:  $ax \cdot cx$
- Outside:  $ax \cdot d$
- Inside:  $b \cdot cx$

# Multiplying polynomials: FOIL

First, Outside, Inside, Last: for polynomials that each have two terms

$$(ax + b) \cdot (cx + d)$$

How do we FOIL?

- First:  $ax \cdot cx$
- Outside:  $ax \cdot d$
- Inside:  $b \cdot cx$
- Last:  $b \cdot d$

# Multiplying polynomials: FOIL

First, Outside, Inside, Last: for polynomials that each have two terms

$$(ax + b) \cdot (cx + d)$$

How do we FOIL?

- First:  $ax \cdot cx$
- Outside:  $ax \cdot d$
- Inside:  $b \cdot cx$
- Last:  $b \cdot d$

What do we get?

# Multiplying polynomials: FOIL

First, Outside, Inside, Last: for polynomials that each have two terms

$$(ax + b) \cdot (cx + d)$$

How do we FOIL?

- First:  $ax \cdot cx$
- Outside:  $ax \cdot d$
- Inside:  $b \cdot cx$
- Last:  $b \cdot d$

What do we get?

$$acx^2 + adx + bcx + bd$$

# Longer polynomials

They work the same way, just keep track of all the terms. (FOIL is the same as distributing)

$$(2x^4 + 5x^3) \cdot (8x^2 + x + 3) = ?$$

# Longer polynomials

They work the same way, just keep track of all the terms. (FOIL is the same as distributing)

$$(2x^4 + 5x^3) \cdot (8x^2 + x + 3) = ?$$

$$= (2x^4 \cdot 8x^2) + (2x^4 \cdot x) + (2x^4 \cdot 3) + (5x^3 \cdot 8x^2) + (5x^3 \cdot x) + (5x^3 \cdot 3)$$

$$= 16x^6 + 2x^5 + 6x^4 + 40x^5 + 5x^4 + 15x^3$$

$$= 16x^6 + 42x^5 + 11x^4 + 15x^3$$

# Polynomial practice

Find the products:

- $(x^2 + 3) \cdot (x - 2)$
- $(3p + 4q) \cdot (p - 2q)$

# Factorials

The *factorial* operator (denoted with an exclamation mark !) returns the product of an integer with all lesser integers.

$$x! = x \cdot (x - 1) \cdot (x - 2) \cdot \dots \cdot 2 \cdot 1$$

# Factorials

The *factorial* operator (denoted with an exclamation mark !) returns the product of an integer with all lesser integers.

$$x! = x \cdot (x - 1) \cdot (x - 2) \cdot \dots \cdot 2 \cdot 1$$

An example:  $10! = 10 * 9 * 8 * 7 * \dots * 3 * 2 * 1 = 3628800$

# Factorials

The *factorial* operator (denoted with an exclamation mark !) returns the product of an integer with all lesser integers.

$$x! = x \cdot (x - 1) \cdot (x - 2) \cdot \dots \cdot 2 \cdot 1$$

An example:  $10! = 10 * 9 * 8 * 7 * \dots * 3 * 2 * 1 = 3628800$

Special factorials:

$$1! = 1$$

$$0! = 1$$

# Factorials

The *factorial* operator (denoted with an exclamation mark !) returns the product of an integer with all lesser integers.

$$x! = x \cdot (x - 1) \cdot (x - 2) \cdot \dots \cdot 2 \cdot 1$$

An example:  $10! = 10 * 9 * 8 * 7 * \dots * 3 * 2 * 1 = 3628800$

Special factorials:

$$1! = 1$$

$$0! = 1$$

These come in handy when we do probabilities (combinations and permutations)

# **Let's call it a day**

Homework is online

<https://github.com/shirikov/math-camp-2020>