

CIRCULAR $L(j, k)$ -LABELING NUMBERS OF SQUARE OF PATHS*

Qiong Wu^{1,†} and Wai Chee Shiu^{2,‡}

¹Faculty of Science, Tianjin University of Technology and Education,
Tianjin, China

²Department of Mathematics, Hong Kong Baptist University,
Kowloon Tong, Hong Kong, China

Received: September 12, 2016; Accepted: October 21, 2016

Abstract

Let j, k and σ be positive numbers, a circular σ - $L(j, k)$ -labeling of a graph G is a function $f : V(G) \rightarrow [0, \sigma)$ such that $|f(u) - f(v)|_\sigma \geq j$ if u and v are adjacent, and $|f(u) - f(v)|_\sigma \geq k$ if u and v are at distance two, where $|a - b|_\sigma = \min\{|a - b|, \sigma - |a - b|\}$. The minimum σ such that there exist a circular σ - $L(j, k)$ -labeling of G is called the circular- $L(j, k)$ -labeling number of G and is denoted by $\sigma_{j,k}(G)$. The k -th power G^k of an undirected graph G is a graph with the same set of vertices and an edge between two vertices when their distance in G is at most k . In this paper, the circular $L(j, k)$ -labeling numbers of P_n^2 are determined.

2000 Mathematics Subject Classification: Primary 05C78; Secondary 05C15

Keywords: Circular $L(j, k)$ -labeling, square of path, code assignment

1. Introduction

The rapid growth of computer wireless networks highlighted the scarcity of available codes (such as radio frequencies) for communication with minimum interference. For example, the Packet Radio Network (PRN) is a computer network that using radio frequencies to transmit packet among computers. The two major types of interference in PRN are direct collision (or interference), which caused by the transmission of adjacent stations (computers), and hidden terminal collision (or interference), which caused by distance-two stations that transmit to the same receiving station or receive the data from the same transmitting station.

*This work is supported by Tianjin Research Program of Application Foundation and Advanced Technology, Tianjin Municipal Science and Technology Commission (No. 14JCYBJC43100), National Natural Science Foundation of China (No. 11601391), Faculty Research Grant of Hong Kong Baptist University.

[†]E-mail address: wuqiong@tute.edu.cn.

[‡]E-mail address: wcshiu@math.hkbu.edu.hk (Corresponding author).

Bertossi and Bonuccelli [1] introduced an optimal code assignment to avoid hidden terminal interference and Jin and Yeh [10] further generalized it to the $L(j, k)$ -labeling problem with $j \leq k$.

Let G be a graph and let $V(G)$ and $E(G)$ be its vertex set and edge set, respectively. For any two vertices u and v , let $d_G(u, v)$ (or simply $d(u, v)$) denote the distance (length of a shortest path) between u and v in G . All notation not defined in this article can be found in the book [2].

For positive numbers j and k , an $L(j, k)$ -labeling f of G is an assignment of numbers to vertices of G such that $|f(u) - f(v)| \geq j$ if $uv \in E(G)$, and $|f(u) - f(v)| \geq k$ if $d(u, v) = 2$. The *span* of f is the difference between the maximum and the minimum numbers assigned by f . The $L(j, k)$ -labeling number of G , denoted by $\lambda_{j,k}(G)$, is the minimum span over all $L(j, k)$ -labeling of G .

The introduction of $L(j, k)$ -labeling with $j \geq k$ was motivated from the channel assignment problem [7]. Afterwards, Heuvel, Leese and Shepherd initiated the investigation of circular $L(j, k)$ -labeling of a graph in [8]. They used the “circular distance” (will be defined below) to instead of the “linear distance”.

For a positive real number σ , let $S(\sigma)$ denote the circle obtained from the closed interval $[0, \sigma]$ by identifying 0 and σ into a single point. $S(\sigma)$ is called a σ -cycle. For any $x \in \mathbb{R}$, $[x]_\sigma \in [0, \sigma)$ denotes the remainder of x upon division of σ . The *circular distance* of two points p and q on $S(\sigma)$ is defined as $|p - q|_\sigma = \min\{|p - q|, \sigma - |p - q|\}$.

A *circular σ - $L(j, k)$ -labeling* of G is a function $f : V(G) \rightarrow S(\sigma)$ satisfying $|f(x) - f(y)|_\sigma \geq j$ if $d(x, y) = 1$ and $|f(x) - f(y)|_\sigma \geq k$ if $d(x, y) = 2$. The minimum σ such that there exists a circular σ - $L(j, k)$ -labeling of G is called the *circular $L(j, k)$ -labeling number* of G and is denoted by $\sigma_{j,k}(G)$. $L(j, k)$ -labeling numbers or circular $L(j, k)$ -labeling numbers of graphs were studied in many articles, please refer to [4–6, 11, 12, 14–17, 19, 25] for $j \geq k$ and [1, 3, 10, 13, 20–24] for $j \leq k$.

2. Main Results

The k -th power G^k of an undirected graph G is a graph with the same set of vertices and an edge between two vertices when their distance in G is at most k . G^2 is called the *square* of G .

Lemma 2.1. *Let j and k be two positive numbers with $j \leq k$. Suppose G is a graph and H is an induced subgraph of G . Then $\sigma_{j,k}(G) \geq \sigma_{j,k}(H)$.*

Note that Lemma 2.1 is not true if H is not an induced subgraph. Suppose f is a circular σ - $L(j, k)$ -labeling of G . For any $a \in \mathbb{R}$, define $g(v) = [f(v) - a]_\sigma$ for $v \in V(G)$. Here g is still a circular σ - $L(j, k)$ -labeling of G . In this paper, the path is denoted by $P_n = v_0 v_1 \cdots v_{n-1}$. Without loss of generality, we may always assume v_0 is labeled by 0 when we consider circular $L(j, k)$ -labelings of P_n^2 . Also, in this paper we shall use $[a, b]$ to denote the close interval from a to b . Similar definition are applied for semi-open intervals $(a, b]$ and $[a, b)$.

Lemma 2.2. *Let $a, b \in \mathbb{R}$ and let σ, h and l be positive numbers. Suppose $|a - b| \geq h$ and $|a - b|_\sigma \geq l$. Then $\sigma \geq h + l$.*

Proof. The lemma follows from $l \leq |a-b|_\sigma = \min\{|a-b|, \sigma-|a-b|\} \leq \sigma-|a-b| \leq \sigma-h$. \square

Theorem 2.1. For $j \leq k$, $\sigma_{j,k}(P_4^2) = \max\{2k, 4j\}$.

Proof. Let $\eta = \max\{2k, 4j\}$. Define a circular labeling f for P_4^2 as follows:

If $k \geq 2j$, let $f(v_0) = 0$, $f(v_1) = j + k$, $f(v_2) = j$ and $f(v_3) = k$.

If $j \leq k < 2j$, let $f(v_0) = 0$, $f(v_1) = 3j$, $f(v_2) = j$ and $f(v_3) = 2j$.

It is clearly that f is a circular η - $L(j, k)$ -labeling of P_4^2 for $j \leq k$. Then $\sigma_{j,k}(P_4^2) \leq \max\{2k, 4j\}$.

On the other hand, let f be a circular σ - $L(j, k)$ -labeling for P_4^2 . Note that by our assumption, $f(v_0) = 0$. Since $d(v_0, v_3) = 2$, $f(v_3) \geq k$. By Lemma 2.2, this implies that $\sigma \geq 2k$. Moreover, since any two vertices of P_4^2 are adjacent or of distance two, $\max_{1 \leq i \leq 3} f(v_i) \geq 3j$. Since $|f(v_s) - f(v_t)|_\sigma \geq j$ and $\max\{|f(v_s) - f(v_t)|\} \geq 3j$ for $0 \leq s < t \leq 3$, by Lemma 2.2 we have $\sigma \geq 4j$. Thus, $\sigma \geq \max\{2k, 4j\}$.

Hence, $\sigma_{j,k}(P_4^2) = \max\{2k, 4j\}$. \square

Theorem 2.2. For $j \leq k$, $\sigma_{j,k}(P_5^2) = \max\{j + 2k, 5j\}$.

Proof. Let $\eta = \max\{j + 2k, 5j\}$. Given a circular labeling f for P_5^2 as follows:

If $k \geq 2j$, let $f(v_0) = 0$, $f(v_1) = 2j + k$, $f(v_2) = j$, $f(v_3) = j + k$ and $f(v_4) = k$.

If $j \leq k < 2j$, let $f(v_0) = 0$, $f(v_1) = 4j$, $f(v_2) = j$, $f(v_3) = 3j$ and $f(v_4) = 2j$.

It is easy to verify that f is a circular η - $L(j, k)$ -labeling of P_5^2 for $j \leq k$. Then $\sigma_{j,k}(P_5^2) \leq \max\{j + 2k, 5j\}$.

On the other hand, let f be a circular σ - $L(j, k)$ -labeling for P_5^2 . Since $d(v_0, v_3) = d(v_0, v_4) = 2$ and v_3 and v_4 are adjacent, $\max_{i=3,4} f(v_i) \geq j + k$. Again since $d(v_0, v_i) = 2$ for $i = 3, 4$, by Lemma 2.2 we have $\sigma \geq j + 2k$. Meantime, since any two vertices of P_5^2 are adjacent or of distance two, $\max_{1 \leq i \leq 4} f(v_i) \geq 4j$. Since $|f(v_s) - f(v_t)|_\sigma \geq j$ and $\max\{|f(v_s) - f(v_t)|\} \geq 4j$ for $0 \leq s < t \leq 4$, by Lemma 2.2 we have $\sigma \geq 5j$. Thus, $\sigma \geq \max\{j + 2k, 5j\}$.

Hence, $\sigma_{j,k}(P_5^2) = \max\{j + 2k, 5j\}$ for $j \leq k$. \square

Theorem 2.3. Let j and k be two positive numbers with $j \leq k < 3j$. For $n \geq 6$, $\sigma_{j,k}(P_n^2) = 4j + k$.

Proof. Let $\sigma = 4j + k$. Define a circular labeling f of P_n^2 as follows.

$f(v_i) = [ij]_\sigma$, for $0 \leq i \leq n - 1$. It is easy to verify that f is a circular σ - $L(j, k)$ -labeling of P_n^2 for $j \leq k < 3j$ and $n \geq 6$. Then $\sigma_{j,k}(P_n^2) \leq 4j + k$ for $j \leq k < 3j$ and $n \geq 6$.

On the other hand, since P_6^2 is an induced subgraph of P_n^2 for $n \geq 6$, by Lemma 2.1, we have $\sigma_{j,k}(P_n^2) \geq \sigma_{j,k}(P_6^2)$. It suffices to find a lower bound of $\sigma = \sigma_{j,k}(P_6^2)$.

Let f be a circular σ - $L(j, k)$ -labeling for P_6^2 . Note that $\sigma \leq 4j + k$. As mentioned before $f(v_0) = 0$. By considering the distances of v_i from v_0 , we have $f(v_1), f(v_2) \in [j, \sigma - j]$ and $f(v_3), f(v_4) \in [k, \sigma - k]$. Without loss of generality, we may assume that $f(v_1) < f(v_4)$, otherwise consider the new labeling $[\sigma - f]_\sigma$. Here we have $f(v_1) \in [j, \sigma - 2k]$ and $f(v_4) \in [2j, \sigma - k]$. Since $d(v_1, v_5) = 2$, we have $|f(v_1) - f(v_5)|_\sigma \geq k$, that is, $k \leq |f(v_1) - f(v_5)| \leq \sigma - k$. So we have $f(v_5) \in [j + k, \sigma)$ when $f(v_1) < f(v_5)$ or $f(v_5) \in [0, \sigma - 3k]$ when $f(v_1) > f(v_5)$.

Case 1. Suppose $2j < k < 3j$. Here $\sigma - 3k < 0$. Under this condition, by the same argument above, we also have $f(v_2) < f(v_5)$. Thus $f(v_5) \in [j + k, \sigma)$. This implies that $f(v_2) \in [j, \sigma - k)$. Since the length of $[j, \sigma - 2k]$ is $\sigma - j - 2k \leq 3j - k < j$, $f(v_2), f(v_3) \in [2j, \sigma - k]$. Now $f(v_2), f(v_3), f(v_4) \in [2j, \sigma - k]$. Since v_2, v_3, v_4 induce a K_3 , $\sigma - k - 2j \geq 2j$. Hence $\sigma \geq 4j + k$.

Case 2. Suppose $j \leq k \leq 2j$. We have two cases:

- a. Suppose $f(v_5) \in [0, \sigma - 3k]$. Similarly, we obtain that $f(v_1), f(v_2) \in [k, \sigma - j]$, and $f(v_3), f(v_4) \in [k, \sigma - k]$. We can conclude that $f(v_i) \in [k, \sigma - j]$ for $1 \leq i \leq 4$. Since the distances between any of two of v_1, v_2, v_3, v_4 are at most 2, it means that $\sigma - j - k \geq 3j$. That is, $\sigma \geq 4j + k$.
- b. Suppose $f(v_5) \in [j + k, \sigma)$.
 If $f(v_2) < f(v_5)$, then $f(v_2) \in [j, \sigma - k)$. We can conclude that $f(v_i) \in [j, \sigma - k]$ for $1 \leq i \leq 4$. Similar to the above case, we have $\sigma \geq 4j + k$.
 If $f(v_2) > f(v_5)$, then $f(v_2) \in [j + 2k, \sigma - j]$ and $f(v_5) \in [j + k, \sigma - j - k]$. Combining with $f(v_3), f(v_4) \in [k, \sigma - k]$, we can conclude that $f(v_i) \in [k, \sigma - j]$ for $2 \leq i \leq 5$. Similar to the above case, we have $\sigma \geq 4j + k$.

This completes the proof. \square

Theorem 2.4. Let j and k be two positive numbers with $k \geq 3j$. We have $\sigma_{j,k}(P_6^2) = j + 2k$.

Proof. Define a circular labeling f for P_n^2 as follows:

$f(v_0) = 0, f(v_1) = j, f(v_2) = 2j, f(v_3) = k, f(v_4) = j + k$ and $f(v_5) = 2j + k$.

It is clearly that f is a circular $(j+2k)$ - $L(j, k)$ -labeling for P_6^2 . Then $\sigma_{j,k}(P_6^2) \leq j + 2k$.

On the other hand, since P_5^2 is an induced subgraph of P_6^2 , by Lemma 2.1 and Theorem 2.2, we have $\sigma_{j,k}(P_6^2) \geq \sigma_{j,k}(P_5^2) \geq j + 2k$. Hence, the proof completes. \square

Theorem 2.5. Let j and k be two positive numbers with $k \geq 3j$. We have $\sigma_{j,k}(P_n^2) = \frac{7}{3}k$ for $n \geq 7$.

Proof. Let $\sigma = \frac{7}{3}k$. For $n \geq 7$, define a circular labeling f of P_n^2 by $f(v_i) = [ik/3]_\sigma$ for $0 \leq i \leq n - 1$. It is easy to verify that f is a circular $\frac{7}{3}k$ - $L(j, k)$ -labeling of P_n^2 . Thus, $\sigma_{j,k}(P_n^2) \leq \frac{7}{3}k$.

Since P_7^2 is an induced subgraph of P_n^2 for $n \geq 7$, by Lemma 2.1, we have $\sigma_{j,k}(P_n^2) \geq \sigma_{j,k}(P_7^2)$. It suffices to find a lower bound of $\sigma_{j,k}(P_7^2)$.

Let $\sigma = \sigma_{j,k}(P_7^2)$. Note that $\sigma \leq \frac{7}{3}k$. Let f be a circular σ - $L(j, k)$ -labeling of P_7^2 . We define the traveling distance from v_i to v_j , denoted by $\mathcal{L}(i, j)$, to be the length from $f(v_i)$ to $f(v_j)$ along the clockwise direction in the σ -cycle. In P_7^2 , if we travel the vertices from $v_0, v_4, v_1, v_5, v_2, v_6, v_3$ one by one, then the total traveling distance from v_0 to v_3 is $T = \mathcal{L}(0, 4) + \mathcal{L}(4, 1) + \mathcal{L}(1, 5) + \mathcal{L}(5, 2) + \mathcal{L}(2, 6) + \mathcal{L}(6, 3)$. As $k \leq \mathcal{L}(u, v) \leq \sigma - k$ for $d(u, v) = 2$, we have

$$6k \leq T \leq 6(\sigma - k). \quad (2.1)$$

Since $\sigma \leq \frac{7}{3}k$, we have $2 < \frac{6k}{\sigma} \leq \frac{T}{\sigma} \leq 6 - \frac{6k}{\sigma} < 4$. It implies that we arrive $f(v_3)$ from $f(v_0)$ after rotating the σ -cycle 2 or 3 times.

Suppose we arrive $f(v_3)$ from $f(v_0)$ after rotating the σ -cycle 2 times. Since v_3 and v_0 are at distance 2 and $f(v_0) = 0$, it follows that $|T - 2\sigma|_\sigma \geq k$, that is $2\sigma + k \leq T \leq 3\sigma - k$. From (2.1), we have $6k \leq 3\sigma - k$, and thus, $\sigma \geq \frac{7}{3}k$.

Suppose we arrive $f(v_3)$ from $f(v_0)$ after rotating the σ -cycle 3 times. Since v_3 and v_0 are at distance 2 and $f(v_0) = 0$, it follows that $|T - 3\sigma|_\sigma \geq k$, that is $3\sigma + k \leq T \leq 4\sigma - k$. From (2.1), we have $3\sigma + k \leq 6(\sigma - k)$, and hence $\sigma \geq \frac{7}{3}k$.

Then, we have $\sigma_{j,k}(P_n^2) \geq \sigma_{j,k}(P_7^2) = \sigma \geq \frac{7}{3}k$. The proof completes. \square

References

- [1] A.A. Bertossi and M.A. Bonuccelli, Code assignment for hidden terminal interference avoidance in multihop packet radio networks, *IEEE/ACM Transactions on Networking*, **3** (1995), 441–449.
- [2] J.A. Bondy, U.S.R. Murty, *Graph theory with applications*, New York, MacMillan, 1976.
- [3] T. Calamoneri, A. Pelc, and R. Petreschi, Labeling trees with a condition at distance two, *Discrete Math.*, **306** (2006), 1534–1539.
- [4] T. Calamoneri, The $L(h, k)$ -Labelling Problem: A Survey and Annotated Bibliography, *Comput. J.*, **49** (2009), 585–608.
- [5] J.R. Griggs and X.T. Jin, Recent Progress in Mathematics and Engineering on Optimal Graph Labellings with Distance Conditions, *J. Combin. Optim.*, **14** (2007), 249–257.
- [6] J.R. Griggs and X.T. Jin, Optimal channel assignments for lattices with conditions at distance two, In Proc. 5th Int. Workshop on algorithms for wireless Mobile, Ad Hoc and Sensor Networks (W-MAN '05), Denver, Colorado, April 4–8. IEEE Computer Society, 2005.
- [7] J.R. Griggs and R.K. Yeh, Labelling graphs with a condition at distance 2, *SIAM J. Discrete Math.* **5** (1992), 586–595.
- [8] J. van den Heuvel, R.A. Leese and M.A. Shepherd, Graph labeling and radio channel assignment, *J. Graph Theory*, **29** (1998), 263–283.
- [9] K. R. Jayasree, T. Nicholas, The minimal $L(1, 2)$ -labelings of generalized Petersen graphs, *Int. J. Engrg. Sci. Tech.*, **3** (2011), 318–328.
- [10] X.T. Jin and R.K. Yeh, Graph distance-dependent labeling related to code assignment in computer networks, *Naval Research Logistics*, **52** (2005), 159–164.
- [11] P.C.B. Lam, W. Lin, and J. Wu, $L(j, k)$ -labelings and circular $L(j, k)$ -labelings of products of complete graphs, *J. Combin. Optim.* **14** (2007), 219–227.

-
- [12] R.A. Leese and S.D. Noble, Cyclic labelings with constraints at two distances, *Electron. J. Combin.*, **11** (2004), #R16.
 - [13] W. Lin and Q. Wu, The $(1, k)$ -circular chromatic numbers of lattices, manuscript, 2009.
 - [14] D. Liu, Hamiltonicity and circular distance two labelings, *Discrete Math.*, **232** (2001), 163–169.
 - [15] D. Liu, Sizes of graphs with fixed orders and spans for circular distance two labeling, *Ars Combin.*, **67** (2003), 125–139.
 - [16] D. Liu and X. Zhu, Circular distance two labeling and circular chromatic number, *Ars Combin.*, **69** (2003), 177–183.
 - [17] D. Liu and X. Zhu, Circular distance two labeling and the λ -number for outerplanar graphs, *SIAM J. Discrete Math.*, **19** (2005), 281–293.
 - [18] G. Mohammad, Theorems and computations in circular colorings of graphs, *Ph.D. Thesis*, Simon Fraser University, 2007.
 - [19] B. Mohar, Circular colorings of edge weighted graphs, *J. Graph Theory*, **43** (2003), 107–116.
 - [20] Q. Niu, $L(j, k)$ -labeling of graph and edge span, *M.Phil. Thesis*, Southeast University, 2007.
 - [21] W.C. Shiu and Q. Wu, $L(j, k)$ -labeling number of direct product of path and cycle, *Acta Math. Sin.* **29** (2013), 1437–1448.
 - [22] Q. Wu and W. Lin, Circular $L(j, k)$ -labeling numbers of trees and products of graphs, *J. Southeast University*, **26** (2010), 142–145.
 - [23] Q. Wu, W.C. Shiu, and P.K. Sun, Circular $L(j, k)$ -labeling number of direct product of path and cycle, *J. Comb. Optim.* **27** (2014), 355–368.
 - [24] Q. Wu and W.C. Shiu, $L(j, k)$ -labeling numbers of square paths, preprint (2016).
 - [25] R.K. Yeh, A survey on labeling graphs with a condition at distance two, *Discrete Math.*, **306** (2006), 1217–1231.