



# Ring-magic Labeling of Graphs

Wai Chee SHIU

Department of Mathematics,  
Hong Kong Baptist University, Hong Kong

Richard M. LOW

Department of Mathematics,  
San José State University, San José, USA

June 22, 2007



# Outline

- ★ Background
- ★ Ring-magic
- ★ General Results
- ★ group-magic but not ring-magic
- ★  $\mathbb{Z}_3$ -ring-magic trees
- ★  $V_4$ -ring-magic trees
- ★ Further studies



# Group-magic Labeling of Graphs

Let  $G = (V, E)$  be a connected, simple graph. Let  $A$  be a finite abelian group.


**Definition 1** Suppose that there exists a labeling  $f : E \rightarrow A \setminus \{0\}$  such that the induced vertex labeling  $f^+ : V \rightarrow A$ , defined by

$$f^+(v) = \sum_{uv \in E} f(uv),$$

is a constant map. Then,  $f$  is an ***A-magic labeling*** of  $G$  and  $G$  is ***A-magic***.


# Some Examples

Ex 1.  $C_6$  is  $A$ -magic, for all abelian groups  $A$ .




# Some Examples

Ex 1.  $C_6$  is  $A$ -magic, for all abelian groups  $A$ .




Ex 2.  $P_3$  is not  $A$ -magic, for all abelian groups  $A$ .



# Some Examples (continued)

Ex 3. The following graph is  $\mathbb{Z}_3$ -magic, but not  $\mathbb{Z}_2$ -magic.



A graph is  $\mathbb{Z}_2$ -magic if and only if the degree of each vertex is of the same parity.



# $\mathbb{Z}_n$ -magic Labelings of Graphs

Various classes of graphs have already been studied, either partially or completely. They include:

- ★ Powers of paths (S-M. Lee, F. Saba, G.C. Sun, 1993).



# $\mathbb{Z}_n$ -magic Labelings of Graphs

Various classes of graphs have already been studied, either partially or completely. They include:

- ★ Powers of paths (S-M. Lee, F. Saba, G.C. Sun, 1993).
- ★ Amalgamations of stars and cycles (S-M. Lee, E. Salehi, 2003).





# $\mathbb{Z}_n$ -magic Labelings of Graphs

Various classes of graphs have already been studied, either partially or completely. They include:

- ★ Powers of paths (S-M. Lee, F. Saba, G.C. Sun, 1993).
- ★ Amalgamations of stars and cycles (S-M. Lee, E. Salehi, 2003).
- ★ Trees of diameter at most 4 (S-M. Lee, E. Salehi, H. Sun, 2004).



# $\mathbb{Z}_n$ -magic Labelings of Graphs

Various classes of graphs have already been studied, either partially or completely. They include:

- ★ Powers of paths (S-M. Lee, F. Saba, G.C. Sun, 1993).
- ★ Amalgamations of stars and cycles (S-M. Lee, E. Salehi, 2003).
- ★ Trees of diameter at most 4 (S-M. Lee, E. Salehi, H. Sun, 2004).
- ★ Eulerian graphs (R.M. Low, S-M. Lee, 2004).



# $\mathbb{Z}_n$ -magic Labelings of Graphs

Various classes of graphs have already been studied, either partially or completely. They include:

- ★ Powers of paths (S-M. Lee, F. Saba, G.C. Sun, 1993).
- ★ Amalgamations of stars and cycles (S-M. Lee, E. Salehi, 2003).
- ★ Trees of diameter at most 4 (S-M. Lee, E. Salehi, H. Sun, 2004).
- ★ Eulerian graphs (R.M. Low, S-M. Lee, 2004).
- ★ Maximal planar and maximal outerplanar graphs (S-M. Lee, Y.S. Ho, R.M. Low, 2004).



# $\mathbb{Z}_n$ -magic Labelings of Graphs

Various classes of graphs have already been studied, either partially or completely. They include:

- ★ Powers of paths (S-M. Lee, F. Saba, G.C. Sun, 1993).
- ★ Amalgamations of stars and cycles (S-M. Lee, E. Salehi, 2003).
- ★ Trees of diameter at most 4 (S-M. Lee, E. Salehi, H. Sun, 2004).
- ★ Eulerian graphs (R.M. Low, S-M. Lee, 2004).
- ★ Maximal planar and maximal outerplanar graphs (S-M. Lee, Y.S. Ho, R.M. Low, 2004).
- ★ Products of group-magic graphs (R.M. Low, S-M. Lee, 2006).



# $\mathbb{Z}_n$ -magic Labelings of Graphs

Various classes of graphs have already been studied, either partially or completely. They include:

- ★ Powers of paths (S-M. Lee, F. Saba, G.C. Sun, 1993).
- ★ Amalgamations of stars and cycles (S-M. Lee, E. Salehi, 2003).
- ★ Trees of diameter at most 4 (S-M. Lee, E. Salehi, H. Sun, 2004).
- ★ Eulerian graphs (R.M. Low, S-M. Lee, 2004).
- ★ Maximal planar and maximal outerplanar graphs (S-M. Lee, Y.S. Ho, R.M. Low, 2004).
- ★ Products of group-magic graphs (R.M. Low, S-M. Lee, 2006).
- ★  $N$ -partite graphs (W.C. Shiu, R.M. Low, 2006).



# $\mathbb{Z}_n$ -magic Labelings of Graphs

Various classes of graphs have already been studied, either partially or completely. They include:

- ★ Powers of paths (S-M. Lee, F. Saba, G.C. Sun, 1993).
- ★ Amalgamations of stars and cycles (S-M. Lee, E. Salehi, 2003).
- ★ Trees of diameter at most 4 (S-M. Lee, E. Salehi, H. Sun, 2004).
- ★ Eulerian graphs (R.M. Low, S-M. Lee, 2004).
- ★ Maximal planar and maximal outerplanar graphs (S-M. Lee, Y.S. Ho, R.M. Low, 2004).
- ★ Products of group-magic graphs (R.M. Low, S-M. Lee, 2006).
- ★  $N$ -partite graphs (W.C. Shiu, R.M. Low, 2006).
- ★ Sun graphs (W.C. Shiu, R.M. Low, 2007).

# Ring-magic Labelings of Graphs


Let  $G = (V, E)$  be a connected, simple graph. Let  $R$  be a commutative ring with unity 1.

**Definition 2** Suppose that there exists a labeling  $f : E \rightarrow R \setminus \{0\}$  such that the induced vertex labelings  $f^+ : V \rightarrow R$ , defined by  $f^+(v) = \sum_{uv \in E} f(uv)$ , and  $f^\times : V \rightarrow R$ , defined by  $f^\times(v) = \prod_{uv \in E} f(uv)$ , are constant maps. Then,  $f$  is an  ***$R$ -ring-magic labeling*** of  $G$  and  $G$  is  ***$R$ -ring-magic***.

In this case, the values of  $f^+$  and  $f^\times$  are called the ***additive*** and ***multiplicative  $R$ -magic values*** of  $f$ , respectively.

# Examples of $\mathbb{Z}_3$ -ring-Magic Graphs


Ex 1. A  $\mathbb{Z}_3$ -ring-magic tree.






# Examples of $\mathbb{Z}_3$ -ring-Magic Graphs

Ex 1. A  $\mathbb{Z}_3$ -ring-magic tree.



Ex 2. A  $\mathbb{Z}_3$ -ring-magic graph.





# Some Quick Observations

- ★ A regular graph is  $R$ -ring magic, for any ring  $R$ .



# Some Quick Observations

- ★ A regular graph is  $R$ -ring magic, for any ring  $R$ .
- ★ Let  $A$  be the abelian group associated with ring  $R$ . If  $G$  is not  $A$ -magic, then  $G$  is not  $R$ -ring magic.



# Some Quick Observations

- ★ A regular graph is  $R$ -ring magic, for any ring  $R$ .
- ★ Let  $A$  be the abelian group associated with ring  $R$ . If  $G$  is not  $A$ -magic, then  $G$  is not  $R$ -ring magic.
- ★  $G$  is  $\mathbb{Z}_2$ -ring magic  $\iff$  the degree of each vertex of  $G$  is of the same parity.



# A Few General Results

**Theorem 1** *Let  $R$  be a ring and  $G = (V, E)$  be an  $R$ -ring magic graph of order  $p$ . Let  $h$  and  $k$  be the additive and multiplicative  $R$ -magic values respectively, of an  $R$ -ring magic labeling  $f$ . Then,*

$$hp = 2a \text{ and } k^p = b^2$$

*for some  $a, b \in R$ .*

# A Few General Results

**Theorem 1** *Let  $R$  be a ring and  $G = (V, E)$  be an  $R$ -ring magic graph of order  $p$ . Let  $h$  and  $k$  be the additive and multiplicative  $R$ -magic values respectively, of an  $R$ -ring magic labeling  $f$ . Then,*

$$hp = 2a \text{ and } k^p = b^2$$

*for some  $a, b \in R$ .*

**Proof:**

$$hp = \sum_{v \in V} f^+(v) = \sum_{v \in V} \sum_{uv \in E} f(uv) = 2 \sum_{e \in E} f(e)$$

$$k^p = \prod_{v \in V} f^\times(v) = \prod_{v \in V} \prod_{uv \in E} f(uv) = \left( \prod_{e \in E} f(e) \right)^2$$




# A Few General Results (continued)

**Theorem 2** *Let  $R_1$  be a ring, which contains a subring isomorphic to ring  $R_2$ . If graph  $G$  is  $R_2$ -ring magic, then  $G$  is  $R_1$ -ring magic.*

# A Few General Results (continued)

**Theorem 2** *Let  $R_1$  be a ring, which contains a subring isomorphic to ring  $R_2$ . If graph  $G$  is  $R_2$ -ring magic, then  $G$  is  $R_1$ -ring magic.*

The converse of Theorem 2 does not always hold.



This graph is  $\mathbb{Z}_2$ -ring-magic. Hence it is  $\mathbb{Z}_6$ -ring-magic. However, it is straight-forward to show (using an exhaustive case analysis) that  $G$  is not  $\mathbb{Z}_3$ -group-magic and hence, cannot be  $\mathbb{Z}_3$ -ring-magic.



# A Few General Results (continued)

For a commutative ring  $R$ , let  $U(R)$  denote the group of units.

**Theorem 3** *Suppose that  $f$  is an  $R$ -ring magic labeling of a graph  $G$  and  $u \in U(R)$ , where  $R$  is an integral domain. Then,  $uf$  is an  $R$ -ring magic labeling of  $G \iff o(u) \mid [d(v_i) - d(v_j)]$ , for all  $v_i, v_j \in V(G)$ , where  $o(u)$  is the order of  $u$  in  $U(R)$ .*

# Proof of Theorem 3

**Proof:**  $(uf)^\times(v) = u^{d(v)}k$ , where  $k$  is the multiplicative magic value.

$$u^{d(v_i)}k = u^{d(v_j)}k \iff (u^{d(v_i)} - u^{d(v_j)})k = 0$$

$$\iff u^{d(v_i)} - u^{d(v_j)} = 0$$

$$\iff u^{d(v_i)-d(v_j)} = 1$$

$$\iff o(u) \mid [d(v_i) - d(v_j)].$$


# Example

Ex. Consider the ring  $\mathbb{Z}_5$ . Then,  $U(\mathbb{Z}_5) = \{1, 2, 3, 4\}$  is isomorphic (as a group) to  $\mathbb{Z}_4$ . Element in  $U(\mathbb{Z}_5)$  is of order 1 or 2.

# Example

Ex. Consider the ring  $\mathbb{Z}_5$ . Then,  $U(\mathbb{Z}_5) = \{1, 2, 3, 4\}$  is isomorphic (as a group) to  $\mathbb{Z}_4$ . Element in  $U(\mathbb{Z}_5)$  is of order 1 or 2.



In particular,  $u = 4$  which has order 2.



# Example

Ex. Consider the ring  $\mathbb{Z}_5$ . Then,  $U(\mathbb{Z}_5) = \{1, 2, 3, 4\}$  is isomorphic (as a group) to  $\mathbb{Z}_4$ . Element in  $U(\mathbb{Z}_5)$  is of order 1 or 2.

In particular,  $u = 4$  which has order 2.





# $\mathbb{Z}_n$ -ring Magic Graphs

**Corollary 4** *Let  $G$  be a  $\mathbb{Z}_n$ -ring magic graph of odd order. Then, for each odd prime factor  $r$  of  $n$ , the multiplicative magic value  $k$  must be a square in  $\mathbb{Z}_r$ .*



# $\mathbb{Z}_n$ -ring Magic Graphs

**Corollary 4** *Let  $G$  be a  $\mathbb{Z}_n$ -ring magic graph of odd order. Then, for each odd prime factor  $r$  of  $n$ , the multiplicative magic value  $k$  must be a square in  $\mathbb{Z}_r$ .*

**Proof:** By Theorem 1, there exists  $b \in \mathbb{Z}_n$  such that  $\left(\frac{k}{r}\right) = \left(\frac{k}{r}\right)^p = \left(\frac{k^p}{r}\right) = \left(\frac{b^2}{r}\right) = \left(\frac{b}{r}\right)^2$ . Here  $\left(\frac{a}{r}\right)$  is the Legendre symbol.



# A Natural Question

At this point, it is very natural for us to ask the following question. For a given  $n$ , are there graphs which are  $\mathbb{Z}_n$ -group-magic but not  $\mathbb{Z}_n$ -ring-magic?





# A Natural Question

At this point, it is very natural for us to ask the following question. For a given  $n$ , are there graphs which are  $\mathbb{Z}_n$ -group-magic but not  $\mathbb{Z}_n$ -ring-magic?

The answer to this question is yes for  $n$  being odd prime.



# A Natural Question

At this point, it is very natural for us to ask the following question. For a given  $n$ , are there graphs which are  $\mathbb{Z}_n$ -group-magic but not  $\mathbb{Z}_n$ -ring-magic?

The answer to this question is yes for  $n$  being odd prime. Let  $C_m(y)$  denote the  $m$ -cycle with  $y$  pendants attached to each vertex.



# A Natural Question

At this point, it is very natural for us to ask the following question. For a given  $n$ , are there graphs which are  $\mathbb{Z}_n$ -group-magic but not  $\mathbb{Z}_n$ -ring-magic?

The answer to this question is yes for  $n$  being odd prime. Let  $C_m(y)$  denote the  $m$ -cycle with  $y$  pendants attached to each vertex.

**Theorem 5** *Let  $n$  be an odd prime. Then, there exists an integer  $y$  such that  $C_4(y)$  is  $\mathbb{Z}_n$ -group magic but which is not  $\mathbb{Z}_n$ -ring magic.*

# An Example

Before to prove Theorem 5, we need a technical lemma.

**Lemma 6** *Let  $n$  be an odd prime. Then, there exists  $y \geq 1$  such that  $y \equiv -1 \pmod{n-1}$  and*

$$\left( \frac{y^2 - 2y - 3}{n} \right) = -1.$$

# An Example

Before to prove Theorem 5, we need a technical lemma.

**Lemma 6** *Let  $n$  be an odd prime. Then, there exists  $y \geq 1$  such that  $y \equiv -1 \pmod{n-1}$  and*  
$$\left( \frac{y^2 - 2y - 3}{n} \right) = -1.$$

We also need the following well-known result:

**Theorem A** *Let  $n$  be an odd prime. Then, there are exactly  $(n-1)/2$  quadratic residues of  $n$  and  $(n-1)/2$  quadratic nonresidues of  $n$  among the integers  $1, 2, \dots, n-1$ .*

# An Example

Before to prove Theorem 5, we need a technical lemma.

**Lemma 6** *Let  $n$  be an odd prime. Then, there exists  $y \geq 1$  such that  $y \equiv -1 \pmod{n-1}$  and  $\left(\frac{y^2-2y-3}{n}\right) = -1$ .*

**Proof:**  $y \equiv -1 \pmod{n-1} \Leftrightarrow y = -1 + k(n-1)$ .  
 $\left(\frac{y^2-2y-3}{n}\right) = \left(\frac{(-1-k)^2-2(-1-k)-3}{n}\right) = \left(\frac{k(k+4)}{n}\right)$ .

Theorem A implies the existence of an integer  $w$  ( $1 \leq w \leq n-1$ ) such that  $\left(\frac{w}{n}\right) = 1$  and  $\left(\frac{w+1}{n}\right) = -1$ .

Thus,  $\left(\frac{4w}{n}\right) \left(\frac{4(w+1)}{n}\right) = -1$ . Hence,  $\left(\frac{k(k+4)}{n}\right) = -1$ ,  
where  $k = 4w$ .



# Proof of Theorem 5

Consider the graph  $C_4(y)$ , where  $y$  satisfies the two conditions in Lemma 6.



# Proof of Theorem 5

Consider the graph  $C_4(y)$ , where  $y$  satisfies the two conditions in Lemma 6.

$C_4(y)$  is  $\mathbb{Z}_n$ -group-magic, as all of the pendants can be labeled with 1 and the edges in the cycle labeled  $a$ ,  $1 - a - y$ ,  $a$ , and  $1 - a - y$  respectively.





# Proof of Theorem 5

We now claim that  $C_4(y)$  is not  $\mathbb{Z}_n$ -ring-magic.  
Assume that  $C_4(y)$  has a  $\mathbb{Z}_n$ -ring-magic labeling.



# Proof of Theorem 5

We now claim that  $C_4(y)$  is not  $\mathbb{Z}_n$ -ring-magic.  
Assume that  $C_4(y)$  has a  $\mathbb{Z}_n$ -ring-magic labeling.

Degree of each vertex of  $C_4(y)$  is either 1 or  $y + 2$ .  
Because of Theorem 3 and the first condition in Lemma 6, we can assume without loss of generality that the pendants are labeled 1.

# Proof of Theorem 5

We now claim that  $C_4(y)$  is not  $\mathbb{Z}_n$ -ring-magic.  
Assume that  $C_4(y)$  has a  $\mathbb{Z}_n$ -ring-magic labeling.

Degree of each vertex of  $C_4(y)$  is either 1 or  $y + 2$ .  
Because of Theorem 3 and the first condition in Lemma 6, we can assume without loss of generality that the pendants are labeled 1.

Let  $a$  and  $b$  be the labels of the two non-pendant edges incident to a vertex of degree  $y + 2$  in  $C_4(y)$ . Then, the following two relationships must hold:

- ★  $a + b + y \equiv 1 \pmod{n}$ .
- ★  $ab \equiv 1 \pmod{n}$ .



# Proof of Theorem 5

Equivalent to saying that  $x(1 - y - x) \equiv 1 \pmod{n}$  has a solution, which implies that  $x^2 + (y - 1)x + 1 \equiv 0 \pmod{n}$  has a solution.

# Proof of Theorem 5

Equivalent to saying that  $x(1 - y - x) \equiv 1 \pmod{n}$  has a solution, which implies that  $x^2 + (y - 1)x + 1 \equiv 0 \pmod{n}$  has a solution.

However, this equation has discriminant  $D = (y - 1)^2 - 4(1) = y^2 - 2y - 3$ . Since  $y$  was chosen as to satisfy the conditions of Lemma 6,  $D$  is not a square in  $\mathbb{Z}_n$ .

# Proof of Theorem 5


Equivalent to saying that  $x(1 - y - x) \equiv 1 \pmod{n}$  has a solution, which implies that  $x^2 + (y - 1)x + 1 \equiv 0 \pmod{n}$  has a solution.

However, this equation has discriminant  $D = (y - 1)^2 - 4(1) = y^2 - 2y - 3$ . Since  $y$  was chosen as to satisfy the conditions of Lemma 6,  $D$  is not a square in  $\mathbb{Z}_n$ .

Thus,  $x^2 + (y - 1)x + 1 \equiv 0$  has no solutions in  $\mathbb{Z}_n$ . A contradiction.

# Actual Example

Ex. Let  $n = 7$ .




The system  $a + b = 3 \pmod{7}$ ,  $ab = 1 \pmod{7}$  has no solutions.



# $R$ -ring-Magic Property for Trees

Let  $T$  be a tree. If  $f$  is an  $R$ -ring-magic labeling of  $T$ , then the additive and multiplicative magic values of  $f$  are the same. We call this value the  *$R$ -ring-magic value* of  $T$ .





# Remark on Group-magic Labeling of Graphs

M. Doob studied group-magic graphs since 1974. The codomain of the edge labeling is the whole abelian group. That is, 0 was allowed to label on edges.

Under this concept of group-magic, R.P. Stanley considered (1973-1976)  $\mathbb{Z}$ -magic graphs. He pointed out that the theory of magic labelings could be studies in the general context of linear homogeneous diophantine equations.

# $R$ -ring-Magic Property for Trees

**Definition 3** Let  $R$  be a commutative ring with unity. A graph  $G = (V, E)$  is called  *$R'$ -ring-magic* if there exists a labeling  $f : E \rightarrow R$  such that the induced vertex labelings  $f^+ : V \rightarrow R$ , defined by  $f^+(v) = \sum_{uv \in E} f(uv)$ , and  $f^\times : V \rightarrow R$ , defined by  $f^\times(v) = \prod_{uv \in E} f(uv)$ , are constant maps.



# *R*-ring-Magic Property for Trees

**Theorem B (Shiu, Lam and Lee 2002)** *Let  $T$  be a tree and  $A$  be a ring. Suppose that  $f$  is an  $A'$ -group-magic labeling of  $T$ . If there is an edge  $e$  which is incident to a leaf of  $T$  and  $f(e) = 0$ , then  $f = 0$ .*

**Lemma 7** *Let  $T$  be a tree and  $A$  be a ring. Then,  $T$  has at most one  $A$ -ring-magic labeling with  $A$ -ring-magic value  $k$ .*

**Lemma 8** *Let  $T$  be a tree. Suppose that  $f$  is a  $\mathbb{Z}_n$ -ring-magic labeling of  $T$ , with  $\mathbb{Z}_n$ -ring-magic value 1. Then,  $f = 1$  (i.e., all the values of  $f$  are 1).*



# $\mathbb{Z}_3$ -ring Magic Property for Trees

**Theorem 9** *Let  $T$  be a tree. Then,  $T$  is  $\mathbb{Z}_3$ -ring magic with ring-magic value 1  $\iff d(v) \equiv 1 \pmod{3}$ , for all  $v \in V(T)$ . Moreover, for this case, suppose  $T$  is of order  $p$ . Then,  $p \equiv 2 \pmod{3}$ .*

**Proof:** The first part is easy.



# $\mathbb{Z}_3$ -ring Magic Property for Trees

**Theorem 9** *Let  $T$  be a tree. Then,  $T$  is  $\mathbb{Z}_3$ -ring magic with ring-magic value 1  $\iff d(v) \equiv 1 \pmod{3}$ , for all  $v \in V(T)$ . Moreover, for this case, suppose  $T$  is of order  $p$ . Then,  $p \equiv 2 \pmod{3}$ .*

**Proof:**

Let  $f$  be the  $\mathbb{Z}_3$ -ring-magic labeling with  $\mathbb{Z}_3$ -ring-magic value 1. By the proof of Theorem 1, we have

$$p = \sum_{v \in V} f^+(v) = 2(p - 1) \pmod{3}.$$

Hence,  $p \equiv 2 \pmod{3}$ .



# $\mathbb{Z}_3$ -ring Magic Property for Trees

**Corollary 10** *A tree  $T$  of odd order  $p$  is  $\mathbb{Z}_3$ -ring magic  $\iff p \equiv 5 \pmod{6}$  and  $d(v) \equiv 1 \pmod{3}$ , for all  $v \in V(T)$ .*



# $\mathbb{Z}_3$ -ring Magic Property for Trees

**Theorem 11** *Suppose a tree  $T$  has a  $\mathbb{Z}_3$ -ring magic labeling  $f$  with magic value 2. Let  $v$  be a vertex of  $T$  which is adjacent to pendants. Then,  $T$  is of even order and  $d(v) \equiv 1$  or  $0 \pmod{6}$ .*



# $\mathbb{Z}_3$ -ring Magic Property for Trees

**Theorem 11** *Suppose a tree  $T$  has a  $\mathbb{Z}_3$ -ring magic labeling  $f$  with magic value 2. Let  $v$  be a vertex of  $T$  which is adjacent to pendants. Then,  $T$  is of even order and  $d(v) \equiv 1 \text{ or } 0 \pmod{6}$ .*

**Proof:** Order of  $T$  must be even.

Let  $a$  and  $b$  be the number of 1 and 2 labeled to the edges incident with  $v$ .



# $\mathbb{Z}_3$ -ring Magic Property for Trees

**Theorem 11** *Suppose a tree  $T$  has a  $\mathbb{Z}_3$ -ring magic labeling  $f$  with magic value 2. Let  $v$  be a vertex of  $T$  which is adjacent to pendants. Then,  $T$  is of even order and  $d(v) \equiv 1$  or  $0 \pmod{6}$ .*

**Proof:** Let  $a$  and  $b$  be the number of 1 and 2 labeled to the edges incident with  $v$ .

Clearly  $a + b = d(v) = d$ . Since the ring-magic value is 2, we have  $2^b \equiv 2 \pmod{3}$  and  $a + 2b \equiv 2 \pmod{3}$ . Hence  $b$  is odd. Since  $f(uv) = 2$  for all pendants  $u$  adjacent with  $v$ , this implies that  $b = d$  or  $b = d - 1$ .

# $\mathbb{Z}_3$ -ring Magic Property for Trees

**Theorem 11** *Suppose a tree  $T$  has a  $\mathbb{Z}_3$ -ring magic labeling  $f$  with magic value 2. Let  $v$  be a vertex of  $T$  which is adjacent to pendants. Then,  $T$  is of even order and  $d(v) \equiv 1$  or  $0 \pmod{6}$ .*

**Proof:** Let  $a$  and  $b$  be the number of 1 and 2 labeled to the edges incident with  $v$ .

$$a + 2b \equiv 2 \pmod{3}, b = d \text{ or } b = d - 1.$$

(Case 1). If  $b = d$ , then  $a = 0$  and  $b \equiv 1 \pmod{3}$ .

Since  $b$  is odd,  $d = b \equiv 1 \pmod{6}$ .

(Case 2). If  $b = d - 1$ , then  $a = 1$  and  $b \equiv 2$

$\pmod{3}$ . Since  $b$  is odd,  $d = b + 1 \equiv 0 \pmod{6}$ .



# $\mathbb{Z}_3$ -ring Magic Property for Trees

The converse of Theorem 11 is not true in general.



# $\mathbb{Z}_3$ -ring Magic Property for Trees


The converse of Theorem 11 is not true in general.  
For example, the converse holds in the case where  
 $H = K_{1,7}$ .

# $\mathbb{Z}_3$ -ring Magic Property for Trees

The converse of Theorem 11 is not true in general. For example, the converse holds in the case where  $H = K_{1,7}$ .

However, the following caterpillar  $G$  provides a counter-example which illustrates that the converse does not always hold.

$G$  is of order 18,  $d(v_i) \equiv 1$  or  $0 \pmod{6}$ , but  $G$  is not  $\mathbb{Z}_3$ -group-magic. Hence, it is not  $\mathbb{Z}_3$ -ring-magic.



This caterpillar graph  $G$  is not  $\mathbb{Z}_3$ -ring-magic.



# $V_4$ -ring Magic Property for Trees

Let  $V_4$  be the ring  $\mathbb{Z}_2 \times \mathbb{Z}_2$ .

**Lemma 12** (Lee, Saba, Salehi and Sun, 2002). *A tree  $T$  is  $V_4$ -group magic  $\iff T$  has no vertex of even degree.*



# $V_4$ -ring Magic Property for Trees

Let  $V_4$  be the ring  $\mathbb{Z}_2 \times \mathbb{Z}_2$ .

**Lemma 12** (Lee, Saba, Salehi and Sun, 2002). *A tree  $T$  is  $V_4$ -group magic  $\iff T$  has no vertex of even degree.*

**Lemma 13** *Let  $T$  be a tree of odd order. Then,  $T$  is not  $V_4$ -ring magic.*



# $V_4$ -ring Magic Property for Trees

Let  $V_4$  be the ring  $\mathbb{Z}_2 \times \mathbb{Z}_2$ .

**Lemma 12** (Lee, Saba, Salehi and Sun, 2002). *A tree  $T$  is  $V_4$ -group magic  $\iff T$  has no vertex of even degree.*

**Lemma 13** *Let  $T$  be a tree of odd order. Then,  $T$  is not  $V_4$ -ring magic.*

**Theorem 14** *Let  $T$  be a tree. Then,  $T$  is  $V_4$ -ring magic  $\iff d(v) \equiv 1 \pmod{2}$ , for all  $v \in V(T)$ . Moreover, for this case, suppose  $T$  is of order  $p$ . Then,  $p \equiv 0 \pmod{2}$ .*





# $V_4$ -ring Magic Property for Trees

**Theorem 14** *Let  $T$  be a tree. Then,  $T$  is  $V_4$ -ring magic  $\iff d(v) \equiv 1 \pmod{2}$ , for all  $v \in V(T)$ . Moreover, for this case, suppose  $T$  is of order  $p$ . Then,  $p \equiv 0 \pmod{2}$ .*



# $V_4$ -ring Magic Property for Trees

**Theorem 14** *Let  $T$  be a tree. Then,  $T$  is  $V_4$ -ring magic  $\iff d(v) \equiv 1 \pmod{2}$ , for all  $v \in V(T)$ . Moreover, for this case, suppose  $T$  is of order  $p$ . Then,  $p \equiv 0 \pmod{2}$ .*

**Proof:**

$\Rightarrow$ : By Lemma 12



# $V_4$ -ring Magic Property for Trees

**Theorem 14** *Let  $T$  be a tree. Then,  $T$  is  $V_4$ -ring magic  $\iff d(v) \equiv 1 \pmod{2}$ , for all  $v \in V(T)$ . Moreover, for this case, suppose  $T$  is of order  $p$ . Then,  $p \equiv 0 \pmod{2}$ .*

**Proof:**

$\Rightarrow$ : By Lemma 12

$\Leftarrow$ : Suppose that  $d(v) \equiv 1 \pmod{2}$ , for all  $v \in V(T)$ . Then, the constant map  $f$  which labels every edge of  $T$  with  $x$ , where  $x \in V_4 - \{0\}$ , is a  $V_4$ -ring-magic labeling of  $T$ . Here,  $f^+ = f^\times = x$ .




# Questions on Group-magic

- ★ Let  $k \in \mathbb{Z}$  and  $k \geq 2$ . Are there graphs  $G$  whose integer-magic spectrum  $(\{k \in \mathbb{N} \mid G \text{ is } \mathbb{Z}_k\text{-magic}\})$  is  $k\mathbb{N}$ ?
- ★ Let  $k_1, k_2, \dots, k_n \in \mathbb{Z}$  and  $k_i \geq 2$ . Are there graphs  $G$  whose integer-magic spectrum is  $k_1\mathbb{N} \cup k_2\mathbb{N} \cup \dots \cup k_n\mathbb{N}$ ?
- ★ Are there classes of  $\mathbb{Z}_k$ -magic graphs having only magic-value 0?
- ★ Is it possible to construct  $\mathbb{Z}_k$ -magic graphs which have certain specified magic-values?
- ★ Let  $k \in \mathbb{Z}$  and  $k \geq 2$ . What is the “smallest” graph which has integer-magic spectrum  $\{2, 3, \dots, \} - \{k\mathbb{N}\}$ ?
- ★ Find a  $V_4$ -magic graph which is not  $Z_4$ -magic.



# Questions on Ring-magic

- ★ We can define integer-ring-magic spectrum and ask the same questions as group-magic.
- ★ Find the integer-ring-magic spectrum of some classes of graphs.



★ **Thank you** ★