

# THE INTEGER-MAGIC SPECTRA OF BICYCLIC GRAPHS WITHOUT PENDANT

W.C. SHIU AND RICHARD M. LOW

**ABSTRACT.** Let  $A$  be a non-trivial, finitely-generated abelian group and  $A^* = A - \{0\}$ . A graph is  $A$ -magic if there exists an edge labeling using elements of  $A^*$  which induces a constant vertex labeling of the graph. For connected bicyclic graphs  $G$  (without pendant), we determine the set of numbers  $k \geq 2$ , where  $G$  has a  $\mathbb{Z}_k$ -magic labeling.

## 1. INTRODUCTION

Let  $G = (V, E)$  be a connected simple graph. For any non-trivial, finitely generated abelian group  $A$  (written additively), let  $A^* = A - \{0\}$ . A mapping  $f : E \rightarrow A^*$  is called a *labeling* of  $G$ . Any such labeling induces a map  $f^+ : V \rightarrow A$ , defined by  $f^+(v) = \sum_{uv \in E} f(uv)$ . If there exists a labeling  $f$  whose induced map on  $V$  is a constant map, we say that  $f$  is an  $A$ -magic labeling of  $G$  and that  $G$  is an  $A$ -magic graph. The corresponding constant is called an  $A$ -magic value. The *integer-magic spectrum* of a graph  $G$  is the set  $\text{IM}(G) = \{k \mid G \text{ is } \mathbb{Z}_k\text{-magic and } k \geq 2\}$ . Note that the integer-magic spectrum of a graph is not to be confused with the set of achievable magic values.  $\mathbb{Z}$ -magic (or  $\mathbb{Z}_1$ -magic) graphs were considered by Stanley [29, 30], where he pointed out that the theory of magic labelings could be studied in the general context of linear homogeneous diophantine equations. Doob [1–3] and others [7, 9, 15, 16, 26] have studied  $A$ -magic graphs and  $\mathbb{Z}_k$ -magic graphs were investigated in [4, 6, 8, 10–14, 17–20, 27, 28].

## 2. BICYCLIC GRAPHS

A connected  $(p, p+1)$ -graph  $G$  is called a *bicyclic* graph. A characterization of bicyclic graphs without pendant is given in [25]. For the sake of completeness, we include the results and their proofs (Lemma 2.1, Theorems 2.2 and 2.3, and Corollary 2.4).

---

*Date:* March 28, 2012; *Version:* v.1.0.

2000 *Mathematics Subject Classification.* 05C15.

*Key words and phrases.* group-magic graphs, integer-magic graphs, integer-magic spectra, bicyclic graphs.

**Definition.** A vertex of degree  $k$  is called a  $k$ -vertex. A vertex of degree greater than  $k$  is called a  $k^+$ -vertex.

**Lemma 2.1.** Let  $G$  be a  $(p, p+1)$ -bicyclic graph without pendant. Then, the number of  $2^+$ -vertices in  $G$  is at most two.

*Proof.* It is known that  $\sum_{v \in V(G)} d(v) = 2(p+1)$ , where  $d(v)$  denotes the degree of  $v$ . Let  $x$  be the number of  $2^+$ -vertices in  $G$ . Then,  $2(p-x) + 3x \leq 2(p+1)$ . Hence,  $x \leq 2$ .  $\square$

**Definition.** A *one-point union of two cycles* is a simple graph obtained from two cycles, say  $C_m$  and  $C_n$  where  $m, n \geq 3$ , by identifying one vertex from each cycle. Without loss of generality, we may assume the  $m$ -cycle to be  $u_0 u_1 \cdots u_{m-1} u_0$  and the  $n$ -cycle to be  $u_0 u_m u_{m+1} \cdots u_{n+m-2} u_0$ . We denote this graph by  $U(m, n)$ .

**Definition.** A *theta graph* is the union of three internally disjoint (simple) paths that have the same two distinct end vertices. Without loss of generality, we may assume that  $s, t, r \geq 1$  such that  $P_s = u_0 u_1 \cdots u_s$ ,  $P_t = u_0 u_{s+1} u_{s+2} \cdots u_{s+t-1} u_s$  and  $P_r = u_0 u_{s+t} u_{s+t+1} \cdots u_{s+t+r-2} u_s$ . Note that in this paper,  $P_i$  denotes a path of length  $i$ . We denote this graph by  $\Theta(s, t, r)$ .

**Definition.** A *long dumbbell graph* is a simple graph obtained from two cycles  $C_m$  and  $C_n$ , by joining a path of length  $l$  (again, denoted by  $P_l$  in this paper) for  $m, n \geq 3$  and  $l \geq 1$ . Without loss of generality, we may assume

$$C_m = u_0 u_1 \cdots u_{m-1} u_0, \quad P_l = u_{m-1} u_m \cdots u_{m+l-1}$$

$$\text{and } C_n = u_{m+l-1} u_{m+l} \cdots u_{m+n+l-2} u_{m+l-1}.$$

We denote this graph by  $D(m, n; l)$ .

**Theorem 2.2.** Let  $G$  be a  $(p, p+1)$ -bicyclic graph without pendant. Then,  $G$  contains only one  $2^+$ -vertex if and only if  $G$  is a one-point union of two cycles.

*Proof.* Suppose  $G$  contains only one  $2^+$ -vertex. Let  $d$  be the degree of the  $2^+$ -vertex. Since  $2(p-1) + d = 2(p+1)$ ,  $d = 4$ . Since  $G$  contains one 4-vertex and  $(p-1)$  2-vertices,  $G$  is eulerian and contains two cycles. Hence,  $G$  is a one-point union of two cycles. The converse is clear.  $\square$

**Theorem 2.3.** Let  $G$  be a  $(p, p+1)$ -bicyclic graph without pendant. Then,  $G$  contains two  $2^+$ -vertices if and only if  $G$  is either a long dumbbell graph or a theta graph.

*Proof.* Suppose  $G$  contains only two  $2^+$ -vertices. Let  $d$  be the sum of the degrees of the  $2^+$ -vertices. Since  $2(p-2) + d = 2(p+1)$ ,  $d = 6$ . Since the degree of the  $2^+$ -vertices is greater than 2, the two  $2^+$ -vertices must

be 3-vertices. Since  $G$  contains two 3-vertices and  $(p - 2)$  2-vertices,  $G$  is edge-traceable. The two 3-vertices are connected to each other by joining either one path or three disjoint paths. If the 3-vertices are connected by one path, then two disjoint cycles are incident with these 3-vertices respectively. Hence,  $G$  is a long dumbbell graph. If the vertices are connected by three paths, then  $G$  is a theta graph. The converse is clear.  $\square$

**Corollary 2.4.** *A bicyclic graph without pendant is either a one-point union of two cycles, a long dumbbell graph or a theta graph.*

### 3. MAIN RESULTS

We first recall the the following useful result, found in [16].

**Theorem A.** *Let  $G$  be a  $\mathbb{Z}_k$ -magic graph, with  $k|n$ . Then,  $G$  is a  $\mathbb{Z}_n$ -magic graph.*

Using Theorem A, we obtain the following two theorems which describe the integer-magic spectra of  $U(m, n)$ .

**Theorem 3.1.** *Let  $m, n \geq 3$ . If  $m$  and  $n$  have the same parity, then  $\text{IM}[U(m, n)] = \{2, 3, 4, \dots\}$ .*

*Proof.* Since  $m$  and  $n$  have the same parity, this implies that  $U(m, n)$  is an eulerian graph with an even number of edges. Traveling along an eulerian circuit of  $U(m, n)$ , we label its edges with  $a, -a, a, -a, \dots, a, -a$ , where  $a \neq 0$ . This yields a  $\mathbb{Z}_k$ -magic labeling, for all  $k \in \{2, 3, 4, \dots\}$ .  $\square$

Note that  $U(m, n)$  is also  $\mathbb{Z}$ -magic, when  $m$  and  $n$  have the same parity.

**Theorem 3.2.** *Let  $m, n \geq 3$ . If  $m$  and  $n$  have different parity, then  $\text{IM}[U(m, n)] = 2\mathbb{N}$ .*

*Proof.* Without loss of generality, suppose that  $m$  is even and  $n$  is odd. In order for  $U(m, n)$  to be  $\mathbb{Z}_k$ -magic, we must consecutively label the edges of  $C_m$  with non-zero  $x, y, x, y, \dots, x, y$ . The edges of  $C_n$  must then be consecutively labeled with non-zero  $z, x+y-z, z, x+y-z, \dots, z, x+y-z, z$ . See Figure 1. Then,  $U(m, n)$  has a  $\mathbb{Z}_k$ -magic labeling  $\iff x+y \equiv x+y+2z \pmod{k}$ . In order for  $0 \equiv 2z \pmod{k}$ ,  $k$  must be even. Clearly,  $U(m, n)$  is  $\mathbb{Z}_2$ -magic, since every vertex is of even degree. Using Theorem A, this implies that  $U(m, n)$  is  $\mathbb{Z}_4$ -magic,  $\mathbb{Z}_6$ -magic, etc.  $\square$




FIGURE 1

Next, we establish the integer-magic spectra of long dumbbell graphs.

**Theorem 3.3.** *Let  $m, n \geq 3$ , and  $l \geq 1$ . If  $m \cdot n$  is even, then  $\text{IM}[D(m, n; l)] = \emptyset$ .*

*Proof.* Since  $m \cdot n$  is even, (WLOG) we can assume that  $m$  is even. If  $D(m, n; l)$  has a  $\mathbb{Z}_k$ -magic labeling, then the edges of subgraph  $C_m$  must be consecutively labeled with non-zero  $x, y, x, y, \dots, x, y$  (with its vertices having magic value  $x + y$ ). However, this would force the edge (from the path of length  $l$ ) connecting  $C_m$  to be labeled 0. Hence,  $D(m, n; l)$  is not  $\mathbb{Z}_k$ -magic, for all  $k$ .  $\square$

**Theorem 3.4.** *Let  $m, n \geq 3$ , and  $l \geq 1$ . If  $m \cdot n$  is odd, then  $\text{IM}[D(m, n; l)] = \{3, 4, 5, \dots\}$ .*

*Proof.* Since  $m \cdot n$  is odd,  $m$  and  $n$  must both be odd. We first note that  $D(m, n; l)$  is not  $\mathbb{Z}_2$ -magic, since it contains vertices of different parity. Figure 2 provides a  $\mathbb{Z}_k$ -magic labeling of  $D(m, n; l)$ , for  $k = 4, 5, 6, \dots$ . Figures 3 and 4 provide  $\mathbb{Z}_3$ -magic labelings of  $D(m, n; l)$ , for  $l$  odd and even, respectively.




FIGURE 2.  $\mathbb{Z}_k$ -magic labeling of  $D(m, n; l)$ , for  $k = 4, 5, 6, \dots$




FIGURE 3.  $\mathbb{Z}_3$ -magic labeling of  $D(m, n; l)$ , for odd  $l$




FIGURE 4.  $\mathbb{Z}_3$ -magic labeling of  $D(m, n; l)$ , for even  $l$

□

Lastly, we focus on the integer-magic spectra of theta graphs.

**Theorem 3.5.**  $\Theta(s, t, r)$  is  $\mathbb{Z}_2$ -magic if and only if  $s = t = r = 1$ .

*Proof.* A graph  $G$  is  $\mathbb{Z}_2$ -magic if and only if the degrees of the vertices are of the same parity. □

**Theorem 3.6.** Let  $s \cdot t \cdot r > 1$ . If exactly one of  $s, t$  and  $r$  is odd, then  $\text{IM}[\Theta(s, t, r)] = \{2k \mid k \geq 2\}$ . If exactly one of  $s, t$  and  $r$  is even, then  $\text{IM}[\Theta(s, t, r)] = \{4, 5, 6, \dots\}$ . Otherwise,  $\text{IM}[\Theta(s, t, r)] = \{3, 4, 5, \dots\}$ .

*Proof.* Let  $G = \Theta(s, t, r)$ , where  $s, t, r \geq 1$ . Without loss of generality, let the three internally disjoint (simple) paths (having the same two distinct end vertices) be  $P_s = u_0 u_1 \cdots u_s$ ,  $P_t = u_0 u_{s+1} u_{s+2} \cdots u_{s+t-1} u_s$  and  $P_r = u_0 u_{s+t} u_{s+t+1} \cdots u_{s+t+r-2} u_s$ . Suppose that  $G$  is  $\mathbb{Z}_k$ -magic (with magic labeling  $f$ ) and has  $\mathbb{Z}_k$ -magic value  $m$ . Furthermore, let  $x_1, x_2, \dots, x_s$  be the edge labeling of  $P_s$  (under  $f$ , read from left to right). Similarly, let  $y_1, y_2, \dots, y_t$  and  $z_1, z_2, \dots, z_r$  be the edge labelings of  $P_t$  and  $P_r$ , respectively (under  $f$ , read from left to right).

**CASE 1.** ( $s$  is odd and  $t, r$  are even): In this case, we must have that  $x_1 + y_1 + z_1 = m$  and  $x_1 + (m - y_1) + (m - z_1) = m$ . This implies that

$2x_1 = 0$ . Thus,  $\text{IM}(G) \subseteq \{4, 6, 8, \dots\}$ . Figure 5 gives a  $\mathbb{Z}_{2n}$ -magic labeling of  $G$ , for  $n \geq 2$ .




FIGURE 5.  $\mathbb{Z}_{2n}$ -magic labeling of  $\Theta(s, t, r)$ ,  $n \geq 2$ , for  $s$  odd and  $t, r$  even.

CASE 2. ( $s, t$  are odd and  $r$  is even): In this case, we must have that  $x_1 + y_1 + z_1 = m$  and  $x_1 + y_1 + (m - z_1) = m$ . This implies that  $x_1 + y_1 + z_1 = m$  and  $x_1 + y_1 - z_1 = 0$ . Figure 6 gives a  $\mathbb{Z}_{2n}$ -magic labeling of  $G$ , for  $n \geq 2$ . Figure 7 gives a  $\mathbb{Z}_{2n+1}$ -magic labeling of  $G$ , for  $n \geq 3$ . Figure 8 gives a  $\mathbb{Z}_5$ -magic labeling of  $G$ . It is straight-forward to show (using indirect proof) that  $G$  is not  $\mathbb{Z}_3$ -magic.




FIGURE 6.  $\mathbb{Z}_{2n}$ -magic labeling of  $\Theta(s, t, r)$ ,  $n \geq 2$ , for  $s, t$  odd and  $r$  even.




FIGURE 7.  $\mathbb{Z}_{2n+1}$ -magic labeling of  $\Theta(s, t, r)$ ,  $n \geq 3$ , for  $s, t$  odd and  $r$  even.




FIGURE 8.  $\mathbb{Z}_5$ -magic labeling of  $\Theta(s, t, r)$ , for  $s, t$  odd and  $r$  even.

CASE 3. ( $s, t$ , and  $r$  are odd): Figure 9 gives a  $\mathbb{Z}_k$ -magic labeling of  $G$ , for  $k \geq 3$ .




FIGURE 9.  $\mathbb{Z}_k$ -magic labeling of  $\Theta(s, t, r)$ ,  $k \geq 3$ , for  $s, t$ , and  $r$  odd.

CASE 4. ( $s, t$ , and  $r$  are even): Figure 10 gives a  $\mathbb{Z}_k$ -magic labeling of  $G$ , for  $k \geq 3$ .




FIGURE 10.  $\mathbb{Z}_k$ -magic labeling of  $\Theta(s, t, r)$ ,  $k \geq 3$ , for  $s, t$ , and  $r$  even.

□

## REFERENCES

- [1] M. Doob, On the construction of magic graphs, *Proc. Fifth S.E. Conference on Combinatorics, Graph Theory and Computing* (1974), 361–374.
- [2] M. Doob, Generalizations of magic graphs, *Journal of Combinatorial Theory, Series B*, **17** (1974), 205–217.
- [3] M. Doob, Characterizations of regular magic graphs, *Journal of Combinatorial Theory, Series B*, **25** (1978), 94–104.
- [4] M.C. Kong, S-M Lee, and H. Sun, On magic strength of graphs, *Ars Combinatoria*, **45** (1997), 193–200.
- [5] A. Kotzig and A. Rosa, Magic valuations of finite graphs, *Canad. Math. Bull.*, **13** (1970), 451–461.
- [6] S-M Lee, Yong-Song Ho and R.M. Low, On the integer-magic spectra of maximal planar and maximal outerplanar graphs, *Congressus Numerantium*, **168** (2004), 83–90.
- [7] S-M Lee, A. Lee, Hugo Sun, and Ixin Wen, On group-magic graphs, *JCMCC*, **38** (2001), 197–207.
- [8] S-M Lee and F. Saba, On the integer-magic spectra of two-vertex sum of paths, *Congressus Numerantium*, **170** (2004), 3–15.
- [9] S-M Lee, F. Saba, E. Salehi, and H. Sun, On the  $V_4$ -group magic graphs, *Congressus Numerantium*, **156** (2002), 59–67.
- [10] S-M Lee, F. Saba, and G.C. Sun, Magic strength of the  $k$ -th power of paths, *Congressus Numerantium*, **92** (1993), 177–184.
- [11] S-M Lee and E. Salehi, Integer-magic spectra of amalgamations of stars and cycles, *Ars Combinatoria*, **67** (2003), 199–212.
- [12] S-M Lee, E. Salehi and H. Sun, Integer-magic spectra of trees with diameters at most four, *JCMCC*, **50** (2004), 3–15.
- [13] S-M Lee, L. Valdes, and Yong-Song Ho, On group-magic spectra of trees, double trees and abbreviated double trees, *JCMCC*, **46** (2003), 85–95.
- [14] R.M. Low and S-M Lee, On the integer-magic spectra of tessellation graphs, *Australas. J. Combin.*, **34** (2006), 195–210.



- [15] R.M. Low and S-M Lee, On the products of group-magic graphs, *Australas. J. Combin.*, **34** (2006), 41–48.
- [16] R.M. Low and S-M Lee, On group-magic eulerian graphs, *JCMCC*, **50** (2004), 141–148.
- [17] R.M. Low and L. Sue, Some new results on the integer-magic spectra of tessellation graphs, *Australas. J. Combin.*, **38** (2007), 255–266.
- [18] E. Salehi, Zero-sum magic graphs and their null sets, *Ars Combinatoria*, **82** (2007), 41–53.
- [19] E. Salehi, On zero-sum magic graphs and their null sets, *Bulletin of the Institute of Mathematics, Academia Sinica*, **3** (2008), 255–264.
- [20] E. Salehi and P. Bennett, On integer-magic spectra of caterpillars, *JCMCC*, **61** (2007), 65–71.
- [21] J. Sedláček, On magic graphs, *Math. Slov.*, **26** (1976), 329–335.
- [22] J. Sedláček, Some properties of magic graphs, in *Graphs, Hypergraph, and Bloc Syst. 1976, Proc. Symp. Comb. Anal., Zielona Gora* (1976), 247–253.
- [23] W.C. Shiu, P.C.B. Lam and S-M. Lee, Edge-magicness of the composition of a cycle with a null graph, *Congressus Numerantium*, **132** (1998), 9–18.
- [24] W.C. Shiu, P.C.B. Lam and S-M. Lee, On a construction of supermagic graphs, *JCMCC*, **42** (2002), 147–160.
- [25] W.C. Shiu, M.H Ling and R.M. Low, The edge-graceful spectra of connected bicyclic graphs without pendant, *JCMCC*, **66** (2008), 171–185.
- [26] W.C. Shiu and R.M. Low, Group-magicness of complete  $N$ -partite graphs, *JCMCC*, **58** (2006), 129–134.
- [27] W.C. Shiu and R.M. Low, Integer-magic spectra of sun graphs, *J. Comb. Optim.*, **14** (2007), 309–321.
- [28] W.C. Shiu and R.M. Low,  $\mathbb{Z}_k$ -magic labelings of fans and wheels with magic-value zero, *Australas. J. Combin.*, **45** (2009), 309–316.
- [29] R.P. Stanley, Linear homogeneous diophantine equations and magic labelings of graphs, *Duke Math. J.*, **40** (1973), 607–632.
- [30] R.P. Stanley, Magic labeling of graphs, symmetric magic squares, systems of parameters and Cohen–Macaulay rings, *Duke Math. J.*, **40** (1976), 511–531.
- [31] W.D. Wallis, *Magic Graphs*, Birkhauser Boston, (2001).

DEPARTMENT OF MATHEMATICS, HONG KONG BAPTIST UNIVERSITY, 224 WATERLOO ROAD, KOWLOON TONG, HONG KONG  
*E-mail address:* wcshiu@hkbu.edu.hk

DEPARTMENT OF MATHEMATICS, SAN JOSE STATE UNIVERSITY, SAN JOSE, CA 95192, USA  
*E-mail address:* richard.low@sjsu.edu