

Linear Vertex Arboricity, Independence Number and Clique Cover Number

Peter Che Bor Lam,¹ Wai Chee Shiu,¹ Feng Sun,² Jianfang Wang³ and Guiying Yan.⁴

Abstract

The linear vertex-arboricity of a graph G is defined to the minimum number of subsets into which the vertex-set G can be partitioned so that every subset induces a linear forest. In this paper, we give the upper and lower bounds for sum and product of linear vertex-arboricity with independence number and with clique cover number respectively. All of these bounds are sharp.

Key words and phrases: Linear vertex-arboricity, independent number, clique cover number.

AMS 1991 Subject Classifications: 05C15, 05C35.

1. Introduction

Throughout this paper, all graphs are simple and finite. Let $G = (V, E)$ be a graph. A subset of V is called an LV -set if it induces a linear forest in G . A partition of V is called an LV -partition if every subset in the partition is an LV -set. *Linear-vertex arboricity* of G , denoted by $\rho'(G)$, is the smallest number of subsets into which the vertex set V can be partitioned so that the partition is an LV -partition.

A clique is a subset of V such that its induced subgraph is a complete graph. A clique is *maximum* if no other clique of G is of larger order. *Clique number* of G , denoted by $\omega(G)$, is the number of vertices in a maximum clique of G . A *clique cover* of size k is a partition of the vertex set V into V_1, V_2, \dots, V_k , such that each V_i is a clique, $1 \leq i \leq k$. The *clique cover number* of G , denoted by $\vartheta(G)$, is the number of cliques in a smallest clique cover of G . An *independent* set is a subset of vertices, no two of which are adjacent. $\alpha(G)$ is the number of vertices in an independent set of maximum order. It is also called the *independence number* of G .

In [1], Chartrand et al introduced the concept of vertex-arboricity as a generalization of vertex coloring. Linear vertex-arboricity of graphs was first mentioned by Harary [2].

¹Department of Mathematics, Hong Kong Baptist University. Work is partially supported by FRG, HKBU, and RGC Grant Hong Kong.

²Rally International Inc., Forest Park, IL 60130, USA.

³Institute of Applied Mathematics, Chinese Academy of Sciences, and Asia-Pacific Operational Research Center, Beijing, China. Work is partially supported by the National Science Foundation of China and Tian Yuan Grant No. 19136014

⁴Institute of Applied Mathematics, Chinese Academy of Sciences, Beijing, China.

This is followed by a series of papers by other researchers. For examples, see [3] to [6]. In this paper, we consider the sum and product of linear vertex-arboricity with given independence number and given clique cover number respectively. We shall obtain various lower and upper bounds, all of which are sharp. For terms and notations not defined in the paper, refer to [7].

2. Lemmas

In this section, we shall present some Lemmas. The first Lemma can be found in [6].

Lemma 2.1 Suppose M is a maximum clique of a graph $G = (V, E)$, and N is a clique in $\langle V \setminus M \rangle$. Then there exists a one-to-one mapping $f : N \rightarrow M$ such that for any $n \in N$, $nf(n) \notin E$.

The following Lemma is a direct consequence of Lemma 2.1.

Lemma 2.2 For any graph $G = (V, E)$, we can construct a sequence of cliques Q_1, Q_2, \dots, Q_h with the following properties:

1. $|Q_i| \geq |Q_j| \geq 1$ if $j > i$,
2. $Q_i \cap Q_j = \emptyset$ if $i \neq j$,
3. Q_i is a maximum clique in $G - \bigcup_{j=1}^{i-1} Q_j$,
4. $\bigcup_{j=1}^h Q_j = V$, and
5. For each $1 \leq i \leq h$, we may label elements of Q_i by v_i^k , where $1 \leq k \leq |Q_i|$, so that $v_j^k v_{j+1}^k \notin E$ for $1 \leq j \leq h-1$ and $1 \leq k \leq |Q_{j+1}|$.

A sequence of cliques of $G = (V, E)$ is called a *standard sequence* of cliques, or simply a *standard sequence* if it satisfies the above conditions. The following Lemma can also be found in [6].

Lemma 2.3 Suppose a standard sequence is embedded into $\mathbb{Z}^+ \times \mathbb{Z}^+$ so that the j -th clique lies in the j -column, and for each clique, the k -th vertex lies in the k -th row. Then any four vertices in four adjacent columns and in the same row is an *LV-set*.

Lemma 2.4 If G is a graph of order n , then

$$\rho'(G) + \vartheta(G) \leq n + 1, \quad (1)$$

$$\rho'(G) \cdot \vartheta(G) \leq \lfloor (n + 3)^2 / 8 \rfloor. \quad (2)$$

Proof We shall use mathematical induction to obtain the upper bounds for (1) and (2). We can verify that $\rho'(G) + \vartheta(G) \leq n + 1$ for $n = 1$ or 2 . Suppose that $\rho'(G) + \vartheta(G) \leq n + 1$ holds for $n = 1, 2, \dots, k - 1$, where $k \geq 3$. Consider any graph G of order k . If G is edgeless, then $\rho'(G) + \vartheta(G) = 1 + k$. Suppose G has at least one edge uv . Let $G' = \langle V \setminus \{u, v\} \rangle$ and $G'' = \langle \{u, v\} \rangle$. Then

$$\rho'(G) + \vartheta(G) \leq \rho'(G') + \vartheta(G') + \rho'(G'') + \vartheta(G'') \leq k + 1,$$

which establishes the upper bound of (1).

To establish the upper bound of (2), we first verify that $\rho'(G) \cdot \vartheta(G) \leq \lfloor (n + 3)^2 / 8 \rfloor$ for $1 \leq n \leq 16$. Suppose that $\rho'(G) \cdot \vartheta(G) \leq \lfloor (n + 3)^2 / 8 \rfloor$ for $1 \leq n \leq k - 1$, where $k \geq 17$. Consider any graph G of order k . Let Q_1, Q_2, \dots, Q_h be a standard sequence of cliques. We shall denote the number of s -cliques in this sequence by $f(s)$.

Suppose $|Q_4| \geq 4$. Let $Q_i = \{u_i^1, u_i^2, u_i^3, u_i^4, \dots\}$ for $i = 1, 2, 3, 4$. Because the sequence is standard, $A_j = \{u_1^j, u_2^j, u_3^j, u_4^j\}$ are LV -sets for $j = 1, 2, 3, 4$. Let $G^* = \langle V - (\bigcup_{j=1}^4 A_j) \rangle$. Then

$$\begin{aligned} \rho'(G) \cdot \vartheta(G) &\leq \{\rho'(G^*) + 4\} \{\vartheta(G^*) + 4\} \\ &\leq \rho'(G^*) \cdot \vartheta(G^*) + 4\{\rho'(G^*) + \vartheta(G^*)\} + 16 \\ &\leq \frac{(k - 16 + 3)^2}{8} + 4\{k - 16 + 1\} + 16 \\ &\leq \frac{(k + 3)^2}{8}. \end{aligned}$$

So we may assume that $\sum_{r \geq 4} f(r) \leq 3$. Suppose $\sum_{r \geq 4} f(r) = 1$ and $w = |Q_1|$. Also suppose the numbers of 3-, 2- and 1-cliques are x , y and z respectively. Then

$$k = w + 3x + 2y + z, \quad (3)$$

$$\rho'(G) \leq (w + 1)/2 + \epsilon_1 + (3x + 2y + \epsilon_2)/4, \quad (4)$$

$$\vartheta(G) \leq (1 - \epsilon_1) + x + y + z, \quad (5)$$

where $\epsilon_1 = 0$ if there exists one vertex in Q_1 which is not adjacent to vertices in any 1-clique of the standard sequence, and $\epsilon_1 = 1$ otherwise. If $\epsilon_1 = 1$ then $\vartheta(G) \leq x + y + z$.

To obtain (4), we partition Q_1 into $\frac{w}{2}$ 2-subsets if w is even, and $\frac{w-1}{2}$ 2-subsets plus one 1-subset if w is odd. By Lemma 2.3, we may partition vertices in 3- and 2-cliques into sets of four vertices along the lines $y = 1, 2$ and 3 , each inducing one linear forest in G . In the right side of (4), ϵ_2 is a correctional term to allow for subsets having less than four vertices. Substituting (3) into (4) and (5), we get

$$\begin{aligned}\rho'(G) \cdot \vartheta(G) &\leq \frac{(k + w - z + 2 + \epsilon_2 + 4\epsilon_1)(k - w - x + z + 2 - 2\epsilon_1)}{8} \\ &\leq \frac{\{k + 2 + (\epsilon_2 - x + 2\epsilon_1)/2\}^2}{8}.\end{aligned}$$

The Lemma follows if we manage to chose ϵ_2 to satisfy $\epsilon_2 \leq 2 + x - 2\epsilon_1$. In the following cases/sub-cases, let $x + y \equiv t \pmod{4}$, where $0 \leq t \leq 3$.

Case 1 $x = 0$.

Sub-case 1.1 $\epsilon_1 = 1$

If $t = 0$, then we may choose $\epsilon_2 = 0$ because the number of 2-cliques is a multiple of 4. If $t = 1$, then four vertices of Q_1 and one 2-clique may form two LV -sets (Figure 1a). Therefore

$$\begin{aligned}\rho'(G) &\leq (w - 4 + 1)/2 + \epsilon_1 + 2 + (3x + 2y - 2)/4 \\ &\leq (w + 1)/2 + \epsilon_1 + (3x + 2y)/4,\end{aligned}$$

and we may choose $\epsilon_2 = 0$.




Figure 1 - $x = 0$.

If $t = 2$ or 3 , then two vertices of Q_1 and two or three 2-cliques respectively, may be partitioned into two LV -sets, (Figure 1b and 1c). So

$$\begin{aligned}\rho'(G) &\leq (w - 2 + 1)/2 + \epsilon_1 + 2 + (3x + 2y - 4)/4 \\ &= (w + 1)/2 + \epsilon_1 + (3x + 2y)/4,\end{aligned}$$

and we may also choose $\epsilon_2 = 0$.

Sub-case 1.2 $\epsilon_1 = 0$

If $t = 0$ then $\epsilon_2 = 0$ because the number of 2-cliques is a multiple of 4. If $t = 1$ or 3 then $\epsilon_2 = 2$ because the total number of vertices in 2-cliques plus 2 is a multiple of 4. If $t = 2$, then three vertices of Q_1 together two 2-cliques and all the 1-cliques may be partitioned into three LV -sets (Figure 1d). So

$$\begin{aligned}\rho'(G) &\leq (w - 3 + 1)/2 + 3 + (3x + 2y - 4)/4 \\ &= (w + 1)/2 + \epsilon_1 + (3x + 2y + 2)/4,\end{aligned}$$

and we may also choose $\epsilon_2 = 2$.

Case 2 $x = 1$.

Sub-case 2.1 $\epsilon_1 = 1$

If $t = 0$, then two vertices of Q_1 together with one vertex in the 3-clique forms one LV -set (Figure 2a), and

$$\begin{aligned}\rho'(G) &\leq (w - 2 + 1)/2 + \epsilon_1 + 1 + (3x + 2y - 1)/4 \\ &= (w + 1)/2 + \epsilon_1 + (3x + 2y - 1)/4.\end{aligned}$$




Figure 2 - $x = 1$, $\epsilon_1 = 1$.

If $t = 1$, then three vertices of Q_1 together with vertices in the 3-clique two form two LV -sets (Figure 2b), and

$$\begin{aligned}\rho'(G) &\leq (w - 3 + 1)/2 + \epsilon_1 + 2 + (3x + 2y - 3)/4 \\ &= (w + 1)/2 + \epsilon_1 + (3x + 2y - 1)/4.\end{aligned}$$

If $t = 2$, then four vertices of Q_1 together with vertices in the 3-clique and one 2-clique to form three LV -sets (Figure 2c), and

$$\begin{aligned}\rho'(G) &\leq (w - 4 + 1)/2 + \epsilon_1 + 3 + (3x + 2y - 5)/4 \\ &= (w + 1)/2 + \epsilon_1 + (3x + 2y - 1)/4.\end{aligned}$$

If $t = 3$, then four vertices of Q_1 together with vertices in the 3-clique and two 2-clique to form three LV -sets (Figure 2d), and

$$\begin{aligned}\rho'(G) &\leq (w - 4 + 1)/2 + \epsilon_1 + 3 + (3x + 2y - 7)/4 \\ &= (w + 1)/2 + \epsilon_1 + (3x + 2y - 3)/4.\end{aligned}$$

So for any t , we may also choose $\epsilon_2 = 0$.

Sub-case 2.2 $\epsilon_1 = 0$

For any value of t , argument of sub-case 2.1 is still valid if we replace ϵ_1 by 1, and we get

$$\begin{aligned}\rho'(G) &\leq (w + 1)/2 + 1 + (3x + 2y - 1)/4 \\ &= (w + 1)/2 + (3x + 2y + 3)/4.\end{aligned}$$

Therefore for any t , we may choose $\epsilon_2 = 3$.

Cases where $x \geq 2$, and where $\sum_{r \geq 4} f(r) = 0$, 2 and 3 may be dealt with in the similar manner. ■

3. Main results

Theorem 3.1 If $G = (V, E)$ is a graph of order n , then

$$\lceil \sqrt{2n} \rceil \leq \rho'(G) + \alpha(G) \leq n + 1, \quad (6)$$

$$\lceil n/2 \rceil \leq \rho'(G) \cdot \alpha(G) \leq \lfloor (n + 3)^2/8 \rfloor. \quad (7)$$

Proof Let $\rho'(G) = k$. It follows that $\alpha(G) \geq n/(2k)$ and $\rho'(G) \cdot \alpha(G) \geq n/2$. Therefore

$$\rho'(G) + \alpha(G) \geq k + n/(2k) = \left[\sqrt{k} - \sqrt{n/(2k)} \right]^2 + \sqrt{2n},$$

from which $\rho'(G) + \alpha(G) \geq \lceil \sqrt{2n} \rceil$ follows. The upper bounds follows from Lemma 2.4 because the clique cover number is in general not less than the independence number. ■

The bounds in Theorem 3.1 are sharp. For the upper bound of (6), equality holds for a null graph G on n vertices, and for the lower bound of (7), equality holds for a complete graph on n vertices. Stronger results on sharpness of other bounds are obtained in the following two theorems.

Theorem 3.2 For any positive integer n , there exists a graph G of order n such that

$$\rho'(G) + \alpha(G) = \lceil \sqrt{2n} \rceil.$$

Proof Let m be the positive integer such that $2(m-1)^2 < n \leq 2m^2$ and let $l = n - 2(m-1)^2$, then $0 < l \leq 2m^2 - 2(m-1)^2 = 4m - 2$.

If $l \leq 2m - 2$, then $2m - 2 < \sqrt{2n} < 2m - 1$. We construct $G = A_1 \vee A_2 \vee \cdots \vee A_{m-1}$, where A_i are paths of order $2m$ for $1 \leq i \leq m - 2$ and of order $l + 2$ for $i = m - 1$. We can see that $|G| = n$, $\rho'(G) = m - 1$, $\alpha(G) = m$, $\rho'(G) + \alpha(G) = 2m - 1 = \lceil \sqrt{2n} \rceil$.

If $2m - 1 \leq l$, then $2m - 1 < \sqrt{2n} \leq 2m$. We construct $G = A_1 \vee A_2 \vee \cdots \vee A_m$, where A_i are paths of order $2m$ for $1 \leq i \leq m - 1$ and of order $l + 2 - 2m$ for $i = m$. We can see that $|G| = n$, $\rho'(G) = m$, $\alpha(G) = m$, $\rho'(G) + \alpha(G) = 2m = \lceil \sqrt{2n} \rceil$. ■

Theorem 3.3 For any positive integer $n \geq 4$, there exists a graph G of order n such that

$$\rho'(G) \cdot \alpha(G) = \left\lfloor \frac{(n+3)^2}{8} \right\rfloor. \quad (8)$$

Proof Let $G = N_t \vee K_{n-t}$, where N_t is a null graph of order t . Then $\alpha(G) = t$ and $\rho'(G) = \lfloor (n+3-t)/2 \rfloor$. Set $t = (n+2+s)/2$, where $s \equiv n \pmod{4}$, then $\alpha(G) = (n+2+s)/2$, $\rho'(G) = (n+4-s)/4$, and

$$\rho'(G) \cdot \alpha(G) = \frac{(n+3)^2 - (s-1)^2}{8} \quad (9)$$

Because $\rho'(G) \cdot \alpha(G)$ is an integer and $(s-1)^2/8 \leq 1/2$, (8) follows from (9). ■

For any graph G , we have $\omega(\bar{G}) = \alpha(G)$. The following corollary is a direct consequence of Theorem 3.1.

Corollary 3.4 For any graph G of order n ,

$$\begin{aligned} \lceil \sqrt{2n} \rceil &\leq \rho'(G) + \omega(\bar{G}) \leq n + 1, \\ \lfloor n/2 \rfloor &\leq \rho'(G) \cdot \omega(\bar{G}) \leq \lfloor (n+3)^2/8 \rfloor. \end{aligned}$$

and all of the above bounds are sharp.

Now we can show that the upper bounds and lower bounds for sum and product of linear vertex-arboricity and clique cover number is the same as those in Theorem 3.1.

Theorem 3.5 If G is a graph of order n , then

$$\lceil \sqrt{2n} \rceil \leq \rho'(G) + \vartheta(G) \leq n + 1, \quad (10)$$

$$\lceil n/2 \rceil \leq \rho'(G) \cdot \vartheta(G) \leq \lfloor (n+3)^2/8 \rfloor. \quad (11)$$

and all of the bounds are sharp.

Proof The upper bounds of both (10) and (11) is given in Lemma 2.4. The lower bounds of both (10) and (11) follows from Theorem 3.1 and the fact that $\vartheta(G) \geq \alpha(G)$ for any graph G . ■

References

- [1] G. CHARTRAND, D. P. GELLER and S. HEDETNIEMI, A Generalization of the Chromatic Number. *Proc. Camb. Phil. Soc.*, 1968, **64** (2), 365-371.
- [2] F. HARARY, Conditional Colorability in Graphs. *First Colorado Symposium on Graph Theory*. Wiley-Interscience, New York, 1984, 123-138.
- [3] Y. ALAVI, J. LIU and J. WANG. On Linear Vertex-arboricity of Complementary Graphs. *J. Graph Theory*, 1994, **18** (3), 315-322.
- [4] Y. ALAVI, D. GREEN, J. LIU and J. WANG. On Linear Vertex-arboricity of Graphs. *Congressus Numeratim*, 1991, **82**, 187-192.
- [5] Y. ALAVI, D. R. LICK, J. LIU and J. WANG. Bounds for the Linear Vertex-arboricity and Domination Number of Graphs. *Vishna International Journal of Graph Theory*, 1992, **1** (2), 95-102.
- [6] Y. ALAVI, P. C. B. LAM, D. R. LICK, P. ERDŐS, J. LIU and J. WANG, Upper Bounds on Linear-Vertex Arboricity of Complementary Graphs. To appear in *Utilitas Math*.

- [7] D. B. WEST, *Introduction to Graph Theory*, Prentice-Hall, London 1996.
- [8] M. C. GOLUBIC, *Algorithmic Graph Theory and Perfect Graphs*, Academic Press, New York, 1980.