

The neighbor expanded sum distinguishing index of Halin graphs*

Jingjing Huo^a, Wai Chee Shiu^b, Weifan Wang^c

^aDepartment of Mathematics,
Hebei University of Engineering, Handan 056038, China.

^bDepartment of Mathematics,
Hong Kong Baptist University, Kowloon Tong, Hong Kong, China.

^cDepartment of Mathematics,
Zhejiang Normal University, Jinhua 321004, China.

Abstract

Let $G = (V, E)$ be a graph. A total k -weighting c of G is a function $c : V(G) \cup E(G) \rightarrow \{1, 2, \dots, k\}$. For $x \in V(G)$, define $w(x) = \sum_{y \in N(x)} (c(xy) + c(y))$. A total k -weighting c of G is called neighbor expanded

sum distinguishing (nesd for short) if $w(u) \neq w(v)$ for every $uv \in E(G)$. The smallest value of k for which such a nesd total k -weighting of G exists is called the neighbor expanded sum distinguishing index of G , denoted by $\text{nesd}_t(G)$. In this paper, we prove that $\text{nesd}_t(G) \leq 3$ for any Halin graph G . Furthermore, $\text{nesd}_t(G) = 2$ for the cubic Halin graphs G .

Key words: Halin graphs, total weighting, neighbor expanded sum distinguishing index.

AMS Subject Classifications: 05C15.

1 Introduction

Graphs considered in this paper are finite and simple. The terminologies and notation used but undefined in this paper can be found in [1]. Let $V(G)$ and $E(G)$ denote the vertex set and the edge set of a graph G , respectively.

*Research was done while the first author visited Department of Mathematics, Hong Kong Baptist University in 2015. Partially supported by NSFC (No.11371328, No.11501161, No.11701136) and NSFHB (No.A2016402164), Tianjin Research Program of Application Foundation and Advanced Technology (No.14JCYBJC43100), National Natural Science Foundation of China (No.11601391). E-mails: jingjinghuo@hotmail.com (Huo); shiu1958@gmail.com; wcshiu@math.hkbu.edu.hk; (Shiu, corresponding author); ww@zjnu.cn (Wang).

We call $T(G) = V(G) \cup E(G)$ the total set of G . Let $N(v)$ denote the set of neighbors of v in G and let $d(v) = |N(v)|$ denote the degree of v in G . Let $\Delta(G)$ and $\delta(G)$ denote the maximum degree and the minimum degree of a vertex in G , respectively. A 3-regular graph is called a *cubic graph*. For $n \geq 3$, an n -cycle is a cycle of length n . An n -cycle is odd (or even) if n is odd (or even). For every $n \geq 3$, a *wheel* W_n is the graph obtained from an n -cycle by adding a new vertex w and joining it to every vertex of the n -cycle. Such vertex w is called the *center* of the wheel. A wheel W_n is odd (or even) if n is odd (or even).

A Halin graph G is a plane graph constructed as follows. Let T be a tree on at least 4 vertices. All vertices of T are either of degree 1, called leaves, or of degree at least 3. Let C be a cycle connecting the leaves of T in such a way that C forms the boundary of the unbounded face. The tree T and the cycle C are called the *characteristic tree* and the *adjoint cycle* of G , respectively. For simplicity, we usually write $G = T \cup C$. The vertices in $V(C)$ and in $V(G) \setminus V(C)$ are called *outer vertices* and *inner vertices* of G , respectively. Let $V_{in}(G)$ denote the set of inner vertices in a Halin graph G . If $|V_{in}(G)| = 1$, then G is just a wheel.

Let $[k]$ denote the set $\{1, 2, \dots, k\}$. A *vertex k -weighting* c of G is a function $c : V(G) \rightarrow [k]$. A vertex k -weighting c of G is called *proper* if $c(u) \neq c(v)$ for every $uv \in E(G)$. A graph G is k -colorable if there exists a proper vertex k -weighting. The chromatic number $\chi(G)$ of G is the smallest value of k such that a graph G is k -colorable.

An *edge k -weighting* c of G is a function $c : E(G) \rightarrow [k]$. For a vertex $x \in V(G)$, we define $\sigma(x) = \sum_{y \in N(x)} c(xy)$. An edge k -weighting c of G is

called *neighbor sum distinguishing* if $\sigma(u) \neq \sigma(v)$ for every $uv \in E(G)$. Let $\text{nsd}_e(G)$ be the smallest value of k so that G admits a neighbor sum distinguishing edge k -weighting.

A graph G is *nice* if no component is isomorphic to K_2 . In [5], Karoński, Luczak and Thomason proposed the following "1-2-3 Conjecture".

Conjecture 1 *If G is a nice graph, then $\text{nsd}_e(G) \leq 3$.*

The best known upper bound on $\text{nsd}_e(G)$ is due to Kalkowski et al., who showed that $\text{nsd}_e(G) \leq 5$ if G is nice (see [4]).

A *total k -weighting* c of G is a function $c : V(G) \cup E(G) \rightarrow [k]$. For a vertex $x \in V(G)$, we define $t(x) = c(x) + \sigma(x) = c(x) + \sum_{y \in N(x)} c(xy)$. A total

k -weighting c of G is called *neighbor sum distinguishing* if $t(u) \neq t(v)$ for every $uv \in E(G)$. Let $\text{nsd}_t(G)$ be the smallest value of k so that G admits a neighbor sum distinguishing total k -weighting.

In [6], Przybyło and Woźniak posed the following "1-2 conjecture".

Conjecture 2 *If G is a connected graph, then $\text{nsd}_t(G) \leq 2$.*

It is known that for every graph G , $\text{nsd}_t(G) \leq 3$ (see [3]).

Recently, Flandrin et al. [2] introduced the neighbor expanded sum distinguishing index. For a vertex $x \in V(G)$, define

$$w(x) = \sum_{y \in N(x)} (c(xy) + c(y)),$$

where c is a total k -weighting of G . The value $w(x)$ is called an *expanded sum at x* . A total k -weighting c of G is called *neighbor expanded sum distinguishing* (nesd for short) if $w(u) \neq w(v)$ for every $uv \in E(G)$. The smallest value of k so that G admits a nesd total k -weighting of G is called the *neighbor expanded sum distinguishing index* of G , and denoted it by $\text{nesd}_t(G)$.

In the same paper, they completely determined the neighbor expanded sum distinguishing indices for paths, cycles, trees, complete graphs and complete bipartite graphs. Furthermore, they proposed the following conjecture.

Conjecture 3 For every graph G , $\text{nesd}_t(G) \leq 2$.

In this paper, we investigate the neighbor expanded sum distinguishing index of Halin graphs. To prove our main results, we need some useful results as follows.

Lemma 1.1 ([2]) Let k be an odd integer and let G be a connected k -colorable graph. Then $\text{nesd}_t(G) \leq k$.

Lemma 1.2 ([7]) If G is a Halin graph, then $\chi(G) \leq 4$ with equality if and only if G is isomorphic to an odd wheel.

2 Halin graphs

In the section, we give an upper bound on the neighbor expanded sum distinguishing index of Halin graphs.

Theorem 2.1 For $n \geq 3$, $\text{nesd}_t(W_n) = 2$.

Proof. Let w be the center of W_n and $C = v_1v_2 \cdots v_n$ be the adjoint cycle of W_n . For convenience, we let $e_i = v_iv_{i+1}$, for $1 \leq i \leq n-1$ and $e_n = v_nv_1$. The proof is split into the following cases:

Case 1: $n = 3m$.

For $m = 1$, we weight the center w with 2 and all the edges of T with 1. Then weight the sequence of $v_1, e_1, v_2, e_2, v_3, e_3$ by 1, 2, 2, 1, 1, 2 in order. Then the expanded sum at w is 7 and the expanded sums of v_1, v_2, v_3 are 10, 8, 9, respectively.

For $m \geq 2$, we weight the center w and all edges of T with 1. Then weight the vertex-edge alternating sequence of $v_1, e_1, v_2, e_2, \dots$,

v_{3m}, e_{3m} by 1, 2, 2, 1, 1, 2 cyclically. Then the expanded sums of the vertices v_1, v_2, \dots, v_{3m} are 9, 7, 8 cyclically, and the expanded sum at w is $3m + 4m = 7m \geq 14$.

Case 2: $n = 3m + 1$.

We weight w and wv_{3m+1} with 1, and all other edges of T with 2. Then weight the vertex-edge alternating sequence of $v_1, e_1, v_2, e_2, \dots, v_{3m+1}, e_{3m+1}$ by 1, 2, 2, 1, 1, 2 cyclically. Then the expanded sums of the vertices v_1, v_2, \dots, v_{3m} are 10, 8, 9 cyclically, the expanded sum at v_{3m+1} is 8, and the expanded sum at w is $6m + 1 + 4m + 1 = 10m + 2 \geq 12$.

Case 3: $n = 3m + 2$.

We weight w and all edges of T with 1. Then weight the vertex-edge alternating sequence of $v_1, e_1, v_2, e_2, \dots, v_{3m+2}, e_{3m+2}$ by 1, 2, 2, 1, 1, 2 cyclically. Then the expanded sums of the vertices $v_2, v_3, \dots, v_{3m+1}$ are 7, 8, 9 cyclically, the expanded sum at v_1 and v_{3m+2} are 9 and 7, respectively, and the expanded sum at w is $3m + 2 + 4m + 3 = 7m + 5 \geq 12$.

This completes the proof. ■

Theorem 2.2 For any Halin graph G , $\text{nesd}_t(G) \leq 3$.

Proof. If G is a wheel, then by Theorem 2.1, we obtain that $\text{nesd}_t(G) = 2$. Otherwise, we can get $\text{nesd}_t(G) \leq 3$ by Lemmas 1.1 and 1.2. Hence the theorem holds. ■

3 Cubic Halin graphs

In the section, we determine exactly the neighbor expanded sum distinguishing index of cubic Halin graphs.

Lemma 3.1 Let $G = T \cup C$ be a cubic Halin graph with $|V_{in}(G)| \geq 5$. Then G contains H_1 or H_2 of Fig. 1 as a subgraph.




Figure 1: Two subgraphs of a Halin graph.

Proof. Let $P = u_0 u_1 \cdots u_l$ be a longest path of T . Then $l \geq 4$. Let $N(u_1) = \{u_2, v_1, v_2\}$. Then v_1 and v_2 are leaves of T . Since $d(u_2) = 3$. Let $N(u_2) = \{u_1, u_3, x\}$. Suppose x is not a leaf of T . By means of the longest path P , x is adjacent to two leaves, say y_1 and y_2 . After renaming suitably, we obtain the subgraph H_1 described in Fig. 1. Suppose x is a leaf of T . We rename it by y_1 . After renaming suitably, we get the subgraph H_2 described in Fig. 1. ■

Suppose c is a total 2-weighting of a cubic graph G . Let $w^c(x)$ denote the expanded sum at x under c , for $x \in V(G)$. Let $c^* = 3 - c$. That means that c^* swaps the weights 1 to 2 and 2 to 1 assigned by c , respectively. Clearly $w^{c^*}(x) = 18 - w^c(x)$. Thus c is a nesd total 2-weighting of G if and only if c^* is a nesd total 2-weighting of G . We call c^* the *dual weighting* of c .

Theorem 3.2 For any cubic Halin graph $G = T \cup C$, $\text{nesd}_t(G) = 2$.

Proof. We prove the theorem by induction on the number of the inner vertices. If $|V_{in}(G)| = 1$, then G is a wheel and we have done by Theorem 2.1. If $2 \leq |V_{in}(G)| \leq 4$, then there are only five Halin graphs G , depicted in Fig. 2, in which a nesd total 2-weighting is established.




Figure 2: Five cubic Halin graphs of small order.

Now assume that $|V_{in}(G)| \geq 5$. By Lemma 3.1, G contains subgraph H_1 or H_2 described in Fig. 1. We shall keep the notation depicted in Fig. 1.

Case 1: G contains the subgraph H_1 .

Since $|V_{in}(G)| \geq 5$, $u_1, u_2, u_3, y_3 \in V_{in}(G)$. Note that x_1 and z_1 are distinct vertices of C . Let $H = G - \{u_1, v_1, v_2, y_1, y_2, y_3\} + \{x_1 u_2, z_1 u_2\}$. Then H is a Halin graph with $\Delta(H) = 3$ and $|V_{in}(H)| = |V_{in}(G)| - 3 \geq 2$. By the induction hypothesis, H admits a nesd total 2-weighting c . We keep the weights of all $x \in T(H) \cap T(G)$. Moreover, $w_H(u_2) \neq w_H(x_1)$ and $w_H(u_2) \neq w_H(z_1)$, where $w_H(x)$ is the expanded sum in H at x . For each case, $w_H(u_2)$ has 3 possible cases which will be shown at the following tables. By symmetry and duality, we have to deal with the following subcases.

- (1.1) $c(x_1) = c(u_2) = c(z_1) = c(x_1u_2) = c(z_1u_2) = 1$. Hence $6 \leq w_H(u_2) \leq 8$. We assign:

Conditions		Original $c(u_2u_3)$	Assignments	
$w_H(u_2)$	$w_H(z_1)$		Special rearrangement	Elements assigned with 2
6	$\neq 9$ $= 9$			$y_1, u_1v_2, y_1y_2, y_2y_3$ $y_1, u_1v_2, y_1y_3, y_2y_3$
7	$\neq 9$ $= 9$			$y_1, v_1v_2, y_1y_2, y_2y_3$ $y_1, v_1v_2, y_1y_3, y_2y_3$
8		2	$c(u_2u_3) = 1,$ $c(u_2) = 2$	$v_2, u_1v_1, u_1v_2, u_2y_3,$ y_1y_2, y_1y_3, y_2y_3

- (1.2) $c(x_1) = c(u_2) = c(x_1u_2) = c(z_1u_2) = 1$ and $c(z_1) = 2$. We assign:

Condition	Assignment
$w_H(u_2)$	Elements assigned with 2
7	$y_1, y_3, v_1v_2, v_2y_1, y_1y_2, y_1y_3, y_2y_3$
8	$v_2, y_3, u_1v_1, u_1v_2, y_1y_3$
9	$v_2, y_3, u_1v_1, v_1v_2, y_2y_3$

- (1.3) $c(x_1) = c(z_1) = c(x_1u_2) = c(z_1u_2) = 1$ and $c(u_2) = 2$. We assign:

$w_H(u_2)$	Elements assigned with 2
6	v_1, y_2, y_1y_3
7	$v_1, y_2, u_1v_1, v_2y_1, y_1y_3, y_2y_3$
8	$v_1, v_2, y_1, y_2, u_1v_2, v_1v_2, v_2y_1, y_2y_3$

- (1.4) $c(x_1) = c(u_2) = c(x_1u_2) = c(z_1) = 1$ and $c(z_1u_2) = 2$. We assign:

$w_H(u_2)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
7			$y_1, y_2, u_1v_1, u_1v_2, u_2y_3, v_2y_1$
8			$v_2, y_2, y_3, u_1v_1, u_1v_2, y_1y_2$
9	2	$c(u_2u_3) = 1,$ $c(u_2) = 2$	$u_1, y_1, u_1v_1, u_2y_3, v_1v_2,$ v_2y_1, y_2y_3, y_2z_1

- (1.5) $c(x_1) = c(u_2) = c(x_1u_2) = 1$ and $c(z_1u_2) = c(z_1) = 2$. We assign:

$w_H(u_2)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
8			$u_1, y_1, y_2, u_2y_3, v_1v_2, v_2y_1, y_1y_3$
9			$u_1, v_2, u_1v_1, u_2y_3, y_1y_2, y_2z_1$
10	2	$c(u_2u_3) = 1,$ $c(u_2) = 2$	$u_1, v_2, u_1u_2, u_1v_1, u_1v_2, u_2y_3,$ $v_1v_2, y_1y_2, y_2y_3, y_2z_1$

- (1.6) $c(x_1) = c(u_2) = c(z_1u_2) = 1$ and $c(x_1u_2) = c(z_1) = 2$. We assign:

$w_H(u_2)$	Elements assigned with 2		
8	$u_1, v_1, v_1v_2, u_2y_3, y_1y_3, y_2y_3$		
9	$u_1, y_1, u_1v_1, u_1v_2, u_2y_3, v_1x_1, v_2y_1, y_1y_3, y_2y_3$		
$w_H(u_2)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
10	2	$c(u_2u_3) = 1,$ $c(u_2) = 2$	$u_1, v_2, y_3, x_1v_1, v_1v_2,$ u_1u_2, y_1y_2, y_2y_3

- (1.7) $c(x_1) = c(x_1u_2) = c(z_1) = 1$ and $c(z_1u_2) = c(u_2) = 2$. We assign:

$w_H(u_2)$	Elements assigned with 2	
7	$v_1, y_2, u_1v_1, v_2y_1, y_1y_3, u_2y_3, y_2z_1$	
8	$v_1, v_2, y_2, u_1v_1, u_1v_2, v_2y_1, y_1y_2, y_1y_3, y_2z_1, u_2y_3$	
9	$v_2, y_2, y_3, x_1v_1, u_1v_1, u_1v_2, v_2y_1, y_1y_3, y_2y_3, y_2z_1$	

(1.8) $c(x_1) = c(u_2) = c(z_1) = 1$ and $c(x_1u_2) = c(z_1u_2) = 2$. We assign:

$w_H(u_2)$	Elements assigned with 2		
8	$u_1, v_1, y_3, v_1v_2, v_2y_1, y_1y_3, y_2z_1$		
9	$u_1, y_1, u_1v_1, u_1v_2, u_2y_3, v_1x_1, v_2y_1, y_1y_3, y_2y_3, y_2z_1$		
$w_H(u_2)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
10	2	$c(u_2u_3) = 1,$ $c(u_2) = 2$	$u_1, v_2, y_3, u_1u_2, v_1v_2,$ $v_1x_1, y_1y_2, y_2y_3, y_2z_1$

(1.9) $c(x_1) = c(x_1u_2) = c(z_1u_2) = 1$ and $c(u_2) = c(z_1) = 2$. We assign:

$w_H(u_2)$	Elements assigned with 2
7	$v_1, y_1, y_3, u_1v_2, v_1v_2, y_1y_2, y_2y_3, y_2z_1$
8	$v_2, y_2, u_1v_2, u_2y_3, v_1x_1, y_1y_3, y_2y_3$
9	$v_2, y_2, y_3, v_1v_2, v_1x_1, y_1y_2$

(1.10) $c(u_2) = c(x_1u_2) = c(z_1u_2) = 1$ and $c(x_1) = c(z_1) = 2$. We assign:

$w_H(u_2)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
8			$v_2, y_3, u_1u_2, u_1v_2, y_1y_3$
9			$u_1, v_2, y_3, v_2y_1, y_1y_2, y_1y_3$
10	2	$c(u_2u_3) = 1,$ $c(u_2) = 2$	$u_1, v_2, y_3, u_1u_2, v_1v_2,$ y_1y_2, y_2y_3

One can check that each total weighting c is a nesd total 2-weighting.

Case 2: G contains the subgraph H_2 .

Since $|V_{in}(G)| \geq 5$, we obtain that $u_1, u_2, u_3 \in V_{in}(G)$. Note that x_1 and z_1 are distinct vertices. Let $H = G - \{u_1, v_1, v_2, y_1\} + \{x_1u_2, z_1u_2\}$. Then H is a Halin graph with $\Delta(H) = 3$ and $|V_{in}(H)| = |V_{in}(G)| - 2 \geq 3$. By the induction hypothesis, H admits a nesd total 2-weighting c . We keep the weights of all $x \in T(H) \cap T(G)$. Moreover, $w_H(u_2) \neq w_H(x_1)$ and $w_H(u_2) \neq w_H(z_1)$. By duality, we have to deal with the following subcases.

(2.1) $c(x_1) = c(u_2) = c(z_1) = c(x_1u_2) = c(z_1u_2) = 1$. We assign:

Conditions				Assignments
$w_H(u_2)$	$w_H(z_1)$	$w_H(u_3)$	$w_H(x_1)$	Elements assigned with 2
6	$\neq 7$	$\neq 7$	$\neq 8$	u_1v_2, v_2y_1
	$= 7$			u_1u_2, u_1v_2
	$\neq 7$			$u_1, u_1u_2, u_2y_1, v_1v_2, v_2y_1$
	$= 7$			$v_2, u_1u_2, u_1v_2, u_2y_1, v_2y_1$
Conditions		Original $c(u_2u_3)$	Assignments	
$w_H(u_2)$	$w_H(u_3)$	$c(u_2u_3)$	Special rearrangement	Elements assigned with 2
7	$\neq 8$ $= 8$			$u_1u_2, u_1v_1, u_1v_2, v_2y_1$ $u_1, u_1u_2, u_1v_2, v_2y_1$
8	$\neq 10$ $= 10$	2	$c(u_2u_3) = 1,$ $c(u_2) = 2$	$v_2, u_1u_2, u_1v_1, u_2y_1$ v_2, u_1v_1

(2.2) $c(u_2) = c(x_1u_2) = c(z_1u_2) = c(z_1) = 1$ and $c(x_1) = 2$. We assign:

$w_H(u_2)$	$w_H(z_1)$	$w_H(x_1)$	Elements assigned with 2
7	$\neq 8$ $= 8$ $= 8$	$\neq 10$ $= 10$	$u_1, v_2, u_1v_1, v_1v_2, u_1v_2, v_2y_1$ $u_1, u_1v_1, u_1v_2, v_1v_2$ u_1v_1, u_1u_2, v_1v_2
8	$\neq 7$ $= 7$		u_1, u_1v_2, v_2y_1 u_1u_2, u_1v_1, u_1v_2

$w_H(u_2)$	$w_H(u_3)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
9	$\neq 8$ $= 8$	2	$c(u_2u_3) = 1, c(u_2) = 2$	$v_2, u_1v_1, u_1v_2, u_2y_1$ $v_2, u_1v_1, u_1v_2, v_2y_1$

(2.3) $c(x_1) = c(z_1) = c(x_1u_2) = c(z_1u_2) = 1$ and $c(u_2) = 2$. We assign:

$w_H(u_2)$	$w_H(u_3)$	$w_H(z_1)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
6	$\neq 8$ $= 8$ $= 8$	$\neq 8$ $= 8$	1	$c(u_2u_3) = 2,$ $c(u_2) = 1$	v_1, y_1 $v_1, u_1u_2, u_2y_1, y_1z_1$ v_1, y_1, u_2y_1

$w_H(u_2)$	$w_H(u_3)$	Elements assigned with 2
7	$\neq 8$ $= 8$	v_1, y_1, u_1v_1 v_1, y_1, u_1u_2, u_1v_1
8	$\neq 10$ $= 10$	$u_1, v_1, y_1, u_1v_1, v_2y_1$ $u_1, v_1, u_1v_1, u_1v_2, y_1z_1$

(2.4) $c(x_1) = c(u_2) = c(x_1u_2) = c(z_1u_2) = 1$ and $c(z_1) = 2$. We assign:

$w_H(u_2)$	$w_H(z_1)$	Elements assigned with 2
7	$\neq 9$ $= 9$	u_2y_1, v_1v_2, v_2y_1 $u_1u_2, u_1v_2, v_1v_2, v_2y_1$
8	$\neq 9$ $= 9$	$v_2, u_1u_2, u_1v_1, v_2y_1$ u_1, u_1v_2, v_1v_2

$w_H(u_2)$	$w_H(u_3)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
9	$\neq 10$ $= 10$	2	$c(u_2u_3) = 1, c(u_2) = 2$	$u_1, u_1v_1, u_1v_2,$ u_2y_1, v_1v_2, v_2y_1 $v_2, u_1u_2, u_1v_1,$ u_1v_2, v_1v_2

(2.5) $c(x_1) = c(u_2) = c(z_1u_2) = c(z_1) = 1$ and $c(x_1u_2) = 2$. We assign:

$w_H(u_2)$	$w_H(u_3)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
7	$\neq 6$ $= 6$			$v_1, u_1v_2, v_1v_2, v_2y_1$ $v_1, u_1u_2, u_1v_1, u_1v_2, u_2y_1$
8	$\neq 7$ $= 7$			$v_1, v_2, u_1v_1, u_1v_2, v_2y_1$ $u_1, v_1, u_2y_1, v_1v_2, v_2y_1$
9	$\neq 11$ $= 11$	2	$c(u_2u_3) = 1,$ $c(u_2) = 2$	$u_1, v_1, v_2, u_1u_2, u_1v_2,$ u_2y_1, v_1v_2, v_2y_1 $v_2, u_1v_1, u_1v_2,$ v_1x_1, v_2y_1

(2.6) $c(x_1) = c(u_2) = c(x_1u_2) = c(z_1) = 1$ and $c(z_1u_2) = 2$. We assign:

$w_H(u_2)$	$w_H(u_3)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
7	$\neq 8$ $= 8$			y_1, u_2y_1, v_1v_2 u_1, y_1, u_2y_1
8	$\neq 9$ $= 9$			$u_1, u_1u_2, v_1v_2, v_2y_1, y_1z_1$ $v_2, u_1v_1, u_1v_2, y_1z_1$
9	$\neq 10$ $= 10$	2	$c(u_2u_3) = 1,$ $c(u_2) = 2$	$u_1, u_1v_1, u_1v_2, v_1v_2,$ v_2y_1, u_2y_1, y_1z_1 $v_2, u_1u_2, u_1v_1,$ u_1v_2, v_1v_2, y_1z_1

(2.7) $c(x_1u_2) = c(z_1u_2) = c(z_1) = 1$ and $c(u_2) = c(x_1) = 2$. We assign:

$w_H(u_2)$	$w_H(u_3)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
7	$\neq 8$ $= 8$	1	$c(u_2u_3) = 2,$ $c(u_2) = 1$	v_1, y_1, u_1u_2 v_1, y_1, u_1v_2, u_2y_1
8	$\neq 10$ $= 10$			$u_1, v_1, y_1, u_1u_2, v_2y_1$ v_1, u_1v_1, y_1z_1
9	$\neq 12$ $= 12$			$u_1, v_1, v_2, y_1, u_1u_2,$ u_1v_2, u_2y_1 v_1, v_2, u_1v_1, y_1z_1

(2.8) $c(u_2) = c(x_1u_2) = c(z_1u_2) = 1$ and $c(x_1) = c(z_1) = 2$. We assign:

$w_H(u_2)$	$w_H(u_3)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
8	$\neq 6$ $= 6$			u_1v_2, v_1v_2, v_2y_1 u_2y_1, v_1v_2
9	$\neq 10$ $= 10$			$u_1, u_1u_2, u_1v_2,$ u_2y_1, v_1v_2, v_2y_1 $v_2, u_1u_2, u_1v_1, u_1v_2, v_2y_1$
10	$\neq 11$ $= 11$	2	$c(u_2u_3) = 1,$ $c(u_2) = 2$	$u_1, v_2, u_1u_2,$ u_2y_1, v_1v_2, v_2y_1 $v_2, u_1v_1, v_1v_2, v_2y_1$

(2.9) $c(x_1) = c(x_1u_2) = c(z_1u_2) = 1$ and $c(z_1) = c(u_2) = 2$. We assign:

$w_H(u_2)$	$w_H(u_3)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
7	$\neq 9$ $= 9$	1	$c(u_2u_3) = 2,$ $c(u_2) = 1$	$v_1, y_1, u_1u_2, u_1v_1, u_1v_2$ u_1, v_1, y_1, u_1u_2
8	$\neq 9$ $= 9$			$v_1, y_1, u_1u_2,$ u_1v_1, u_1v_2, v_1v_2 $u_1, v_1, y_1, u_1u_2,$ u_1v_1, u_1v_2
10	$\neq 10$ $= 10$			$u_1, y_1, u_1v_2,$ v_1v_2, v_1x_1, v_2y_1 $u_1, v_1, y_1, u_1u_2,$ u_1v_1, v_1v_2, v_2y_1

(2.10) $c(u_2) = c(z_1u_2) = c(z_1) = 1$ and $c(x_1) = c(x_1u_2) = 2$. We assign:

$w_H(u_2)$	$w_H(u_3)$	$w_H(x_1)$	Elements assigned with 2
8	$\neq 7$ $= 7$ $= 7$	$\neq 10$ $= 10$	u_2y_1, v_1x_1, v_2y_1 $v_2, u_1v_1, v_1x_1, v_2y_1$ $u_1, v_1, v_2, u_1v_1, u_1u_2, u_2y_1, v_1v_2$

$w_H(u_2)$	$w_H(z_1)$	Elements assigned with 2
9	$\neq 8$ $= 8$	$v_1, v_2, u_1u_2, u_1v_1, u_2y_1$ $u_1, u_1u_2, v_1x_1, v_2y_1$

$w_H(u_2)$	$w_H(u_3)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
10	$\neq 8$ $= 8$	2	$c(u_2u_3) = 1,$ $c(u_2) = 2$	$v_2, u_1v_1, u_2y_1, v_1x_1,$ v_2y_1 $v_1, v_2, u_1u_2, u_1v_1,$ u_2y_1, v_1v_2, v_2y_1

(2.11) $c(u_2) = c(x_1u_2) = c(z_1) = 1$ and $c(x_1) = c(z_1u_2) = 2$. We assign:

$w_H(u_2)$	$w_H(u_3)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
8	$\neq 7$ $= 7$			u_2y_1, v_1v_2, y_1z_1 $y_1, u_1u_2, u_2y_1, v_1v_2, v_2y_1$
9	$\neq 8$ $= 8$			$v_2, u_1u_2, u_1v_1,$ u_1v_2, v_2y_1, y_1z_1 $u_1, v_2, y_1, u_1v_2,$ u_2y_1, v_2y_1
10	$\neq 11$ $= 11$	2	$c(u_2u_3) = 1,$ $c(u_2) = 2$	$u_1, v_2, u_1u_2, u_1v_1,$ u_2y_1, v_2y_1, y_1z_1 $v_2, u_1v_1, v_1v_2,$ v_2y_1, y_1z_1

(2.12) $c(x_1) = c(z_1u_2) = c(z_1) = 1$ and $c(x_1u_2) = c(u_2) = 2$. We assign:

$w_H(u_2)$	$w_H(u_3)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
7	$\neq 8$ $= 8$	1	$c(u_2u_3) = 2,$ $c(u_2) = 1$	v_1, y_1, u_1u_2, v_1x_1 $v_1, y_1, u_1u_2, v_1x_1, v_2y_1$
$w_H(u_2)$	$w_H(u_3)$	Elements assigned with 2		
8	$\neq 9$ $= 9$	$v_1, v_2, y_1, u_1v_2, u_1u_2, v_1x_1$ $v_1, u_1v_1, v_1x_1, y_1z_1$		
9	$\neq 10$ $= 10$	$v_1, v_2, y_1, u_1u_2, u_1v_1, u_1v_2, v_1x_1, v_2y_1$ $u_1, v_1, y_1, u_1u_2, u_1v_2, u_2y_1, v_1v_2, v_1x_1, v_2y_1$		

(2.13) $c(x_1) = c(x_1u_2) = c(z_1) = 1$ and $c(z_1u_2) = c(u_2) = 2$. We assign:

$w_H(u_2)$	$w_H(u_3)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
7	$\neq 10$ $= 10$	1	$c(u_2u_3) = 2,$ $c(u_2) = 1$	$u_1, v_1, y_1, u_1u_2, y_1z_1$ $v_1, y_1, u_1u_2, u_1v_1,$ u_1v_2, y_1z_1
$w_H(u_2)$	$w_H(u_3)$	Elements assigned with 2		
8	$\neq 10$ $= 10$	$u_1, v_1, y_1, u_1u_2, v_1v_2, y_1z_1$ $v_1, y_1, u_1u_2, u_1v_1, v_1v_2, y_1z_1$		
9	$\neq 10$ $= 10$	$v_1, v_2, y_1, u_1u_2, u_1v_1, u_1v_2, v_1v_2, y_1z_1$ $u_1, y_1, u_1v_2, u_2y_1, v_1v_2, v_1x_1, y_1z_1$		

(2.14) $c(x_1) = c(u_2) = c(z_1u_2) = 1$ and $c(x_1u_2) = c(z_1) = 2$. We assign:

$w_H(u_2)$	$w_H(u_3)$	Elements assigned with 2		
8	$\neq 7$ $= 7$	u_2y_1, v_1v_2, v_1x_1 $u_1v_2, v_1v_2, v_1x_1, v_2y_1$		
9	$\neq 8$ $= 8$	$v_2, u_1u_2, u_1v_1, u_1v_2, v_1x_1, v_2y_1$ $u_1, u_1u_2, u_1v_2, u_2y_1, v_1v_2, v_1x_1, v_2y_1$		
$w_H(u_2)$	$w_H(u_3)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
10	$\neq 11$ $= 11$	2	$c(u_2u_3) = 1,$ $c(u_2) = 2$	$u_1, v_2, u_1u_2, u_1v_1,$ u_2y_1, v_1x_1, v_2y_1 $v_2, u_1v_1, u_2y_1, v_1v_2, v_1x_1$

(2.15) $c(x_1) = c(u_2) = c(x_1u_2) = 1$ and $c(z_1u_2) = c(z_1) = 2$. We assign:

$w_H(u_2)$	$w_H(u_3)$	Elements assigned with 2		
8	$\neq 9$ $= 9$	$u_1, y_1, u_1v_2, u_2y_1, v_1v_2$ $u_1, y_1, u_1u_2, u_2y_1, v_1v_2$		
9	$\neq 10$ $= 10$	$u_1, y_1, u_1v_1, u_1v_2, u_2y_1, v_1v_2, v_2y_1$ $v_2, u_1v_1, v_1v_2, y_1z_1$		
$w_H(u_2)$	$w_H(u_3)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
10	$\neq 11$ $= 11$	2	$c(u_2u_3) = 1,$ $c(u_2) = 2$	$u_1, v_2, u_1u_2, u_1v_1,$ u_2y_1, v_1v_2, y_1z_1 $u_1, v_2, u_1v_1, v_1v_2, y_1z_1$

(2.16) $c(x_1) = c(u_2) = c(z_1) = 1$ and $c(x_1u_2) = c(z_1u_2) = 2$. We assign:

$w_H(u_2)$	$w_H(u_3)$	Elements assigned with 2		
8	$\neq 6$ $= 6$	$u_1v_2, v_1v_2, v_1x_1, v_2y_1, y_1z_1$ $u_1, v_1, y_1, u_1u_2, u_1v_2, v_1v_2, u_2y_1, v_2y_1$		
9	$\neq 8$ $= 8$	$u_1v_1, u_2y_1, v_1v_2, v_1x_1, v_2y_1, y_1z_1$ $u_1, v_1, v_2, y_1, u_2y_1, v_1v_2, v_2y_1$		
$w_H(u_2)$	$w_H(u_3)$	Original $c(u_2u_3)$	Special rearrangement	Elements assigned with 2
10	$\neq 9$ $= 9$	2	$c(u_2u_3) = 1,$ $c(u_2) = 2$	$u_1, v_1, v_2, u_1u_2, u_1v_1,$ $u_1v_2, v_1v_2, v_2y_1, y_1z_1$ $v_2, u_1v_1, v_1v_2, v_1x_1,$ v_2y_1, y_1z_1

One may check that each total weighting c defined above is a nest total 2-weighting. This completes the proof. ■

References

- [1] J.A. Bondy, U.S.R. Murty, *Graph Theory with Applications*, Macmillan, London, 1976.
- [2] E. Flandrin, H. Li, A. Marczyk, J.-F. Saclé, M. Woźniak, A note on neighbor expanded sum distinguishing index, *Discuss. Math. Graph Theory*, **37** (2017), 29–37.
- [3] M. Kalkowski, A note on the 1,2-conjecture, (PhD Thesis, Adam Mickiewicz University in Poznań 2010), to appear in *Elect. J. Combin.*
- [4] M. Kalkowski, M. Karoński, F. Pfender, Vertex-coloring edge-weightings: towards the 1-2-3-conjecture. *J. Combin. Theory, Ser. B*, **100** (2010), 347–349.
- [5] M. Karoński, T. Łuczak, A. Thomason, Edge weights and vertex colours. *J. Combin. Theory, Ser. B*, **91** (2004), 151–157.
- [6] J. Przybyło, M. Woźniak, On a 1,2 conjecture, *Discrete Math. Theoret. Comput. Sci.*, **12**(1) (2010), 101–108.
- [7] W. Wang, K.-W. Lih, List coloring Halin graphs, *Ars Combin.*, **77** (2005), 53–63.