



The bounds of Full Friendly Index Sets of $C_m \times C_n$

Wai Chee SHIU and Man Ho LING

wcshiu@hkbu.edu.hk

Department of Mathematics,
Hong Kong Baptist University, Hong Kong
June 24-27, 2007

Outline

- ★ Labeling and induced labeling
- ★ Background and some known results
- ★ General properties
- ★ Upper bounds
- ★ Lower bounds
- ★ Further studies



Labeling and its induced labeling

Let $G = (V, E)$ be a connected, simple graph. Let A be a finite abelian group.



Labeling and its induced labeling

Let $G = (V, E)$ be a connected, simple graph. Let A be a finite abelian group.

A labeling $f : V \rightarrow A$ induces an edge labeling $f^* : E \rightarrow A$, defined by

$$f^*(xy) = f(x) + f(y),$$

for each $xy \in E$.

Labeling and its induced labeling

Let $G = (V, E)$ be a connected, simple graph. Let A be a finite abelian group.

A labeling $f : V \rightarrow A$ induces an edge labeling $f^* : E \rightarrow A$, defined by

$$f^*(xy) = f(x) + f(y),$$

for each $xy \in E$.

For each $i \in A$, let $v_f(i) = |f^{-1}(i)|$ and $e_f(i) = |f^{*-1}(i)|$.

A-cordial

A labeling is said to be *A-friendly* if $|v_f(i) - v_f(j)| \leq 1$ for all $i, j \in A$.

A-cordial

A labeling is said to be *A-friendly* if $|v_f(i) - v_f(j)| \leq 1$ for all $i, j \in A$.

An *A*-friendly labeling f is said to be *A-cordial* if $|e_f(i) - e_f(j)| \leq 1$ for all $i, j \in A$.

A-cordial

A labeling is said to be *A-friendly* if $|v_f(i) - v_f(j)| \leq 1$ for all $i, j \in A$.

An *A*-friendly labeling f is said to be *A-cordial* if $|e_f(i) - e_f(j)| \leq 1$ for all $i, j \in A$.

A graph G is *A-cordial* if there is an *A*-cordial labeling of G .

This concept was introduced by Hovey in 1991 which is a generalization of cordial graphs introduced by Cahit (when $A = \mathbb{Z}_2$).



Some known results in 1987

1987 Cahit ($A = \mathbb{Z}_2$):

1. Every tree is cordial.
2. K_n is cordial if and only if $n \leq 3$.
3. $K_{m,n}$ is cordial for all m, n .
4. Wheel $W_n = K_1 \vee C_{n-1}$ is cordial if and only if $n \not\equiv 0 \pmod{4}$.
5. C_n is cordial if and only if $n \not\equiv 2 \pmod{4}$.



Some known results in 1989-1999

1989 Lee, Ho and Shee

1. Characterized cordial generalized Petersen graphs completely.
2. Investigated unicyclic graphs.
3. Investigated the construction of cordial graphs by cartesian product and composition.



Some known results in 1989-1999

1989 Lee, Ho and Shee

1. Characterized cordial generalized Petersen graphs completely.
2. Investigated unicyclic graphs.
3. Investigated the construction of cordial graphs by cartesian product and composition.

1991 Lee and Liu investigated cordial complete k -partite graphs.



Some known results in 1989-1999

1989 Lee, Ho and Shee

1. Characterized cordial generalized Petersen graphs completely.
2. Investigated unicyclic graphs.
3. Investigated the construction of cordial graphs by cartesian product and composition.

1991 Lee and Liu investigated cordial complete k -partite graphs.

1997 Kuo, Chang and Kwong determined all m and n for which mK_n is cordial.



Some known results in 1989-1999

1989 Lee, Ho and Shee

1. Characterized cordial generalized Petersen graphs completely.
2. Investigated unicyclic graphs.
3. Investigated the construction of cordial graphs by cartesian product and composition.

1991 Lee and Liu investigated cordial complete k -partite graphs.

1997 Kuo, Chang and Kwong determined all m and n for which mK_n is cordial.

1999 Seoud and Abdel proved certain cylinder graphs are cordial.

Some known results in 2006-2007

2006, Chartrand, Lee and Zhang introduced the *friendly index set* of G :

$$\text{FI}(G) = \{|i_f(G)| \mid f \text{ is friendly}\},$$

where $i_f(G) = e_f(1) - e_f(0)$.



Some known results in 2006-2007

2006, Chartrand, Lee and Zhang introduced the *friendly index set* of G :

$$\text{FI}(G) = \{|i_f(G)| \mid f \text{ is friendly}\},$$

where $i_f(G) = e_f(1) - e_f(0)$.

2007, Lee and Ng determined the friendly index sets of complete bipartite graphs and cycles;

Some known results in 2006-2007

2006, Chartrand, Lee and Zhang introduced the *friendly index set* of G :

$$\text{FI}(G) = \{|i_f(G)| \mid f \text{ is friendly}\},$$

where $i_f(G) = e_f(1) - e_f(0)$.

2007, Lee and Ng determined the friendly index sets of complete bipartite graphs and cycles;

Kwong, Lee and Ng determined the friendly index sets of disjoint union of two cycles;

Some known results in 2006-2007

2006, Chartrand, Lee and Zhang introduced the *friendly index set* of G :

$$\text{FI}(G) = \{|i_f(G)| \mid f \text{ is friendly}\},$$

where $i_f(G) = e_f(1) - e_f(0)$.

2007, Lee and Ng determined the friendly index sets of complete bipartite graphs and cycles;

Kwong, Lee and Ng determined the friendly index sets of disjoint union of two cycles;

Salehi and Lee found the friendly index sets of paths.



Full friendly index sets

Now, Shiu and Kwong introduce the *full friendly index set* of G :

$$\text{FFI}(G) = \{i_f(G) \mid f \text{ is friendly}\}.$$



Full friendly index sets

Now, Shiu and Kwong introduce the *full friendly index set* of G :

$$\text{FFI}(G) = \{i_f(G) \mid f \text{ is friendly}\}.$$

Hence G is cordial if and only if $\{-1, 0, 1\} \cap \text{FFI}(G) \neq \emptyset$.



Full friendly index sets

Now, Shiu and Kwong introduce the *full friendly index set* of G :

$$\text{FFI}(G) = \{i_f(G) \mid f \text{ is friendly}\}.$$

The full friendly index sets of P_n and $P_2 \times P_n$ were determined.



Full friendly index sets

Now, Shiu and Kwong introduce the *full friendly index set* of G :

$$\text{FFI}(G) = \{i_f(G) \mid f \text{ is friendly}\}.$$

The full friendly index sets of P_n and $P_2 \times P_n$ were determined.

Namely

$$\text{FFI}(P_n) = \{2i + 1 - n \mid 1 \leq i \leq n - 1\},$$

Full friendly index sets

Now, Shiu and Kwong introduce the *full friendly index set* of G :

$$\text{FFI}(G) = \{i_f(G) \mid f \text{ is friendly}\}.$$

The full friendly index sets of P_n and $P_2 \times P_n$ were determined.

Namely

$$\text{FFI}(P_n) = \{2i + 1 - n \mid 1 \leq i \leq n - 1\},$$

$$\text{FFI}(P_2 \times P_{2n}) = \{2i + 2 - 6n \mid 2 \leq i \leq 6n - 2, i \neq 3, i \neq 6n - 3\},$$

Full friendly index sets

Now, Shiu and Kwong introduce the *full friendly index set* of G :

$$\text{FFI}(G) = \{i_f(G) \mid f \text{ is friendly}\}.$$

The full friendly index sets of P_n and $P_2 \times P_n$ were determined.

Namely

$$\text{FFI}(P_n) = \{2i + 1 - n \mid 1 \leq i \leq n - 1\},$$

$$\text{FFI}(P_2 \times P_{2n}) = \{2i + 2 - 6n \mid 2 \leq i \leq 6n - 2, i \neq 3, i \neq 6n - 3\},$$

$$\text{FFI}(P_2 \times P_{2n+1}) = \{2i - 1 - 6n \mid 3 \leq i \leq 6n + 1, i \neq 6n\},$$

General properties

For a fixed labeling f , a vertex v is called a *k -vertex* if $f(v) = k$ and an edge e is called a *k -edge* if $f^*(e) = k$.

General properties

For a fixed labeling f , a vertex v is called a **k -vertex** if $f(v) = k$ and an edge e is called a **k -edge** if $f^*(e) = k$.

Lemma 1 *If f is a labeling of a (p, q) -graph G , then $i_f(G) \leq q$.*

General properties

For a fixed labeling f , a vertex v is called a **k -vertex** if $f(v) = k$ and an edge e is called a **k -edge** if $f^*(e) = k$.

Lemma 1 *If f is a labeling of a (p, q) -graph G , then $i_f(G) \leq q$.*

Lemma 2 *An odd cycle C in a graph with labeling f contains at least one 0-edge.*

Vertical and horizontal cycles of $C_m \times C_n$




Figure 1: 5 vertical cycles and 3 horizontal cycles of $C_5 \times C_3$.

Upper bounds

Corollary 3 *If f is a (friendly) labeling of the graph $C_m \times C_n$, then $i_f(C_m \times C_n) \leq 2mn$.*

Upper bounds

Theorem 4 *If f is a (friendly) labeling of the graph $C_m \times C_n$, then $i_f(C_m \times C_n) \leq 2mn - 2m$ for even m and odd n .*

Upper bounds

Theorem 4 *If f is a (friendly) labeling of the graph $C_m \times C_n$, then $i_f(C_m \times C_n) \leq 2mn - 2m$ for even m and odd n .*

Proof: $C_m \times C_n$ contains at least m edge disjoint odd (vertical) cycles.

By Lemma 2, we have $e_f(0) \geq m$ and so

$e_f(1) \leq 2mn - m$. Hence,

$i_f(C_m \times C_n) \leq 2mn - 2m$. □

Note that $i_f(C_m \times C_n) \leq 2mn - 2n$ when m is odd and n is even. Therefore, we consider the case when m is even and n is odd only.

Upper bounds

Theorem 5 *If f is a (friendly) labeling of the graph $C_m \times C_n$, then $i_f(C_m \times C_n) \leq 2mn - 2m - 2n$ for odd m and n .*


Upper bounds

Theorem 5 *If f is a (friendly) labeling of the graph $C_m \times C_n$, then $i_f(C_m \times C_n) \leq 2mn - 2m - 2n$ for odd m and n .*

Proof: $C_m \times C_n$ contains at least $m + n$ edge disjoint odd cycles (m vertical and n horizontal). By Lemma 2, we have $e_f(0) \geq m + n$ and so $e_f(1) \leq 2mn - m - n$. Hence,
 $i_f(C_m \times C_n) \leq 2mn - 2m - 2n.$ □

Upper bounds are sharp


Example 1



$$i_f(C_6 \times C_4) = 48$$

Upper bounds are sharp


Example 1



$$i_f(C_6 \times C_3) = 24$$

Upper bounds are sharp

Example 1



$$i_f(C_5 \times C_3) = 14$$



Lower bounds

Let f be any labeling of a graph G .
Let a cycle C be a subgraph of G .

Lower bounds

Let f be any labeling of a graph G .

Let a cycle C be a subgraph of G .

C is called *mixing* (under f), if there is two vertices $u, v \in V(C)$ such that $f(u) = 1$ and $f(v) = 0$.

Clearly, a mixing cycle contains at least one 1-edge.

Lower bounds

Let f be any labeling of a graph G .

Let a cycle C be a subgraph of G .

C is called **mixing** (under f), if there is two vertices $u, v \in V(C)$ such that $f(u) = 1$ and $f(v) = 0$.

Clearly, a mixing cycle contains at least one 1-edge.

C is called **1-full cycle** (under f), if any vertex of C is 1-vertex.

C is called **0-full cycle** (under f), if any vertex of C is 0-vertex.

The bracket will be omitted if there is no ambiguity.

Lower bounds

Lemma 6 (Shiu and Kwong, 2007) *For any labeling, the number of 1-edges in a mixing cycle is even.*

Due to isomorphism, we may assume that $n \leq m$.

Lower bounds

Lemma 6 (Shiu and Kwong, 2007) *For any labeling, the number of 1-edges in a mixing cycle is even.*

Due to isomorphic, we may assume that $n \leq m$.

Theorem 7 *Let f be a friendly labeling of $C_m \times C_n$. If m is even, then $i_f(C_m \times C_n) \geq 4n - 2mn$.*



Lower bounds

Proof of Theorem 7

r : number of horizontal 1-full cycles

s : number of horizontal 0-full cycles

By the property of friendly labeling, $0 \leq r, s \leq \frac{n}{2}$.

Lower bounds

Proof of Theorem 7

r : number of horizontal 1-full cycles

s : number of horizontal 0-full cycles

Suppose $r = s = 0$.

$C_m \times C_n$ contains at least n edge disjoint mixing cycles.

Lower bounds

Proof of Theorem 7

r : number of horizontal 1-full cycles

s : number of horizontal 0-full cycles

Suppose either $r = 0$ or $s = 0$, without loss of generality, we may assume $r \neq 0$ and $s = 0$.

Lower bounds

Proof of Theorem 7

r : number of horizontal 1-full cycles

s : number of horizontal 0-full cycles

Suppose either $r = 0$ or $s = 0$, without loss of generality, we may assume $r \neq 0$ and $s = 0$.

\Rightarrow the number of horizontal mixing cycles is $n - r$,
and hence there exist at least $\lceil \frac{mn/2}{(n-r)} \rceil$ vertical mixing
cycles since $\frac{mn}{2}$ 0-vertices distribute in $n - r$
horizontal cycles.

Lower bounds

Proof of Theorem 7

r : number of horizontal 1-full cycles

s : number of horizontal 0-full cycles

Suppose either $r = 0$ or $s = 0$, without loss of generality, we may assume $r \neq 0$ and $s = 0$.

\Rightarrow the number of horizontal mixing cycles is $n - r$, and hence there exist at least $\lceil \frac{mn/2}{(n-r)} \rceil$ vertical mixing cycles since $\frac{mn}{2}$ 0-vertices distribute in $n - r$ horizontal cycles.

\Rightarrow at least

$$n - r + \lceil \frac{mn}{2(n-r)} \rceil \geq n - \frac{n}{2} + \frac{m}{2} \frac{n}{(n-r)} \geq \frac{m+n}{2} \geq n$$
 edge disjoint mixing cycles.

Lower bounds

Proof of Theorem 7

r : number of horizontal 1-full cycles

s : number of horizontal 0-full cycles

If $r \neq 0$ and $s \neq 0$, then there exist $m \geq n$ vertical mixing cycles.

Lower bounds

Proof of Theorem 7

r : number of horizontal 1-full cycles

s : number of horizontal 0-full cycles

For each case, the number of edge disjoint mixing cycles in $C_m \times C_n$ is at least n .

Lower bounds

Proof of Theorem 7

r : number of horizontal 1-full cycles

s : number of horizontal 0-full cycles


For each case, the number of edge disjoint mixing cycles in $C_m \times C_n$ is at least n .

By Lemma 6, we get $e_f(1) \geq 2n$ and $e_f(0) \leq 2mn - 2n$.

Hence, $i_f(C_m \times C_n) \geq 4n - 2mn$. □

Lower bounds is sharp


Example 2



$$i_f(C_6 \times C_4) = -32$$

Lower bounds is sharp

Example 2




$$i_f(C_6 \times C_3) = -24$$

Lower bounds

Theorem 8 *Let f be a friendly labeling of the graph $C_m \times C_n$. If m is odd, then*
$$i_f(C_m \times C_n) \geq 4n + 4 - 2mn.$$

Lower bounds are sharp


Example 3



$$i_f(C_5 \times C_4) = -20$$

Lower bounds are sharp


Example 3



$$i_f(C_5 \times C_3) = -14$$

Further studies

- ★ Determine the $\text{FFI}(C_m \times C_n)$
- ★ $P_m \times P_n$ for $m \geq 3$.
- ★ Unicyclic graphs.
- ★ Composition graphs, for example $C_m \circ N_n$.
- ★ Cylinder graph, i.e., $C_m \times P_n$.
- ★ Other products graphs.



END
Thank you