

Vol. 3, no. 1, 2003

ISSN 1311 – 6797

# INTERNATIONAL MATHEMATICAL JOURNAL

## Editorial Board

M. Amamiya ( Japan )	S. Kichenassamy ( France )
J. M. Ash ( USA )	F. – H. Lin ( USA )
T. Colding ( USA )	J. Lyness ( USA )
M. Fliess ( France )	K. Ono ( USA )
A. Fokas ( UK )	I. Prigogine ( Belgium )
K. Fuller ( USA )	L. Reichel ( USA )
J. Glimm ( USA )	P. C. Sabatier ( France )
W. Goldman ( USA )	M. Tang ( USA )
F. C. Graham ( USA )	W. Veech ( USA )
T. Hall ( Australia )	J. Vigo – Aguiar ( Spain )
D. Hong ( USA )	X. Zhou ( USA )

Managing Editor: Emil Minchev

International Scientific Publications

## *International Mathematical Journal*

**Aims and scopes:** The journal will publish carefully selected original research papers in the area of pure and applied mathematics as follows: combinatorics, order, lattices, ordered algebraic structures, general mathematical systems, number theory, field theory and polynomials, commutative rings and algebras, algebraic geometry, linear and multi linear algebra, matrix theory, associative and nonassociative rings, category theory, homological algebra, K – theory, group theory, Lie algebras, real functions, measure and integration, functions of complex variables, potential theory, special functions, ordinary differential equations, partial differential equations, finite difference and functional equations, approximation and expansion, Fourier analysis, abstract harmonic analysis, integral transforms, operational calculus, integral equations, functional analysis, operator theory, calculus of variation and optimal control, optimization, geometry, differential geometry, topology, manifolds, probability theory and stochastic processes, statistics, numerical analysis, computer science, mechanics, thermodynamics, quantum theory, relativity, economics, programming, games, mathematical biology, information and communication, circuits.

**Call for papers:** The authors are cordially invited to submit papers in triplicate to the Managing Editor: Emil Minchev. Manuscripts submitted to this journal will be considered for publication with the understanding that the same work has not been published and is not under consideration for publication elsewhere.

**Managing Editor:** Dr. Emil Minchev, Department of Mathematics, Faculty of Education, Chiba University, Yayoi – cho 1 – 33, Inage – ku, Chiba 263 – 8522, Japan

e – mail: eminchev@hotmail.com

© 2003 Intern. Math. Journal. All rights reserved.

ISSN: 1523-9032 (print) ISSN: 1523-9032 (electronic)

Volume 3 Number 1, March 2003  
Editor-in-Chief: Dr. Peter C.B. Lam, Department of Mathematics, Hong Kong Baptist University, Kowloon Tong, Hong Kong.

Volume 3 Number 2, June 2003  
Editor-in-Chief: Dr. Peter C.B. Lam, Department of Mathematics, Hong Kong Baptist University, Kowloon Tong, Hong Kong.

Volume 3 Number 3, September 2003  
Editor-in-Chief: Dr. Peter C.B. Lam, Department of Mathematics, Hong Kong Baptist University, Kowloon Tong, Hong Kong.

Volume 3 Number 4, December 2003  
Editor-in-Chief: Dr. Peter C.B. Lam, Department of Mathematics, Hong Kong Baptist University, Kowloon Tong, Hong Kong.

## More on the Generalized Fibonacci Numbers and Associated Bipartite Graphs

W.C. Shiu and Peter C.B. Lam

Department of Mathematics, Hong Kong Baptist University,  
224 Waterloo Road, Kowloon Tong, Hong Kong.

### Abstract

For a positive integer  $k \geq 2$ , the  $k$ -Fibonacci sequence  $\{f_n^{(k)}\}$  is defined by  $f_n^{(k)} = f_{n-1}^{(k)} + f_{n-2}^{(k)} + \cdots + f_{n-k}^{(k)}$ , for  $n \geq k$ , with initial value  $f_0^{(k)} = f_1^{(k)} = \cdots = f_{k-2}^{(k)} = 0$ ,  $f_{k-1}^{(k)} = 1$ . For a fixed  $\alpha = (a_1, a_2, \dots, a_m)$ , the  $(k, \alpha)$ -sequence is defined by  $s(\alpha)_n^{(k)} = \sum_{i=1}^m a_i f_{n-1+k-i}^{(k)}$  for  $k \geq 2$ ,  $m \geq 1$  and  $n \geq 1$ . In this paper, we consider the relationship between  $s(\alpha)_n^{(k)}$  and perfect matchings of a bipartite graph.

**AMS 2000 MSC** : 15A15, 05C50, 05C30

**Keywords** : Permanent,  $k$ -Fibonacci sequence,  $(k, \alpha)$ -Fibonacci sequence, perfect matching

### 1. Introduction

Let  $A = (a_{i,j})$  be a square matrix of order  $n$  over a ring  $R$ . The *permanent* of  $A$  is defined by

$$\text{per}(A) = \sum_{\sigma \in S_n} \prod_{i=1}^n a_{i,\sigma(i)},$$

where  $S_n$  denotes the symmetric group on  $n$  letters. It is easy to see that for any square matrix  $A$  and any permutation matrices  $P$  and  $Q$ ,  $\text{per}(A) = \text{per}(PAQ)$ . Let

$A_{i,j}$  be the matrix obtained from a square matrix  $A = (a_{i,j})$  by deleting the  $i$ -th row and the  $j$ -th column. Then it is also easy to see that  $\text{per}(A) = \sum_{k=1}^n a_{i,k} \text{per}(A_{i,k}) = \sum_{k=1}^n a_{k,j} \text{per}(A_{k,j})$  for any  $i, j$ .

In this paper, all undefined terminologies and symbols of graph can be found in [1]. Let  $G$  be a bipartite graph with bipartition  $(X, Y)$ . If  $G$  contains a perfect matching, then  $|X| = |Y|$ . Let  $A$  be an adjacency matrix of  $G$ . It is known [7] that the number of perfect matchings (or 1-factor) of  $G$  is  $\sqrt{\text{per}(A)}$ . Namely, if  $|X| = |Y| = n$  then  $A = \begin{pmatrix} O & B \\ B^T & O \end{pmatrix}$  for some square matrix  $B$  of order  $n$ , where  $O$  is the zero matrix of order  $n$ . Such matrix  $B$  is called a *bipartite adjacency matrix*. We shall denote the graph  $G$  as  $G(B)$ . Note that the matrix  $B$  is not unique. The number of perfect matchings of  $G(B)$  is  $\text{per}(B)$ , see [7].

Let  $\{F_n\}$  be the Fibonacci sequence, i.e.,  $F_0 = 0$ ,  $F_1 = 1$  and  $F_n = F_{n-1} + F_{n-2}$  for  $n \geq 2$ .

The  $k$ -Fibonacci sequence  $\{f_n^{(k)}\}$  for positive integer  $k \geq 2$  is defined recursively by

$$f_n^{(k)} = f_{n-1}^{(k)} + f_{n-2}^{(k)} + \cdots + f_{n-k}^{(k)}, \text{ for } n \geq k,$$

with initial value  $f_0^{(k)} = f_1^{(k)} = \cdots = f_{k-2}^{(k)} = 0$ ,  $f_{k-1}^{(k)} = 1$ . The number  $f_n^{(k)}$  is called the  $n$ -th  $k$ -Fibonacci number. It is known that [6]

$$f_j^{(k)} = 2^{j-k}, \text{ for } k \leq j \leq 2k-1.$$

Note that  $\{f_n^{(2)}\}$  is the Fibonacci sequence.

The  $k$ -Lucas sequence  $\{l_n^{(k)}\}$  is defined by  $l_n^{(k)} = f_{n-1}^{(k)} + f_{n+k-1}^{(k)}$  and  $l_n^{(k)}$  is called the  $n$ -th  $k$ -Lucas number. It is known that  $l_j^{(k)} = 2^{j-1}$ ,  $1 \leq j \leq k-1$ , and  $l_k^{(k)} = 1 + 2^{k-1}$ , see [6]. More about Lucas sequence can be found in [3]. Note that  $\{l_n^{(2)}\}$  is the Lucas sequence.

A matrix is said to be a  $(0, 1)$ -matrix if each of its entries is either 0 or 1. Suppose  $n$  and  $k$  are positive integers. Let  $T_n = (t_{i,j})$  be an  $n \times n$  tridiagonal  $(0, 1)$ -matrix, where  $t_{i,j} = 1$  if and only if  $|j-i| \leq 1$ . Let  $U_n^{(k)} = (u_{i,j})$  be an  $n \times n$  upper triangular  $(0, 1)$ -matrix, where  $u_{i,j} = 1$  if and only if  $2 \leq j-i \leq k-1$  if  $k \leq n$  and  $U_n^{(k)} = U_n^{(n)}$  if  $k > n$ . Let  $\mathcal{F}^{(n,k)} = T_n + U_n^{(k)}$  and let  $\mathcal{C}^{(n,k)} = \mathcal{F}^{(n,k)} + E_{1,k+1} - \sum_{j=2}^k E_{1,j}$  for  $n \geq 3$ , where  $E_{i,j}$  denotes the  $n \times n$  matrix with 1 at the  $(i, j)$ -th entry and zeros elsewhere.

In [4, 5], Lee et al. found a class of bipartite graphs whose number of perfect matchings is  $f_n^{(k)}$  and prove the following result.

**Theorem 1.1:** For  $n \geq 2$ , the number of perfect matchings of  $G(\mathcal{F}^{(n,k)})$  is  $f_{n-1+k}^{(k)}$ .

In [2], Brualdi proved the following result:

**Theorem 1.2:** *For  $n \geq 2$ , let  $A^{(n)} = I + U^{(n)}$ . Then  $\text{per}(A^{(n)}) = 2^{n-1}$ .*

Making use of Theorem 1.2, Lee [6] proved the following result.

**Theorem 1.3:** *For  $n \geq 3$ , the number of perfect matchings of  $G(\mathcal{C}^{(n,k)})$  is  $l_{n-1}^{(k)}$ .*

In this paper, we shall show in Section 2 that the permanent of a special matrix is a linear combination of  $k$ -Fibonacci numbers. By making use of this result we obtain the number of perfect matchings of a larger class of bipartite graphs. Theorems 1.1 to 1.3 are special cases of this result. Moreover, in Section 3 we shall use this permanent to obtain the number of perfect matchings of certain bipartite graph which is not isomorphic to the graphs studied in [6].

## 2. Main results

For a fixed  $\alpha = (a_1, a_2, \dots, a_m) \in R^m$ , where  $R$  is a ring. We define the  $(k, \alpha)$ -sequence by

$$s(\alpha)_n^{(k)} = a_1 f_{n+k-2}^{(k)} + a_2 f_{n+k-3}^{(k)} + \cdots + a_m f_{n+k-m-1}^{(k)} = \sum_{i=1}^m a_i f_{n-1+k-i}^{(k)}, \quad k \geq 2, n \geq 1.$$

The number  $s(\alpha)_n^{(k)}$  is called the  $n$ -th  $(k, \alpha)$ -number. Note that, if  $\alpha = (1, \dots, 1) \in \mathbb{Z}^k$ , then  $s(\alpha)_n^{(k)}$  is the  $(n-1+k)$ -th  $k$ -Fibonacci number  $f_{n-1+k}^{(k)}$ ; if  $\alpha = (1, 0, \dots, 0, 1) \in \mathbb{Z}^{k+1}$ , then  $s(\alpha)_n^{(k)}$  is the  $(n-1)$ -st  $k$ -Lucas number  $l_{n-1}^{(k)}$ .

**Theorem 2.1:** *Suppose  $n, k \geq 2$ . Let*

$$B_n = \left( \begin{array}{c|cccc} a_1 & a_2 & \cdots & a_n \\ \hline 1 & & & & \\ 0 & & & & \\ \vdots & & & \mathcal{F}^{(n-1,k)} & \\ 0 & & & & \end{array} \right),$$

for some elements  $a_1, a_2, \dots, a_n$  in a ring  $R$ . Then  $\text{per}(B_n) = \sum_{i=1}^n a_i f_{n-1+k-i}^{(k)}$ .

**Proof:** We shall prove the theorem by mathematical induction on  $n$ . Since

$$\text{per}(B_2) = a_1 + a_2 = a_1 f_k^{(k)} + a_2 f_{k-1}^{(k)},$$

the theorem is true for  $n = 2$ .

Assume that the theorem is true for some  $n \geq 2$ . Expanding the permanent by the first column and by Theorem 1.1 and the induction assumption, we have

$$\begin{aligned}
\text{per}(B_{n+1}) &= a_1 \text{per}(\mathcal{F}^{(n,k)}) + \text{per} \left( \begin{array}{c|cccc} a_2 & a_3 & \cdots & a_{n+1} \\ \hline 1 & & & & \\ 0 & & & & \\ \vdots & & & & \\ 0 & & & & \mathcal{F}^{(n-1,k)} \end{array} \right) \\
&= a_1 f_{n+k-1} + \sum_{i=1}^n a_{i+1} f_{n-1+k-i}^{(k)} \\
&= a_1 f_{n+k-1} + \sum_{i=2}^{n+1} a_i f_{n+k-i}^{(k)} = \sum_{i=1}^{n+1} a_i f_{n+k-i}^{(k)}.
\end{aligned}$$

Thus, the theorem is true for each  $n \geq 2$ . ■

**Corollary 2.2:** For a fixed  $m \geq 1$ , suppose  $n, k \geq 2$  and  $n \geq m$ . Let

$$\mathcal{S}_\alpha^{(n,k)} = \left( \begin{array}{c|ccccc} a_1 & a_2 & \cdots & a_m & 0 & \cdots 0 \\ \hline 1 & & & & & \\ 0 & & & & & \mathcal{F}^{(n-1,k)} \\ \vdots & & & & & \\ 0 & & & & & \end{array} \right).$$

Then the number of perfect matching of  $G(\mathcal{S}_\alpha^{(n,k)})$  is the  $n$ -th  $(k, \alpha)$ -number with  $\alpha = (a_1, a_2, \dots, a_m)$ .

Applying Corollary 2.2 and Theorem 2.1 by choosing  $\alpha = (1, 1, \dots, 1) \in \mathbb{Z}^k$  for  $n > k$  and by choosing  $a_i = 1$  for all  $i = 1, 2, \dots, n$  for  $n \leq k$  respectively, we get Theorem 1.1. Applying Corollary 2.2 and Theorem 2.1 by choosing  $\alpha = (1, 0, \dots, 0, 1) \in \mathbb{Z}^{k+1}$  for  $n > k$  and by choosing  $a_1 = 1$  and  $a_i = 0$  for all  $i = 2, \dots, n$  for  $n \leq k$  respectively, we get Theorem 1.3.

### 3. Other results

From Theorem 1.3, the number of perfect matchings of  $G(\mathcal{C}^{(n,2)})$  is  $l_{n-1}^{(2)}$ . In [6], there is a bipartite graph  $G$ , which is not isomorphic to  $G(\mathcal{C}^{(n,2)})$  and whose number of perfect matchings is also  $l_{n-1}^{(2)}$ . Namely  $G = G(B^{(n)})$ , where  $B^{(n)} = T_n + E_{1,3} - E_{2,3} + E_{2,4} - E_{3,4}$  for  $n \geq 4$ . Now we shall show another bipartite graph whose number of perfect matchings is  $l_{n-1}^{(2)}$  too.

Let  $C^{(n)} = T_n - E_{2,3} + E_{1,5}$  for  $n \geq 5$ . It is easy to see that both  $G(C^{(n)})$  and  $G(\mathcal{C}^{(n,2)})$  contain exactly one vertex of degree 4 when  $n \geq 6$ . It is easy to show that  $G(C^{(6)})$  is not isomorphic to  $G(\mathcal{C}^{(6,2)})$ . Let  $a$  and  $b$  be the vertices of degree 4 in  $G(C^{(n)})$  and  $G(\mathcal{C}^{(n,2)})$  respectively. For  $n \geq 7$ , since  $b$  is adjacent to a vertex of degree 2 but  $a$  is not,  $G(C^{(n)})$  is not isomorphic to  $G(\mathcal{C}^{(n,2)})$ . Since  $G(B^{(n)})$  does not contain any vertex of degree 4 when  $n \geq 6$ ,  $G(C^{(n)})$  is not isomorphic to  $G(B^{(n)})$ . Note that  $G(C^{(5)})$  is isomorphic to  $G(B^{(5)})$ .

Expanding the permanent by the first column and by Theorem 2.1 we have

$$\begin{aligned}
 \text{per}(C^{(n)}) &= \text{per} \left( \begin{array}{cc|ccccc} 1 & 1 & 0 & 0 & 1 & 0 & 0 & \cdots \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & \cdots \\ 0 & 1 & 1 & 1 & 0 & 0 & 0 & \cdots \\ 0 & 0 & 1 & 1 & 1 & 0 & 0 & \cdots \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & \cdots \\ \hline 0 & 0 & 0 & 0 & 1 & & & \\ 0 & 0 & 0 & 0 & 0 & \mathcal{F}^{(n-5,k)} & & \\ \vdots & \vdots & \vdots & \vdots & \vdots & & & \end{array} \right) \\
 &= \text{per} \left( \begin{array}{c|cccc} 1 & 0 & 0 & \cdots \\ \hline 1 & & & \\ 0 & \mathcal{F}^{(n-2,k)} & & \\ \vdots & & & \end{array} \right) + \text{per} \left( \begin{array}{c|ccccc} 1 & 0 & 0 & 1 & 0 & \cdots \\ \hline 1 & & & & & \\ 0 & & & \mathcal{F}^{(n-2,k)} & & \\ \vdots & & & & & \end{array} \right) \\
 &= f_{n-1}^{(2)} + f_{n-1}^{(2)} + f_{n-4}^{(2)} = f_{n-1}^{(2)} + f_{n-2}^{(2)} + f_{n-3}^{(2)} + f_{n-4}^{(2)} = f_n^{(2)} + f_{n-2}^{(2)} = l_{n-1}^{(2)}
 \end{aligned}$$

Thus the number of perfect matchings of the bipartite graph  $G(C^{(n)})$  is  $l_{n-1}^{(2)}$ .

## References

- [1] J.A. Bondy and U.S.R. Murty, *Graph Theory with Applications*, Macmillan (1976).
- [2] R.A.Brualdi, An interesting face of the polytope of doubly stochastic matrices, *Linear Algebra Appl.*, **17** (1985), 5-18.
- [3] R. Honsberger, *Mathematical Germ III*, Math. Assoc. Amer. (1985).
- [4] G.Y. Lee and S.G. Lee, A note on generalized Fibonacci numbers, *Fibonacci Quarterly*, **33(3)** (1995), 273-278.
- [5] G.Y. Lee, S.G. Lee and H.G. Shin, On the  $k$ -generalized Fibonacci matrix  $Q_k$ , *Linear Algebra and its Applications*, **251** (1997), 73-88.
- [6] G.Y. Lee,  $k$ -Lucas numbers and associated bipartite, *Linear Algebra and its Applications*, **320** (2000), 51-61.
- [7] H. Minc, *Permanents*, *Encyclopedia of Mathematics and its Applications*, Addison-Wesley, New York (1978).

Received: August 21, 2002

## Contents

A. Cossidente, V. Napolitano, <i>On some complete intersections in <math>PG(3, q)</math></i>	1
W. C. Shiu, Peter C. B. Lam, <i>More on the generalized Fibonacci numbers and associated bipartite graphs</i> .....	5
E. Ballico, <i>Special projections of projective varieties</i> .....	11
E. Ballico, A. Cossidente, <i>Polar hypersurfaces and projective curves</i>	13
S. Mecheri, <i>Orthogonality and derivation ranges</i> .....	21
S. Mecheri, <i>Another version of Anderson's inequality in norm – ideal</i>	29
R. A. Mollin, K. Cheng, <i>Matrices and continued fractions</i> .....	41
Shu – Cherng Fang, Guangzhi Li, <i>On the resolution of finite fuzzy relation equations</i> .....	59
K. B. Kulasekera, C. Gallagher, <i>Comparisons of two and three dimensional regression surfaces</i> .....	73