

Group Magicness of Complete N -partite Graphs

W.C. Shiu

Department of Mathematics, Hong Kong Baptist University
224 Waterloo Road, Kowloon Tong, Hong Kong
wcshiu@hkbu.edu.hk

Richard M. Low*

Department of Mathematics, San Jose State University
San Jose, CA 95192 USA
low@math.sjsu.edu

Abstract

Let A be a non-trivial abelian group. We call a graph $G = (V, E)$ A -magic if there exists a labeling $f : E(G) \rightarrow A \setminus \{0\}$ such that the induced vertex set labeling $f^+ : V(G) \rightarrow A$, defined by $f^+(v) = \sum f(u, v)$ where the sum is over all $(u, v) \in E(G)$, is a constant map. In this paper, we show that K_{k_1, k_2, \dots, k_n} ($k_i \geq 2$) is A -magic, for all A where $|A| \geq 3$.

1 Introduction.

Let $G = (V, E)$ be a connected, simple graph. For any nontrivial abelian group A (written additively), let $A^* = A \setminus \{0\}$. A function $f : E(G) \rightarrow A^*$ is called a *labeling* of G . Any such labeling induces a map $f^+ : V(G) \rightarrow A$, defined by $f^+(v) = \sum f(u, v)$ where the sum is over all $(u, v) \in E(G)$. If there exists a labeling f whose induced map on $V(G)$ is a constant map, we say that f is an A -magic labeling and that G is an A -magic graph. The *integer-magic spectrum* of a graph G is the set $\text{IM}(G) = \{k : G \text{ is } \mathbb{Z}_k\text{-magic and } k \geq 1\}$. By convention, \mathbb{Z} -magic graphs are considered to be \mathbb{Z}_1 -magic.

\mathbb{Z} -magic graphs were considered by Stanley [19,20], where he pointed out that the theory of magic labelings could be studied in the general context of linear homogeneous diophantine equations. Doob [1,2,3] and others [7,8,14] have studied A -magic graphs and \mathbb{Z}_k -magic graphs were investigated in [4,6,9,10,11,12,13].

*This paper was presented at the 18th Midwest Conference on Combinatorics, Cryptography, and Computing; Rochester, New York (October 29, 2004).

Within the mathematical literature, various definitions of magic graphs have been introduced. The original concept of an A -magic graph is due to J. Sedláček [15,16], who defined it to be a graph with real-valued edge labeling such that (i) distinct edges have distinct nonnegative labels, and (ii) the sum of the labels of the edges incident to a particular vertex is the same for all vertices. Previously, Kotzig and Rosa [5] had introduced yet another definition of a magic graph. Over the years, there has been great research interest in graph labeling problems. The interested reader is directed to Wallis' [21] recent monograph on magic graphs.

2 Basic definitions and notation.

In the study of edge-magic labelings, Shiu, Lam and Lee [17] introduced the following notation. Suppose $f : E(G) \rightarrow X$ is a mapping (i.e., an edge labeling of G). The *labeling matrix* for f , denoted by $\mathcal{L}_f(G)$, is the matrix whose rows and columns are named by the vertices of G and defined in the following way: the (u, v) -entry is $f(uv)$ if $uv \in E$, and is $*$ otherwise. If f is an A -magic labeling of G , then $\mathcal{L}_f(G)$ is an A -magic labeling matrix of G . Note that the row sum of an A -magic labeling matrix is the A -magic value corresponding to the labeling f .

Thus, finding an A -magic labeling of G is equivalent to finding an A -magic labeling matrix $\mathcal{L}_f(G)$, where each row sum (as well as column sum) is the same constant value. In the context of row and column sums, entries with an $*$ are treated as 0.

A graph is called *fully magic* if it is A -magic, for every abelian group A . A graph is called *non-magic* if for every abelian group A , it is not A -magic.

In this paper, we analyze the group-magicness property for the class of complete n -partite graphs.

3 Main results.

First, let us make a few observations. They are straight-forward to verify and can be found in [14].

Observations:

1. A graph G is \mathbb{Z}_2 -magic iff every vertex of G is of the same parity.
2. An eulerian graph G having an even number of edges is A -magic.
3. If A_1 is a subgroup of A and graph G is A_1 -magic, then G is A -magic.

We now characterize the abelian groups A , for which $K_{m,n}$ is A -magic. Let f be a labeling of the complete bipartite graph $K_{m,n}$. Then, $\mathcal{L}_f(K_{m,n}) = \begin{pmatrix} \star_m & B \\ B^T & \star_n \end{pmatrix}$, where B is an $m \times n$ matrix, \star_m and \star_n are square matrices of order m and n respectively with all entries being $*$.

Theorem 1. Let m and n be even. Then, $K_{m,n}$ has an A -magic labeling with magic value 0, for all A .

Proof. Suppose $a \in A \setminus \{0\}$. Let S be an $m \times n$ matrix defined by $S_{i,j} = (-1)^{i+j}a$, where $S_{i,j}$ denotes the (i,j) -entry of S . Then, the row sums and the column sums of S are zero. Clearly, $\begin{pmatrix} \star_m^m & S \\ S^T & \star_n^n \end{pmatrix}$ is an A -magic labeling matrix of $K_{m,n}$, with A -magic value 0. \square

Definition. The matrix S defined in the proof above is called an $m \times n$ zero-sum $(a, -a)$ -matrix.

The integer-magic spectrum of $K_{1,n}$ has been found [11]. For convenience, we state the result here.

Theorem B. $K_{1,1}$ is fully magic and $K_{1,2}$ is non-magic. For $n \geq 3$,

$$\text{IM}(K_{1,n}) = \bigcup_{p|(n-1)} p\mathbb{N}.$$

It is straight-forward to verify the following lemma.

Lemma 1. For $n \geq 3$, $K_{1,n}$ is V_4 -magic if and only if n is odd.

So we may assume $m \geq 3$.

Lemma 2. Let A be an abelian group of order at least 3. Then, there exist $a, b, c \in A \setminus \{0\}$ (not necessarily distinct) such that $a + b + c = 0$.

Proof. It suffices to consider three cases, namely: $A = \mathbb{Z}, \mathbb{Z}_k$ for $k \geq 3$, or V_4 .

If $A = \mathbb{Z}$, then it is obvious. If $A = \mathbb{Z}_k$, then choose $a = b = 1$ and $c = -2$. If $A = V_4$, then choose $a = (1, 0)$, $b = (0, 1)$ and $c = (1, 1)$. \square

Theorem 3. Suppose m is odd, with $m \geq 3$ and $n \geq 2$. For any abelian group A where $|A| \geq 3$, $K_{m,n}$ has an A -magic labeling with magic value 0.

Proof. Let $a, b, c \in A \setminus \{0\}$ be chosen in the same manner as discussed in the proof of Lemma 2.

CASE 1. n is even.

Let $B = \begin{pmatrix} C \\ D \end{pmatrix}$, where C is an $(m-3) \times n$ zero-sum $(a, -a)$ -matrix and D is a $3 \times n$ matrix defined by

$$D_{i,j} = \begin{cases} (-1)^j a & \text{if } i = 1; \\ (-1)^j b & \text{if } i = 2; \\ (-1)^j c & \text{if } i = 3. \end{cases}$$

Note that if $m = 3$, then C does not appear. Then, $\mathcal{L}_f(K_{m,n}) = \begin{pmatrix} \star_m^m & B \\ B^T & \star_n^n \end{pmatrix}$ is an A -magic labeling matrix of $K_{m,n}$, for $A = \mathbb{Z}, \mathbb{Z}_k$ ($k \geq 3$), and V_4 . By

Observation 3, $K_{m,n}$ is A -magic, for all A where $|A| \geq 3$.

CASE 2. n is odd.

Then, $n \geq 3$. Let B be a matrix of the following form:

$$B = \begin{pmatrix} C_1 & D_1^T \\ D_1 & E \end{pmatrix},$$

where C_1 is an $(m-3) \times (n-3)$ zero-sum $(a, -a)$ -matrix, D_1 is a $3 \times (n-3)$ matrix defined similarly as in the proof of the previous case, and E is a Latin square of order 3 defined as follows:

$$E = \begin{pmatrix} a & b & c \\ c & a & b \\ b & c & a \end{pmatrix}.$$

Then, $\mathcal{L}_f(K_{m,n}) = \begin{pmatrix} \star_m^m & B \\ B^T & \star_n^n \end{pmatrix}$ is an A -magic labeling matrix of $K_{m,n}$, for $A = \mathbb{Z}, \mathbb{Z}_k$ ($k \geq 3$), and V_4 . By Observation 3, $K_{m,n}$ is A -magic, for all A where $|A| \geq 3$.

It is clear that the row sums and the column sums of these A -magic labeling matrices are zero. \square

Here a few examples which illustrate Theorem 3.

EXAMPLE 1. $m = 3$ and $n = 4$. Then,

$$B = \begin{pmatrix} -a & a & -a & a \\ -b & b & -b & b \\ -c & c & -c & c \end{pmatrix}, \text{ and } \mathcal{L}_f(K_{m,n}) = \begin{pmatrix} \star_3^3 & B \\ B^T & \star_4^4 \end{pmatrix}.$$

EXAMPLE 2. $m = 3$ and $n = 3$. Then,

$$B = \begin{pmatrix} a & b & c \\ c & a & b \\ b & c & a \end{pmatrix}, \text{ and } \mathcal{L}_f(K_{m,n}) = \begin{pmatrix} \star_3^3 & B \\ B^T & \star_3^3 \end{pmatrix}.$$

EXAMPLE 3. $m = 5$ and $n = 4$. Then,

$$B = \begin{pmatrix} a & -a & a & -a \\ -a & a & -a & a \\ -a & a & -a & a \\ -b & b & -b & b \\ -c & c & -c & c \end{pmatrix}, \text{ and } \mathcal{L}_f(K_{m,n}) = \begin{pmatrix} \star_5^5 & B \\ B^T & \star_4^4 \end{pmatrix}.$$

EXAMPLE 4. $m = 5$ and $n = 2$. Then,

$$B = \begin{pmatrix} a & -a \\ -a & a \\ -a & a \\ -b & b \\ -c & c \end{pmatrix}, \text{ and } \mathcal{L}_f(K_{m,n}) = \begin{pmatrix} \star_5^5 & B \\ B^T & \star_2 \end{pmatrix}.$$

We conclude by showing that K_{k_1, k_2, \dots, k_n} ($k_i \geq 2$) is A -magic, for all A where $|A| \geq 3$. First, recall the following definitions and notation.

Definition. A graph G is n -partite, $n \geq 1$, if it is possible to partition $V(G)$ into n subsets V_1, V_2, \dots, V_n such that every element of $E(G)$ joins a vertex of V_i to a vertex of V_j , $i \neq j$.

Definition. A complete n -partite graph G is an n -partite graph with partite sets V_1, V_2, \dots, V_n having the added property that if $u \in V_i$ and $v \in V_j$, $i \neq j$, then $uv \in E(G)$.

Notation. A complete n -partite graph G with partite sets V_1, V_2, \dots, V_n , where $|V_i| = k_i$, is denoted by K_{k_1, k_2, \dots, k_n} .

We now establish the following result.

Theorem 4. For $n \geq 2$, the complete n -partite graph K_{k_1, k_2, \dots, k_n} with $k_i \geq 2$, is A -magic, for all A where $|A| \geq 3$.

Proof. There are $\binom{n}{2}$ ways of choosing a pair from the partite sets V_1, V_2, \dots, V_n . For each pair, apply a labeling on the corresponding edge set, using either Theorem 1 or Theorem 3. \square

References

- [1] M. Doob, *On the construction of magic graphs*, Proc. Fifth S.E. Conference on Combinatorics, Graph Theory and Computing (1974), 361-374.
- [2] M. Doob, *Generalizations of magic graphs*, Journal of Combinatorial Theory, Series B, **17** (1974), 205-217.
- [3] M. Doob, *Characterizations of regular magic graphs*, Journal of Combinatorial Theory, Series B, **25** (1978), 94-104.
- [4] M.C. Kong, S-M Lee, and H. Sun, *On magic strength of graphs*, Ars Combinatoria, **45** (1997), 193-200.

- [5] A. Kotzig and A. Rosa, *Magic valuations of finite graphs*, Canad. Math. Bull., **13** (1970), 451–461.
- [6] S-M Lee, Yong-Song Ho and R.M. Low, *On the integer-magic spectra of maximal planar and maximal outerplanar graphs*, preprint.
- [7] S-M Lee, A. Lee, Hugo Sun, and Ixin Wen, *On group-magic graphs*, Journal of Combinatorial Mathematics and Combinatorial Computing, **38** (2001), 197–207.
- [8] S-M Lee, F. Saba, E. Salehi, and H. Sun, *On the V_4 -group magic graphs*, Congressus Numerantium, **156** (2002), 59–67.
- [9] S-M Lee, F. Saba, and G. C. Sun, *Magic strength of the k -th power of paths*, Congressus Numerantium, **92** (1993), 177–184.
- [10] S-M Lee and E. Salehi, *Integer-magic spectra of amalgamations of stars and cycles*, Ars Combinatoria, **67** (2003), 199–212.
- [11] S-M Lee, E. Salehi and H. Sun, *Integer-magic spectra of trees with diameters at most four*, preprint.
- [12] S-M Lee, L. Valdes, and Yong-Song Ho, *On group-magic spectra of trees, double trees and abbreviated double trees*, Journal of Combinatorial Mathematics and Combinatorial Computing, **46** (2003), 85–96.
- [13] S-M Lee and J. Wang, *On the integer-magic spectra of honeycomb graphs*, preprint.
- [14] R.M. Low and S-M Lee, *On group-magic eulerian graphs*, Journal of Combinatorial Mathematics and Combinatorial Computing, **50** (2004), 141–148.
- [15] J. Sedláček, *On magic graphs*, Math. Slov., **26** (1976), 329–335.
- [16] J. Sedláček, *Some properties of magic graphs*, in Graphs, Hypergraph, and Bloc Syst. 1976, Proc. Symp. Comb. Anal., Zielona Gora (1976), 247–253.
- [17] W.C. Shiu, P.C.B. Lam and S-M Lee, *Edge-magicness of the composition of a cycle with a null graph*, Congressus Numerantium, **132** (1998), 9–18.
- [18] W.C. Shiu, P.C.B. Lam and S-M Lee, *On a Construction of Supermagic Graphs*, Journal of Combinatorial Mathematics and Combinatorial Computing, **42** (2002), 147–160.
- [19] R.P. Stanley, *Linear homogeneous diophantine equations and magic labelings of graphs*, Duke Math. J., **40** (1973), 607–632.
- [20] R.P. Stanley, *Magic labeling of graphs, symmetric magic squares, systems of parameters and Cohen-Macaulay rings*, Duke Math. J., **40** (1976), 511–531.
- [21] W.D. Wallis, *Magic Graphs*, Birkhauser Boston, (2001).