

### Mid-Semester Test

Name : \_\_\_\_\_

Adm. No. : \_\_\_\_\_

Class : \_\_\_\_\_

Class S/N : \_\_\_\_\_

Date : \_\_\_\_\_

Time allowed : 1 hour

Maximum mark : 100

#### Instructions

Answer all 4 questions. Take  $g = 9.80 \text{ m/s}^2$ .


This question paper consists of 3 printed pages including 1 page of formulae.

You are reminded that cheating during this test is a serious offence.


All working in support of your answer must be shown. Answers must be to appropriate significant figures.

- 
1. a) In the below equation, the SI units of  $x$  and  $x_0$  are in metres,  $t$  and  $t_0$  are in seconds,  $v_0$  is m/s and  $a$  is  $\text{m/s}^2$ . Using the dimensional analysis show whether the equation is homogenous or not.
- $$x = x_0 + v_0(t - t_0) + \frac{1}{2}a(t - t_0)^2$$
- b) A particle is under a constant force  $\mathbf{F} = (-2\mathbf{i} + 4\mathbf{j} + \mathbf{k})$  as it moves from a point (5, 1, 2) to another point (-2, 3, 4). Find the work done on the particle.
- c) What is the angle between the displacement and force vectors in 1(b)?
- (25 marks)

2. In the figure below, car A and car B, moves towards each other in adjacent lanes and parallel to an  $x$  axis. At time  $t = 0$ , car A is at  $x = -35.0 \text{ m}$  and accelerates uniformly from rest at  $2 \text{ m/s}^2$  while car B is at  $x = 270 \text{ m}$  moving at a constant speed of  $20.0 \text{ m/s}$ .
- a) When do the cars meet?
- b) Where do the cars meet?
- c) What is the speed of car A when it crosses car B?
- d) Sketch the displacement-time plots of the two cars on the same graph.



3. The position vector of a particle of mass 2.0 kg moving on the  $x$ - $y$  plane is  $\vec{r} = 2t \hat{i} + 2\sin(\pi t / 4) \hat{j}$ , with  $\vec{r}$  in metres and  $t$  in seconds. Calculate in component form the
- particle's average velocity during the first second of its journey.
  - particle's velocity at  $t = 1.0$  s.
  - force acting on the particle at  $t = 1.0$  s.
- (25 marks)
4. The two blocks ( $A = 16$  kg and  $B = 88$  kg) in the figure below are not attached to each other. The coefficient of static friction between the two blocks is  $\mu_s = 0.33$ , but the surface beneath the larger block is frictionless.
- Draw the free-body diagram for blocks  $A$  and  $B$  if a force  $F$  acts on  $A$  as shown in the figure.
  - What is the magnitude of the minimum force  $F$  required to keep block  $A$  from slipping down block  $B$ ?
  - If  $F$  acts on block  $B$  instead of block  $A$ , what is the magnitude of  $F$  so that block  $A$  does not fall off block  $B$ ?



(25 marks)

\*\*\*\*\* End of Paper \*\*\*\*\*

**Formula sheet**

Name: \_\_\_\_\_ Admin. No.: \_\_\_\_\_ Seat No.: \_\_\_\_\_

<p><b><u>Kinematics</u></b></p> $v_x = v_{0x} + a_x t$ $v_x^2 = v_{0x}^2 + 2a_x(x - x_0)$ $x = x_0 + v_{0x}t + \frac{1}{2}a_x t^2$ $\vec{v} = \frac{d\vec{r}}{dt}, \quad \vec{a} = \frac{d\vec{v}}{dt}$ $y = (\tan \theta)x - \left(\frac{g}{2v^2 \cos^2 \theta}\right)x^2$ $R = \frac{v^2 \sin 2\theta}{g}$ <p><b><u>Dynamics</u></b></p> $\vec{F} = m \frac{d\vec{v}}{dt} = m\vec{a}, F = \mu N$ $a = \frac{dv}{dt}, a = \frac{v^2}{r}, F = m \frac{v^2}{r}$ $\vec{J} = \int \vec{F} dt = \Delta \vec{p}$ $W = \int \vec{F} \cdot d\vec{r}, W_{net} = K_f - K_i$ $KE = \frac{1}{2}mv^2, PE = mgh$ $P = \frac{W}{t}, P = \frac{dW}{dt}$ <p><b><u>Linear momentum</u></b></p> $m_1 \vec{u}_1 + m_2 \vec{u}_2 = m_1 \vec{v}_1 + m_2 \vec{v}_2$ <p><b><u>Static electricity</u></b></p> $F = k \frac{q_1 q_2}{r^2}, k = \frac{1}{4\pi\epsilon_0}$ $F = qE$ $V = k \frac{q}{r}, U = qV$ $\Phi_E = \oint \vec{E} \cdot d\vec{A} = \frac{q}{\epsilon_0}$ $V = Ed, W = qV, E = \frac{kq}{r^2}$	<p><b><u>Current electricity</u></b></p> $Q = It \quad V = IR$ $P = VI = I^2 R = \frac{V^2}{R}$ <p><b><u>Magnetism &amp; electromagnetism</u></b></p> $\vec{F} = q\vec{v} \times \vec{B} \quad \vec{F} = i\vec{L} \times \vec{B}$ $e.m.f. = -N \frac{d\Phi_B}{dt} \quad \Phi_B = BA$ <p><b><u>Thermodynamics</u></b></p> $\Delta U = Q - W$ $W = \int p dV$ $Q_V = nC_V \Delta T \quad \text{const vol}$ $Q_p = nC_p \Delta T \quad \text{const pressure}$ $Q = mC \Delta T$ $Q = mL$ <p><b><u>Ideal Gas</u></b></p> $pV = nRT$ $pV^\gamma = c \text{ (adiabatic)}$ $\gamma = \frac{C_p}{C_v}, C_p - C_v = R$ $W = pV \ln \frac{V_2}{V_1} = nRT \ln \frac{V_2}{V_1}$ $W = \frac{1}{\gamma - 1} (p_1 V_1 - p_2 V_2)$ <p><b><u>Rotational Motion</u></b></p> $\omega = \frac{d\theta}{dt}, \quad \alpha = \frac{d\omega}{dt}$ $\omega = \omega_0 + \alpha t$ $\omega^2 = \omega_0^2 + 2\alpha(\theta - \theta_0)$ $\theta = \theta_0 + \omega_0 t + \frac{1}{2}\alpha t^2$ $I = \sum_i m_i r_i^2, I = \int r^2 dm, K = \frac{1}{2} I \omega^2$	<p><b><u>SHM &amp; waves</u></b></p> $T = \frac{1}{f} \quad v = f\lambda \quad \omega = 2\pi f$ $\omega = \frac{2\pi}{T} \quad k = \frac{2\pi}{\lambda}$ $\omega = \sqrt{k/m} \quad \omega = \sqrt{g/L}$ $x = A \cos(\omega t + \phi)$ $x = A \sin(\omega t + \phi)$ $y(x, t) = A \cos(\omega t + kx)$ $y(x, t) = A \sin(\omega t + kx)$ $\frac{\partial^2 y}{\partial x^2} = \frac{1}{v^2} \frac{\partial^2 y}{\partial t^2}$ <p><b><u>Circuits</u></b></p> $R = R_1 + R_2 + R_3 + \dots \quad \text{series}$ $\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots \quad \text{parallel}$ $\frac{1}{C} = \frac{1}{C_1} + \frac{1}{C_2} + \frac{1}{C_3} + \dots \quad \text{series}$ $C = C_1 + C_2 + C_3 + \dots \quad \text{parallel}$ $Q = CV \quad U = \frac{1}{2} CV^2$ <p><b><u>Constants</u></b></p> <p>Charge on electron  <math>e = -1.60 \times 10^{-19} \text{ C}</math></p> <p>Coulomb's constant  <math>k = 9 \times 10^9 \text{ N m}^2 \text{ C}^{-2}</math></p> <p>Ideal gas constant  <math>R = 8.314 \text{ J mol}^{-1} \text{ K}^{-1}</math></p> <p>Mass of proton  <math>m_p = 1.67 \times 10^{-27} \text{ kg}</math></p> <p>Mass of electron  <math>m_e = 9.11 \times 10^{-31} \text{ kg}</math></p> <p>Permeability of free space  <math>\mu_0 = 4\pi \times 10^{-7} \text{ N A}^{-2}</math></p> <p>Permittivity of free space  <math>\epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2}</math></p> <p>Speed of light in vacuum  <math>c = 3.00 \times 10^8 \text{ m s}^{-1}</math></p>
--	--	---