

jee2022-paper1

1. considering only the principal values of the inverse trigonometric functions, the value of $\frac{3}{2} \cos^{-1} \sqrt{\frac{2}{2+\pi^2}} + \frac{1}{4} \sin^{-1} \frac{2\sqrt{2}\pi}{2+\pi^2} + \tan^{-1} \frac{\sqrt{2}}{\pi}$ is _____
2. Let α be a positive real number. let $f: \mathbb{R} \rightarrow \mathbb{R}$ and $g: (\alpha, \infty) \rightarrow \mathbb{R}$ be the functions defined by $f(x) = \sin(\frac{\pi x}{12})$ and $g(x) = \frac{2 \log_e(\sqrt{x} - \sqrt{\alpha})}{\log_e(e^{\sqrt{x}} - e^{\sqrt{\alpha}})}$ then the value of $\lim_{x \rightarrow \alpha} f(g(x))$ is _____
3. In a study about a pandemic, data of 900 persons was collected. It was found that

- 190 persons had symptom of fever,
- 220 persons had symptom of cough,
- 220 persons had symptom of breathing problem,
- 330 persons had symptom of fever or cough or both,
- 350 persons had symptom of cough or breathing problem or both,
- 340 persons had symptom of fever or breathing problem or both,
- 30 persons had all three symptoms

If a person is chosen randomly from these 900 persons, then the probability that the person has at most one symptom is _____

4. Let z be a complex number with non-zero imaginary part. If $\frac{2+3z+4z^2}{2-3z+4z^2}$ is a real number, then the value of $|z|^2$ is _____
5. Let \bar{z} denote the complex conjugate of a complex number z and let $i = \sqrt{-1}$. In the set of complex numbers, the number of distinct roots of the equation $\bar{z} - z^2 = i(\bar{z} + z^2)$ is _____
6. Let l_1, l_2, \dots, l_{100} be consecutive terms of an arithmetic progression with common difference d_1 , and let w_1, w_2, \dots, w_{100} be consecutive terms of another arithmetic progression with common difference d_2 , where $d_1 d_2 = 10$. For each $i = 1, 2, \dots, 100$, let R_i be a rectangle with length L_i , width W_i and area A_i . If $A_{51} - A_{50} = 1000$, then the value of $A_{100} - A_{90}$ is _____

7. the number of 4-digit integers in the closed interval $[2022, 4482]$ formed by using the digits 0, 2, 3, 4, 6, 7 is
8. Let ABC be the triangle with $AB = 1$, $AC = 3$ and $\angle BAC = \frac{\pi}{2}$. If a circle of radius $r > 0$ touches the sides AB, AC and also touches internally the circumcircle of the triangle ABC, then the value of r is _____
9. consider the equation

$$\int_1^e \frac{(\log_e x)^{\frac{1}{2}}}{x(a - (\log_e x)^{\frac{3}{2}})^2} dx = 1, a \in (-\infty, 0) \cup (1, \infty)$$
which of the following statements is/are TRUE?
- A. No a satisfies the above equation
B. An integer a satisfies the above equation
C. An irrational number a satisfies the above equation
D. More than one a satisfy the above equation
10. Let a_1, a_2, a_3, \dots be an arithmetic progression with $a_1 = 7$ and common difference 8. Let T_1, T_2, T_3, \dots be such that $T_1 = 3$ and $T_{n+1} - T_n = a_n$ for $n \geq 1$. Then, which of the following is/are TRUE?
- A. $T_{20} = 1604$
B. $\sum_{k=1}^{20} T_k = 10510$
C. $T_{30} = 3454$
D. $\sum_{k=1}^{30} T_k = 357610$
11. Let P_1 and P_2 be two planes given by
- $P_1 : 10x + 15y + 12z - 60 = 0$,
 - $P_2 : -2x + 5y + 4z - 20 = 0$.

Which of the following straight lines can be an edge of some tetrahedron whose two faces lie on P_1 and P_2

- A. $\frac{x-1}{0} = \frac{y-1}{0} = \frac{z-1}{5}$
B. $\frac{x-6}{-5} = \frac{y}{2} = \frac{z}{3}$
C. $\frac{x}{-2} = \frac{y-4}{5} = \frac{z}{4}$
D. $\frac{x}{1} = \frac{y-4}{-2} = \frac{z}{3}$
12. Let S be the reflection of a point Q with respect to the plane given by
 $\uparrow r = -(t+p)\hat{i} + t\hat{j} + (1+p)\hat{k}$
where t, p are real parameters and $\hat{i}, \hat{j}, \hat{k}$ are the unit vectors along the three positive coordinate axes. If the position vectors of Q and S are $10\hat{i} + 15\hat{j} + 20\hat{k}$ and $\alpha\hat{i} + \beta\hat{j} + \gamma\hat{k}$ respectively, then which of the following is/are TRUE ?

- A. $3(\alpha + \beta) = -101$
 B. $3(\beta + \gamma) = -71$
 C. $3(\gamma + \alpha) = -86$
 D. $3(\alpha + \beta + \gamma) = -121$
13. Consider the parabola $y^2 = 4x$. Let S be the focus of the parabola. A pair of tangents drawn to the parabola from the point p = (-2, 1) meet the parabola at P_1 and P_2 . Let Q_1 and Q_2 be points on the lines SP_1 and SP_2 respectively such that PQ_1 is perpendicular to SP_1 and PQ_2 is perpendicular to SP_2 . Then, which of the following is/are TRUE ?
- A. $SQ_1 = 2$
 B. $Q_1Q_2 = \frac{3\sqrt{10}}{5}$
 C. $PQ_1 = 3$
 D. $SQ_2 = 1$
14. Let $|M|$ denote the determinant of a square matrix M. Let $g: [0, \frac{\pi}{2}] \rightarrow \mathbb{R}$ be the function defined by $g(\theta) = \sqrt{f(\theta) - 1} + \sqrt{f(\frac{\pi}{2} - \theta) - 1}$ where
- $$f(\theta) = \frac{1}{2} \begin{vmatrix} 1 & \sin \theta & 1 \\ -\sin \theta & 1 & \sin \theta \\ -1 & -\sin \theta & 1 \end{vmatrix} + \begin{vmatrix} \sin \pi & \cos(\theta + \frac{\pi}{4}) & \tan(\theta - \frac{\pi}{4}) \\ \sin(\theta - \frac{\pi}{4}) & -\cos \frac{\pi}{2} & \log_e(\frac{4}{\pi}) \\ \cot(\theta + \frac{\pi}{4}) & \log_e(\frac{\pi}{4}) & \tan \pi \end{vmatrix}$$
- Let p(x) be a quadratic polynomial whose roots are the maximum and minimum values of the function $g(\theta)$ and $p(2) = 2 - \sqrt{2}$. Then, which of the following is/are TRUE ?
- A. $p(\frac{3+\sqrt{2}}{4}) < 0$
 B. $p(\frac{1+3\sqrt{2}}{4}) > 0$
 C. $p(\frac{5\sqrt{2}-1}{4}) > 0$
 D. $p(\frac{5-\sqrt{2}}{4}) < 0$
15. consider the following lists:
- | List-I | List-II |
|---|------------------------|
| (I) $x \in [-\frac{2\pi}{3}, \frac{2\pi}{3}] : \cos x + \sin x = 1$ | (P) has two elements |
| (II) $x \in [-\frac{5\pi}{18}, \frac{5\pi}{18}] : \sqrt{3} \tan 3x = 1$ | (Q) has three elements |
| (III) $x \in [-\frac{6\pi}{5}, \frac{6\pi}{5}] : 2 \cos(2x) = \sqrt{3}$ | (R) has four elements |
| (IV) $x \in [-\frac{7\pi}{4}, \frac{7\pi}{4}] : \sin x - \cos x = 1$ | (S) has five elements |
| (V) | (T) has six elements |
- The correct option is :

- A. $(I) \rightarrow (P); (II) \rightarrow (S); (III) \rightarrow (P); (IV) \rightarrow (S)$

- B. $(I) \rightarrow (P); (II) \rightarrow (P); (III) \rightarrow (T); (IV) \rightarrow (R)$
 C. $(I) \rightarrow (Q); (II) \rightarrow (P); (III) \rightarrow (T); (IV) \rightarrow (S)$
 D. $(I) \rightarrow (Q); (II) \rightarrow (S); (III) \rightarrow (P); (IV) \rightarrow (R)$
16. Two players, P_1 and P_2 , play a game against each other. In every round of the game, each player rolls a fair die once, where the six faces of the die have six distinct numbers. Let x and y denote the readings on the die rolled by P_1 and P_2 , respectively. If $x > y$, then P_1 scores 5 points and P_2 scores 0 point. If $x = y$, then each player scores 2 points. If $x < y$, then P_1 scores 0 point and P_2 scores 5 points. Let X_i and Y_i be the total scores of P_1 and P_2 , respectively, after playing the i^{th} round.

LIST-I

- (I) Probability of $(X_2 \geq Y_2)$ is
 (II) Probability of $(X_2 > Y_2)$ is
 (III) Probability of $(X_3 = Y_3)$ is
 (IV) Probability of $(X_3 > Y_3)$ is
 (V)

LIST-II

- $(P) = \frac{3}{8}$
 $(Q) = \frac{11}{16}$
 $(R) = \frac{5}{16}$
 $(S) = \frac{355}{864}$
 $(T) = \frac{77}{432}$

the correct option is:

- A. $(I) \rightarrow (Q); (II) \rightarrow (R); (III) \rightarrow (T); (IV) \rightarrow (S)$
 B. $(I) \rightarrow (Q); (II) \rightarrow (R); (III) \rightarrow (T); (IV) \rightarrow (S)$
 C. $(I) \rightarrow (P); (II) \rightarrow (R); (III) \rightarrow (Q); (IV) \rightarrow (S)$
 D. $(I) \rightarrow (P); (II) \rightarrow (R); (III) \rightarrow (Q); (IV) \rightarrow (T)$
17. Let p, q and r be nonzero real numbers that are the 10^{th} , 100^{th} , and 1000^{th} terms of a harmonic progression, respectively. Consider the following system of linear equations:

$$x + y + z = 1$$

$$10x + 100y + 1000z = 0$$

$$qrx + pry + pqz = 0$$

LIST-I

- (I) If $\frac{q}{r} = 10$, then the system of linear equations has
 (II) If $\frac{p}{r} \neq 100$, then the system of linear equations has
 (III) If $\frac{p}{q} \neq 10$, then the system of linear equations has
 (IV) If $\frac{p}{q} = 10$, then the system of linear equations has

LIST-II

- (P) $x = 0, y = \frac{10}{9}, z = -\frac{1}{9}$ as a solution
 (Q) $x = \frac{10}{9}, y = -\frac{1}{9}, z = 0$ as a solution
 (R) infinitely many solutions
 (S) no solution

- A. $(I) \rightarrow (T); (II) \rightarrow (R); (III) \rightarrow (S); (IV) \rightarrow (T)$
 B. $(I) \rightarrow (Q); (II) \rightarrow (S); (III) \rightarrow (S); (IV) \rightarrow (R)$

- C. $(I) \rightarrow (Q); (II) \rightarrow (R); (III) \rightarrow (P); (IV) \rightarrow (R)$
D. $(I) \rightarrow (T); (II) \rightarrow (S); (III) \rightarrow (P); (IV) \rightarrow (T)$

18. consider the ellipse $\frac{x^2}{4} + \frac{y^2}{3} = 1$ let $H(\alpha, 0), 0 < \alpha < 2$, be a point. A straight line drawn through H parallel to the y-axis crosses the ellipse and its auxiliary circle at points E and F respectively, in the first quadrant. The tangent to the ellipse at the point E intersects the positive x-axis at a point G. Suppose the straight line joining F and the origin makes an angle ϕ with the positive x-axis

LIST-I

- (I) if $\varphi = \frac{3}{4}$, then the area of the triangle FGH is
(II) if $\varphi = \frac{\pi}{3}$, then the area of the triangle FGH is
(III) if $\varphi = \frac{\pi}{6}$, then the area of the triangle FGH is
(IV) if $\varphi = \frac{\pi}{12}$, then the area of the triangle FGH is
(V)

LIST-II

- (P) $\frac{(\sqrt{3}-1)^4}{8}$
(Q) 1
(R) $\frac{3}{4}$
(S) $\frac{1}{2\sqrt{3}}$
(T) $\frac{3\sqrt{3}}{2}$

- A. $(I) \rightarrow (R); (II) \rightarrow (S); (III) \rightarrow (Q); (IV) \rightarrow (P)$
B. $(I) \rightarrow (R); (II) \rightarrow (T); (III) \rightarrow (S); (IV) \rightarrow (P)$
C. $(I) \rightarrow (Q); (II) \rightarrow (T); (III) \rightarrow (S); (IV) \rightarrow (P)$
D. $(I) \rightarrow (Q); (II) \rightarrow (S); (III) \rightarrow (Q); (IV) \rightarrow (P)$