


1.2



generalized coordinates

$$q = \begin{bmatrix} \theta_1 \\ \theta_2 \end{bmatrix}$$

$$L = KE_1 + KE_2 - PE_1 - PE_2$$

$$KE_1 = \frac{1}{2} m_1 v_1^2 + \frac{1}{2} I_1 \omega_1^2$$

$$\omega_1 = \dot{\theta}_1, \quad v_1^2 = \dot{x}_{r1}^2 + \dot{y}_{r1}^2$$

$$x_{r1} = r_1 \sin \theta_1 \Rightarrow \dot{x}_{r1} = r_1 \dot{\theta}_1 \cos \theta_1$$

$$y_{r1} = r_1 \cos \theta_1 \Rightarrow \dot{y}_{r1} = -r_1 \dot{\theta}_1 \sin \theta_1$$

$$v_1^2 = r_1^2 \dot{\theta}_1^2$$

$$KE_1 = \frac{1}{2} m_1 r_1^2 \dot{\theta}_1^2 + \frac{1}{2} I_1 \dot{\theta}_1^2$$


$$PE_1 = m g r_1 \cos \theta_1$$

$$KE_2 = \frac{1}{2} m_2 v_2^2 + \frac{1}{2} I_2 \omega_2^2$$

$$\omega_2 = \dot{\theta}_1 + \dot{\theta}_2, \quad v_2^2 = \dot{x}_{r2}^2 + \dot{y}_{r2}^2$$

$$x_{r2} = l_1 \sin \theta_1 + r_2 \sin(\theta_1 + \theta_2)$$

$$\dot{x}_{r2} = l_1 \dot{\theta}_1 \cos \theta_1 + r_2 (\dot{\theta}_1 + \dot{\theta}_2) \cos(\theta_1 + \theta_2)$$



$$y_m = l_1 \cos \theta_1 + r_2 \cos (\theta_1 + \theta_2)$$

$$\dot{y}_m = -l_1 \dot{\theta}_1 \sin \theta_1 - r_2 (\dot{\theta}_1 + \dot{\theta}_2) \sin (\theta_1 + \theta_2)$$

$$\begin{aligned} v_2^2 &= l_1^2 \dot{\theta}_1^2 \cos^2 \theta_1 + r_2^2 (\dot{\theta}_1 + \dot{\theta}_2)^2 \cos^2 (\theta_1 + \theta_2) + \\ &\quad 2 l_1 \dot{\theta}_1 r_2 (\dot{\theta}_1 + \dot{\theta}_2) \cos \theta_1 \cos (\theta_1 + \theta_2) + l_1^2 \dot{\theta}_1^2 \sin^2 \theta_1 + \\ &\quad r_2^2 (\dot{\theta}_1 + \dot{\theta}_2)^2 \sin^2 (\theta_1 + \theta_2) + 2 l_1 \dot{\theta}_1 r_2 (\dot{\theta}_1 + \dot{\theta}_2) \sin \theta_1 \sin (\theta_1 + \theta_2) \end{aligned}$$

$$= l_1^2 \dot{\theta}_1^2 + r_2^2 (\dot{\theta}_1 + \dot{\theta}_2)^2 + 2 l_1 \dot{\theta}_1 r_2 (\dot{\theta}_1 + \dot{\theta}_2) \cos \theta_2$$

$$\begin{aligned} KE_2 &= \frac{1}{2} m_2 l_1^2 \dot{\theta}_1^2 + \frac{1}{2} m_2 r_2^2 (\dot{\theta}_1 + \dot{\theta}_2)^2 + m_2 l_1 \dot{\theta}_1 r_2 (\dot{\theta}_1 + \dot{\theta}_2) \cos \theta_2 \\ &\quad + \frac{1}{2} I_2 (\dot{\theta}_1 + \dot{\theta}_2)^2 \end{aligned}$$

$$PE_2 = m_2 g (l_1 \cos \theta_1 + r_2 \cos (\theta_1 + \theta_2))$$

$$PE_2 = m_2 g l_1 \cos \theta_1 + m_2 g r_2 \cos (\theta_1 + \theta_2)$$

$$L = (KE_1 + KE_2) - (PE_1 + PE_2)$$

$$KE_1 = \frac{1}{2} m_1 r_1^2 \dot{\theta}_1^2 + \frac{1}{2} I_1 \dot{\theta}_1^2$$

$$PE_1 = m_1 g r_1 \cos \theta_1$$

$$\begin{aligned} L &= \frac{1}{2} m_1 r_1^2 \dot{\theta}_1^2 + \frac{1}{2} I_1 \dot{\theta}_1^2 + \frac{1}{2} m_2 l_1^2 \dot{\theta}_1^2 + \frac{1}{2} m_2 r_2^2 (\dot{\theta}_1 + \dot{\theta}_2)^2 \\ &\quad + m_2 l_1 \dot{\theta}_1 r_2 (\dot{\theta}_1 + \dot{\theta}_2) \cos \theta_2 + \frac{1}{2} I_2 (\dot{\theta}_1 + \dot{\theta}_2)^2 - m_1 g r_1 \cos \theta_1 \\ &\quad - m_2 g l_1 \cos \theta_1 - m_2 g r_2 \cos (\theta_1 + \theta_2) \end{aligned}$$

$$\begin{aligned} \frac{\partial L}{\partial \dot{\theta}_1} &= m_1 r_1^2 \dot{\theta}_1 + I_1 \dot{\theta}_1 + m_2 l_1^2 \dot{\theta}_1 + m_2 r_2^2 (\dot{\theta}_1 + \dot{\theta}_2) + m_2 l_1 \dot{\theta}_1 r_2 \cos \theta_2 \\ &\quad + m_2 l_1 r_2 (\dot{\theta}_1 + \dot{\theta}_2) \cos \theta_2 + I_2 (\dot{\theta}_1 + \dot{\theta}_2) \end{aligned}$$

$$\frac{\partial L}{\partial \theta_1} = m_1 g r_1 \sin \theta_1 + m_2 g l_1 \sin \theta_1 + m_2 g r_2 \sin(\theta_1 + \theta_2)$$

$$\frac{\partial L}{\partial \dot{\theta}_1} = m_1 r_1^2 \dot{\theta}_1 + I_1 \dot{\theta}_1 + m_2 l_1^2 \dot{\theta}_1 + m_2 r_2^2 \dot{\theta}_1 + m_2 l_1 \dot{\theta}_1 r_2 \cos \theta_2 + m_2 l_1 r_2 \dot{\theta}_1 \cos \theta_2 + m_2 l_1 r_2 \dot{\theta}_2 \cos \theta_2 + I_2 (\dot{\theta}_1 + \dot{\theta}_2)$$

$$\begin{aligned} \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{\theta}_1} \right) &= m_1 r_1^2 \ddot{\theta}_1 + I_1 \ddot{\theta}_1 + m_2 l_1^2 \ddot{\theta}_1 + m_2 r_2^2 \ddot{\theta}_1 + m_2 r_2^2 \ddot{\theta}_2 + \\ & m_2 l_1 \ddot{\theta}_1 r_2 \cos \theta_2 - m_2 l_1 \dot{\theta}_1 \dot{\theta}_2 r_2 \sin \theta_2 + m_2 l_1 r_2 \ddot{\theta}_1 \cos \theta_2 \\ & - m_2 l_1 r_2 \dot{\theta}_1 \dot{\theta}_2 \sin \theta_2 + m_2 l_1 r_2 \ddot{\theta}_2 \cos \theta_2 \\ & - m_2 l_1 r_2 \dot{\theta}_2^2 \sin \theta_2 + I_2 (\ddot{\theta}_1 + \ddot{\theta}_2) \end{aligned}$$

$$U_1 = \ddot{\theta}_2 [m_2 r_2^2 + m_2 r_2 l_1 \cos \theta_2 + I_2] - \dot{\theta}_2 [l_1 m_2 r_2 \dot{\theta}_1 \sin \theta_2 + l_1 m_2 r_2 \sin \theta_2 (\dot{\theta}_1 + \dot{\theta}_2)]$$

$$\begin{aligned} & \ddot{\theta}_1 [m_2 l_1^2 + 2 m_2 \cos \theta_2 l_1 r_2 + m_1 r_1^2 + m_2 r_2^2 + I_1 + I_2] \\ & - g l_1 m_2 \sin \theta_1 - g m_1 r_1 \sin \theta_1 - m_2 g r_2 \sin(\theta_1 + \theta_2) \end{aligned}$$

$$\begin{aligned} L &= \frac{1}{2} m_1 r_1^2 \dot{\theta}_1^2 + \frac{1}{2} I_1 \dot{\theta}_1^2 + \frac{1}{2} m_2 l_1^2 \dot{\theta}_1^2 + \frac{1}{2} m_2 r_2^2 (\dot{\theta}_1 + \dot{\theta}_2)^2 \\ & + m_2 l_1 \dot{\theta}_1 r_2 (\dot{\theta}_1 + \dot{\theta}_2) \cos \theta_2 + \frac{1}{2} I_2 (\dot{\theta}_1 + \dot{\theta}_2)^2 - m_1 g r_1 \cos \theta_1 \\ & - m_2 g l_1 \cos \theta_1 - m_2 g r_2 \cos(\theta_1 + \theta_2) \end{aligned}$$

$$\frac{\partial L}{\partial \dot{\theta}_2} = m_2 r_2^2 (\dot{\theta}_1 + \dot{\theta}_2) + m_2 l_1 \dot{\theta}_1 r_2 \cos \theta_2 + I_2 (\dot{\theta}_1 + \dot{\theta}_2)$$

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{\theta}_2} \right) = m_2 r_2^2 (\ddot{\theta}_1 + \ddot{\theta}_2) + m_2 l_1 \ddot{\theta}_1 r_2 \cos \theta_2 - m_2 l_1 \dot{\theta}_1 \dot{\theta}_2 r_2 \sin \theta_2 + I_2 (\ddot{\theta}_1 + \ddot{\theta}_2)$$

$$\frac{\partial L}{\partial \theta_2} = -m_2 l_1 \dot{\theta}_1 r_2 (\dot{\theta}_1 + \dot{\theta}_2) \sin \theta_2 + m_2 g r_2 \sin(\theta_1 + \theta_2)$$

$$u_2 = m_2 r_2^2 (\ddot{\theta}_1 + \ddot{\theta}_2) + m_2 l_1 \ddot{\theta}_1 r_2 \cos \theta_2 - m_2 l_1 \dot{\theta}_1 \dot{\theta}_2 r_2 \sin \theta_2 + I_2 (\ddot{\theta}_1 + \ddot{\theta}_2) + m_2 l_1 \dot{\theta}_1 r_2 (\dot{\theta}_1 + \dot{\theta}_2) \sin \theta_2 - m_2 g r_2 \sin(\theta_1 + \theta_2)$$

$$u_2 = \ddot{\theta}_2 [m_2 r_2^2 + I_2] + \ddot{\theta}_1 [m_2 r_2^2 + m_2 l_1 r_2 \cos \theta_2 + I_2] - g m_2 r_2 \sin(\theta_1 + \theta_2) + l_1 m_2 r_2 \dot{\theta}_1 (\dot{\theta}_1 + \dot{\theta}_2) \sin \theta_2 - l_1 m_2 r_2 \dot{\theta}_1 \dot{\theta}_2 \sin \theta_2$$

Manipulator equation:

$$M(q)\ddot{q} + C(q, \dot{q})\dot{q} + g(q) = \tau$$

$$M = \begin{bmatrix} m_2 l_1^2 + 2m_2 \cos \theta_2 l_1 r_2 + m_1 r_1^2 + m_2 r_2^2 + I_1 + I_2 & m_2 r_2^2 + m_2 r_2 l_1 \cos \theta_2 + I_2 \\ m_2 r_2^2 + m_2 l_1 r_2 \cos \theta_2 + I_2 & m_2 r_2^2 + I_2 \end{bmatrix}$$

$$g = \begin{bmatrix} -g l_1 m_2 \sin \theta_1 - g m_1 r_1 \sin \theta_1 - m_2 g r_2 \sin(\theta_1 + \theta_2) \\ -g m_2 r_2 \sin(\theta_1 + \theta_2) \end{bmatrix}$$

$$C = \begin{bmatrix} 0 & l_1 m_2 r_2 \dot{\theta}_1 \sin \theta_2 + l_1 m_2 r_2 \sin \theta_2 (\dot{\theta}_1 + \dot{\theta}_2) \\ l_1 m_2 r_2 (\dot{\theta}_1 + \dot{\theta}_2) \sin \theta_2 - l_1 m_2 r_2 \dot{\theta}_2 \sin \theta_2 & 0 \end{bmatrix}$$

M should be symmetric and positive definite matrix.

$$M = M^T \quad (\therefore \text{symmetric})$$

A symmetric matrix with all positive eigen values is positive definite matrix

This is verified with attached matlab code

b)

$$X = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} \theta_1 \\ \theta_2 \\ \dot{\theta}_1 \\ \dot{\theta}_2 \end{bmatrix}$$

$$\dot{X} = \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \\ \dot{x}_3 \\ \dot{x}_4 \end{bmatrix} = \begin{bmatrix} \ddot{\theta}_1 \\ \ddot{\theta}_2 \\ \dot{\theta}_1 \\ \dot{\theta}_2 \end{bmatrix} = \begin{bmatrix} x_3 \\ x_4 \\ \dot{\theta}_1 \\ \dot{\theta}_2 \end{bmatrix} = F(\theta_1, \theta_2, \dot{\theta}_1, \dot{\theta}_2, u_1, u_2)$$

$-\theta_1$

$$\begin{aligned} & (l_2^2 u_1 - l_2^2 u_2 + m_2 r_2^2 \ddot{u}_1 - m_2 r_2^2 \ddot{u}_2 + l_1^2 m_2^2 r_2^3 \dot{\theta}_1^2 \sin(\theta_2) + \\ & l_1^2 m_2^2 r_2^3 \dot{\theta}_2^2 \sin(\theta_2) + g l_1^2 m_2^2 r_2^2 \sin(\theta_1) + l_2^2 g l_1^2 m_2 \sin(\theta_1) + \\ & l_2^2 g m_1 r_1 \sin(\theta_1) - l_1^2 m_2^2 r_2^2 u_2 \cos(\theta_2) + \\ & 2 l_1^2 m_2^2 r_2^3 \dot{\theta}_1 \dot{\theta}_2 \sin(\theta_2) + \\ & l_1^2 m_2^2 r_2^2 \dot{\theta}_1^2 \cos(\theta_2) \sin(\theta_2) - g l_1^2 m_2^2 r_2^2 \sin(\theta_1 + \\ & \theta_2) \cos(\theta_2) + l_2^2 l_1^2 m_2^2 r_2 \dot{\theta}_1^2 \sin(\theta_2) + l_2^2 l_1^2 m_2^2 r_2 \dot{\theta}_2^2 \sin(\theta_2) + \\ & g m_1^2 m_2 r_1 r_2^2 \sin(\theta_1) + 2 l_2^2 l_1^2 m_2^2 r_2 \dot{\theta}_1 \dot{\theta}_2 \sin(\theta_2)) / (- \\ & l_1^2 m_2^2 r_2^2 \cos(\theta_2)^2 + l_1^2 m_2^2 r_2^2 + l_2^2 l_1^2 m_2 + m_1^2 m_2 r_1^2 r_2^2 + l_1^2 m_2 r_2^2 + \\ & l_2^2 m_1 r_1^2 + l_1^2 l_2) \end{aligned}$$

$$\ddot{\theta}_2 =$$

$$\begin{aligned} & -(l_2^2 u_1 - l_1^2 u_2 - l_2^2 u_2 - l_1^2 m_2^2 u_2 - m_1^2 r_1^2 u_2 + m_2 r_2^2 \ddot{u}_1 - m_2 r_2^2 \ddot{u}_2 + \\ & l_1^2 m_2^2 r_2^3 \dot{\theta}_1^2 \sin(\theta_2) + l_1^2 m_2^2 r_2^3 \dot{\theta}_2^2 \sin(\theta_2) - g l_1^2 m_2^2 r_2^2 \sin(\theta_1 + \theta_2) - \\ & l_1^2 g m_2^2 r_2^2 \sin(\theta_1 + \theta_2) + g l_1^2 m_2^2 r_2^2 \sin(\theta_1) + l_2^2 g l_1^2 m_2 \sin(\theta_1) + \\ & l_2^2 g m_1 r_1 \sin(\theta_1) + l_1^2 m_2^2 r_2^2 u_1 \cos(\theta_2) - 2 l_1^2 m_2^2 r_2^2 u_2 \cos(\theta_2) + \\ & 2 l_1^2 m_2^2 r_2^3 \dot{\theta}_1 \dot{\theta}_2 \sin(\theta_2) + \\ & 2 l_1^2 m_2^2 r_2^2 \dot{\theta}_1^2 \cos(\theta_2) \sin(\theta_2) + \\ & l_1^2 m_2^2 r_2^2 \dot{\theta}_2^2 \cos(\theta_2) \sin(\theta_2) - g l_1^2 m_2^2 r_2^2 \sin(\theta_1 + \\ & \theta_2) \cos(\theta_2) + g l_1^2 m_2^2 r_2^2 \cos(\theta_2) \sin(\theta_1) - g m_1^2 m_2 r_1^2 r_2^2 \sin(\theta_1 + \theta_2) \\ & + l_1^2 l_1^2 m_2^2 r_2 \dot{\theta}_1^2 \sin(\theta_2) + l_2^2 l_1^2 m_2^2 r_2 \dot{\theta}_1^2 \sin(\theta_2) + \\ & l_2^2 l_1^2 m_2^2 r_2 \dot{\theta}_2^2 \sin(\theta_2) + g m_1^2 m_2 r_1 r_2^2 \sin(\theta_1) + \end{aligned}$$

c) simulation plot time vs $\theta_1, \theta_2, \dot{\theta}_1, \dot{\theta}_2$

