


INDIAN INSTITUTE OF ENGINEERING SCIENCE AND TECHNOLOGY, SHIBPUR
B.TECH 1st SEM MID-SEMESTER (Group V–VIII) EXAMINATION, FEBRUARY 2022
Chemistry (CH1101)

Time: 45 Min.


Full Marks: 30

Answer all questions


1. (a) Which one is more acidic between **A** and **B** and explain.



- (b) Write the product of the following reaction and give the reaction mechanism.



- (c) State the Beer-Lambert Law of absorption of light and write its limitations.
 (d) Calculate the λ_{\max} (in nm) of the following compounds.




[2+3+2+(2×1½)]

2. (a) What are the essential criteria that a chelating agent must satisfy for being considered for chelation therapy? Suggest chelating agents for treatment of any two of the following:
 i) Pb(II), ii) Pu(IV), iii) As (III) toxicities.

- (b) Write down criteria of radioelement for using as radiodiagnostic agent and radiotherapeutic agent. Give examples of radionuclides used as radiodiagnostic agent and radiotherapeutic agent.


[5+5]

3. (a) Consider the following sequential reaction scheme:



Assuming that only reactant A is present at t=0, what is the expected time dependence of [P] using the steady-state approximation?

- (b) Consider the inter-conversion of the two conformations of a species:



The reaction is first order in each direction, with an equilibrium constant of 10^4 . The activation energy for the conversion of the B→A is 42 kJ mol⁻¹. Assuming an Arrhenius pre-exponential factor of 10¹² s⁻¹, what is the expected observed reaction rate constant at 298K if one were to initiate this reaction starting with only the reactant A at initial condition?

[6+4]