

Answer Key and Marking Scheme

Q1

- (A) i) The line is $x - 2y + 3 = 0$. The optimization problem to be minimized to obtain the closest point is given by

$$\begin{aligned} &\min (4 - x)^2 + (2 - y)^2 \\ &\text{subject to } x - 2y + 3 = 0 \end{aligned}$$

(0.5 marks)

Construct the lagrangian equation as

$$L(x, y, \lambda) = (4 - x)^2 + (2 - y)^2 + \lambda(x - 2y + 3) = 0$$

Solve the equation $\nabla L(x, y, \lambda) = \mathbf{0}$

$$2(x - 4) + \lambda = 0, 2(y - 2) - 2\lambda = 0, x - 2y + 3 = 0 \quad (0.5 \text{ marks})$$

$$\text{Then , } x = \frac{17}{5}, y = \frac{16}{5} \quad (0.5 \text{ marks})$$

$$\text{And distance} = \sqrt{(4 - \frac{17}{5})^2 + (2 - \frac{16}{5})^2} = \sqrt{\frac{9}{5}} \quad (0.5 \text{ marks})$$

- ii) The line is $x + 2y + 5 = 0$. The optimization problem to be minimized to obtain the closest point is given by

$$\begin{aligned} &\min (4 - x)^2 + (2 - y)^2 \\ &\text{subject to } x + 2y + 5 = 0 \end{aligned}$$

(0.5 marks)

Construct the lagrangian equation as

$$L(x, y, \lambda) = (4 - x)^2 + (2 - y)^2 + \lambda(x + 2y + 5) = 0$$

Solve the equation $\nabla L(x, y, \lambda) = \mathbf{0}$

$$2(x - 4) + \lambda = 0, 2(y - 2) + 2\lambda = 0, x + 2y + 5 = 0 \quad (0.5 \text{ marks})$$

$$\text{Then , } x = \frac{7}{5}, y = \frac{-16}{5} \quad (0.5 \text{ marks})$$

$$\text{And distance} = \sqrt{(4 - \frac{7}{5})^2 + (2 - \frac{-16}{5})^2} = \sqrt{\frac{169}{5}} \quad (0.5 \text{ marks})$$

- (B) For the given matrix C , we will first find the eigenvalues, so that we can use the trace and determinant property :
Construct equation $\det(C - \lambda I) = 0$. The resultant characteristic equation is given by

$$\det(C - \lambda I) = \lambda^3 - 14\lambda^2 - 9\lambda + 126$$

(0.5 mark)

Factorize to get $\det(C - \lambda I) = (\lambda - 3)(\lambda + 3)(\lambda - 14) = 0$. The eigenvalues of C is given by: $\lambda_1 = 14, \lambda_2 = 3, \lambda_3 = -3$ (0.5 mark)

- i) Recall that if λ is an eigenvalue of C , then λ^n is an eigenvalue of C^n .

Hence, $\lambda_1^6, \lambda_2^6, \lambda_3^6$ are the 3 eigenvalues of C^6 . (0.5 mark)
Hence

$$\begin{aligned} \text{Trace}(C^6) &= \lambda_1^6 + \lambda_2^6 + \lambda_3^6 \\ &= (14)^6 + (3)^6 + (-3)^6 \end{aligned}$$

(0.5 mark)

ii) Recall that $\det(AB) = \det(A)\det(B)$.
Hence $\det(C^7) = \det(C)^7$. (0.5 mark)

We know that $\det(C) = \lambda_1 * \lambda_2 * \lambda_3 = 14 * 3 * (-3) = -126$.

$$\det(C^7) = \det(C)^7 = (-126)^7$$

(0.5 mark)

(C) Given that $\mathbf{a}, \mathbf{b}, \mathbf{c}$ are linearly independent. Hence $\beta_1\mathbf{a} + \beta_2\mathbf{b} + \beta_3\mathbf{c} = \mathbf{0}$ will have only trivial solution $\beta_1 = \beta_2 = \beta_3 = 0$. (0.5 marks)

Now consider $\alpha_1\mathbf{x} + \alpha_2\mathbf{y} + \alpha_3\mathbf{z} = \mathbf{0}$.

$$\alpha_1\mathbf{x} + \alpha_2\mathbf{y} + \alpha_3\mathbf{z} = \alpha_1(\mathbf{b} - \mathbf{c}) + \alpha_2(\mathbf{a} + \mathbf{c}) + \alpha_3(\mathbf{a} - \mathbf{b}) = \mathbf{0}$$

$$\mathbf{a}(\alpha_2 + \alpha_3) + \mathbf{b}(\alpha_1 - \alpha_3) + \mathbf{c}(\alpha_2 - \alpha_1) = \mathbf{0}$$

(1 marks)

Since $\mathbf{a}, \mathbf{b}, \mathbf{c}$ are linearly independent, we can conclude that

$$\alpha_2 + \alpha_3 = 0,$$

$$\alpha_1 - \alpha_3 = 0, ,$$

$$\alpha_2 - \alpha_1 = 0$$

(0.5 marks)

Solving for $\alpha_1, \alpha_2, \alpha_3$ from the 3 equations, we get

$$\alpha_1 = 0, \alpha_2 = 0, \alpha_3 = 0$$

(0.5 marks)

Hence $\{\mathbf{x}, \mathbf{y}, \mathbf{z}\}$ is a linearly independent set.

(0.5 marks)

Q2

(A) i) We have

$$\begin{aligned} L(\boldsymbol{\beta}) &= \frac{1}{2p} \|\mathbf{y} - \boldsymbol{\beta}\|^2 + \lambda \|\mathbf{W}\boldsymbol{\beta}\|^2 \\ &= \frac{1}{p} \left(\sum_{j=1}^p \left[\frac{1}{2} (y_j - \beta_j)^2 + \lambda \|\mathbf{W}\boldsymbol{\beta}\|^2 \right] \right) \\ &= \frac{1}{p} \sum_{j=1}^p L_j(\boldsymbol{\beta}) \text{ where} \\ L_j(\boldsymbol{\beta}) &= \frac{1}{2} (y_j - \beta_j)^2 + \lambda \|\mathbf{W}\boldsymbol{\beta}\|^2 \end{aligned}$$

(1.5 marks)

ii) Now $L_j(\boldsymbol{\beta}) = \frac{1}{2} (y_j - \beta_j)^2 + \lambda \|\mathbf{W}\boldsymbol{\beta}\|^2 = (y_j - \beta_j)^2 + \lambda \boldsymbol{\beta}^T \mathbf{W}^T \mathbf{W} \boldsymbol{\beta}$.

We have

$$\frac{\partial [\frac{1}{2} (y_j - \beta_j)^2]}{\partial \beta_i} = v_i$$

where

$$v_i = \begin{cases} 0 & \text{when } i \neq j \\ -(y_j - \beta_j) & \text{when } i = j \end{cases}$$

(1 mark)

Also

$$\frac{\partial \lambda \boldsymbol{\beta}^T \mathbf{W}^T \mathbf{W} \boldsymbol{\beta}}{\partial \boldsymbol{\beta}} = 2\lambda \boldsymbol{\beta}^T \mathbf{W}^T \mathbf{W} = (2\lambda \mathbf{W}^T \mathbf{W} \boldsymbol{\beta})^T.$$

(0.5 marks)

Thus, we have

$$\nabla L_j(\boldsymbol{\beta}) = (\mathbf{v} + 2\lambda \mathbf{W}^T \mathbf{W} \boldsymbol{\beta})^T, j = 1, \dots, p.$$

where $\mathbf{v} = [v_1, \dots, v_p]^T$ is a p dimensional vector such that

$$v_i = \begin{cases} 0 & \text{when } i \neq j \\ -(y_j - \beta_j) & \text{when } i = j \end{cases}$$

(0.5 marks)

iii) Now $L(\boldsymbol{\beta}) = \frac{1}{p} \sum_{j=1}^p L_j(\boldsymbol{\beta})$. Therefore,

$$\begin{aligned} \nabla L(\boldsymbol{\beta}) &= \frac{1}{p} \sum_{j=1}^p \nabla L_j(\boldsymbol{\beta}) \\ &= -\frac{1}{p} (\mathbf{y} - \boldsymbol{\beta})^T + 2\lambda (\mathbf{W}^T \mathbf{W} \boldsymbol{\beta})^T. \end{aligned}$$

(1.5 marks)

(B) The dimension of the transformed data $= 1 + m + m + m - 2 = 3m - 1$.
(0.5 marks)

$$\begin{aligned}
K(\mathbf{x}, \mathbf{y}) &= \boldsymbol{\phi}(\mathbf{x})^T \boldsymbol{\phi}(\mathbf{y}) \\
&= 1 + x_1 y_1 + \cdots + x_m y_m + x_1^3 y_1^3 + \cdots + x_m^3 y_m^3 + x_1 x_2 x_3 y_1 y_2 y_3 \\
&\quad + x_2 x_3 x_4 y_2 y_3 y_4 + \cdots + x_{m-2} x_{m-1} x_m y_{m-2} y_{m-1} y_m.
\end{aligned}$$

(1 mark)

(C) We have $f(x, y) = (x^2 + y^2)e^{-(x^2+y^2)}$ and A is the set of all possible critical points of f . So,

i) To find critical points,

$$\begin{aligned}
f_x(x, y) &= 2xe^{-(x^2+y^2)}[1 - x^2 - y^2] = 0 \\
\Rightarrow 2x[1 - x^2 - y^2] &= 0 \\
\Rightarrow x = 0 \text{ or } x^2 + y^2 &= 1. \quad (0.5 \text{ marks}) \\
f_y(x, y) &= 2ye^{-(x^2+y^2)}[1 - x^2 - y^2] = 0 \\
\Rightarrow 2y[1 - x^2 - y^2] &= 0 \\
\Rightarrow y = 0 \text{ or } x^2 + y^2 &= 1. \quad (0.5 \text{ marks})
\end{aligned}$$

Therefore, $A = \{(0, 0)\} \cup \{(x, y) \in \mathbb{R}^2 | x^2 + y^2 = 1\}$. (0.5 marks)

ii) To find the nature of $(0, 0)$, consider

$$\begin{aligned}
f_{xx}(x, y) &= 2e^{-(x^2+y^2)} - 4x^2e^{-(x^2+y^2)} - (6x^2 + 2y^2)e^{-(x^2+y^2)} + 4x^2(x^2 + y^2)e^{-(x^2+y^2)} \\
\Rightarrow f_{xx}(0, 0) &= 2. \quad (0.5 \text{ marks}) \\
f_{yy}(x, y) &= 2e^{-(x^2+y^2)} - 4y^2e^{-(x^2+y^2)} - (2x^2 + 6y^2)e^{-(x^2+y^2)} + 4y^2(x^2 + y^2)e^{-(x^2+y^2)} \\
\Rightarrow f_{yy}(0, 0) &= 2. \quad (0.5 \text{ marks}) \\
f_{xy}(x, y) &= f_{yx}(x, y) \\
&= -8xye^{-(x^2+y^2)} + 4xy(x^2 + y^2)e^{-(x^2+y^2)} \\
\Rightarrow f_{xy}(0, 0) &= f_{yx}(0, 0) = 0. \quad (0.5 \text{ marks})
\end{aligned}$$

Clearly $H(0, 0) = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$ and hence positive definite as its eigenvalues are 2, 2. Therefore, the $(0, 0)$ is a point of minima. (0.5marks)

Q3

- (A) i) The claim is not true. (1.5 marks for finding a matrix satisfying all conditions.

1.5 marks for finding RREF of that matrix with \mathbf{C}_1 as one of the pivotal columns.)

- ii) Now

$$\begin{aligned}\mathbf{C}_1 &= \sum_{i=2}^6 \mathbf{C}_i \\ &= \mathbf{C}_2 + \sum_{i=3}^6 \mathbf{C}_i \\ &= 2\mathbf{C}_2.\end{aligned}$$

(0.5 marks)

$$\begin{aligned}\Rightarrow \mathbf{C}_1 - 2\mathbf{C}_2 + 0\mathbf{C}_3 + 0\mathbf{C}_4 + 0\mathbf{C}_5 + 0\mathbf{C}_6 &= \mathbf{0} \\ \Rightarrow [\mathbf{C}_1, \mathbf{C}_2, \mathbf{C}_3, \mathbf{C}_4, \mathbf{C}_5, \mathbf{C}_6][1, -2, 0, 0, 0, 0]^T &= \mathbf{0} \\ \Rightarrow \mathbf{A}[1, -2, 0, 0, 0, 0]^T &= \mathbf{0}.\end{aligned}$$

(0.5 marks)

Similarly

$$\begin{aligned}\mathbf{C}_1 &= \sum_{i=2}^6 \mathbf{C}_i \\ &= \mathbf{C}_2 + \sum_{i=3}^6 \mathbf{C}_i \\ &= 2\sum_{i=3}^6 \mathbf{C}_i.\end{aligned}$$

(0.5 marks)

$$\begin{aligned}\Rightarrow -1\mathbf{C}_1 + 0\mathbf{C}_2 + 2\mathbf{C}_3 + 2\mathbf{C}_4 + 2\mathbf{C}_5 + 2\mathbf{C}_6 &= \mathbf{0} \\ \Rightarrow [\mathbf{C}_1, \mathbf{C}_2, \mathbf{C}_3, \mathbf{C}_4, \mathbf{C}_5, \mathbf{C}_6][-1, 0, 2, 2, 2, 2]^T &= \mathbf{0} \\ \Rightarrow \mathbf{A}[-1, 0, 2, 2, 2, 2]^T &= \mathbf{0}.\end{aligned}$$

(0.5 marks)

- (B) i) The problem is not linearly separable as its difficult to find the linear decision boundary which separates the two classes. (1 mark)

- ii) The Kernel is defined as: $K(x_1, x_2) = (4 + x_1^T x_2)^2$ (0.5 mark)

The Kernel matrix is defined as

$$K = \begin{pmatrix} 144 & 16 & 16 & 16 \\ 16 & 144 & 16 & 16 \\ 16 & 16 & 144 & 16 \\ 16 & 16 & 16 & 144 \end{pmatrix}$$

(1 mark)

- iii)

$$Q(\alpha) = \sum_{i=1}^4 \alpha_i - \frac{1}{2} \left(\sum_{i=1}^4 144\alpha_i^2 + 32\alpha_1\alpha_2 + 32\alpha_3\alpha_4 - 32\alpha_1\alpha_3 - 32\alpha_1\alpha_4 - 32\alpha_2\alpha_3 - 32\alpha_2\alpha_4 \right)$$

$$\frac{\partial Q(\alpha)}{\partial \alpha_i} = 0$$

We get system of equations:

$$144\alpha_1 + 16\alpha_2 - 16\alpha_3 - 16\alpha_4 = 1$$

$$16\alpha_1 + 144\alpha_2 - 16\alpha_3 - 16\alpha_4 = 1$$

$$-16\alpha_1 - 16\alpha_2 + 144\alpha_3 + 16\alpha_4 = 1$$

$$-16\alpha_1 - 16\alpha_2 + 16\alpha_3 + 144\alpha_4 = 1$$

solving above equations we get $\alpha_i = \frac{1}{128}$ (0.5 marks)

$w = (0, 0, 22.6, 0, 0, 0)$ and $b = 0$

The decision boundary is given by

$$w^T \phi(x) + b = 0$$

$$x_1 x_2 = 0 \quad (2 \text{ marks})$$

Q4

- (A) i) The Lagrangian is given by $L(x, \lambda) = x_1^2 + x_2^2 - 4x_1 - 4x_2 + \lambda_1(x_1^2 - x_2) + \lambda_2(x_1 + x_2 - 2)$
 $\frac{\partial L}{\partial x_1} = 0, \frac{\partial L}{\partial x_2} = 0$

$$x_1 = \frac{(4-\lambda_2)}{2(1+\lambda_1)} \text{ and } x_2 = \frac{\lambda_1-\lambda_2+4}{2} \quad (1 \text{ mark})$$

Substituting above x_1 and x_2 values in $L(x, \lambda)$

$$L(\lambda) = \frac{(4-\lambda_2)^2}{4(1+\lambda_1)^2} + \frac{(\lambda_1-\lambda_2+4)^2}{4} - 4\frac{(4-\lambda_2)}{2(1+\lambda_1)} - 4\frac{\lambda_1-\lambda_2+4}{2} + \lambda_1(\frac{(4-\lambda_2)^2}{4(1+\lambda_1)^2} - \frac{\lambda_1-\lambda_2+4}{2}) + \lambda_2(\frac{(4-\lambda_2)}{2(1+\lambda_1)} + \frac{\lambda_1-\lambda_2+4}{2} - 2) \quad (1 \text{ mark})$$

Dual is given by: $\max_{\lambda} L(\lambda)$

- ii) Dual and primal objective function value will be the same as the primal objective function is convex and the constraints are convex. (2 marks)
- (B) i) the dimension of each training sample is 1×6 as the covariance matrix is having 6 eigen values so that means each training sample will have 6 columns. (1 mark)
- ii) total variance= sum of all eigen values=23.330 if we want to retain 95% variance then we should keep top 3 eigen values as
 $(12 + 6.8 + 3.5) * 100 / (23.330) = 95.7$ (1.5 marks)

if we want to retain 99% variance then we should keep top 4 eigen values as $(12 + 6.8 + 3.5 + 1) * 100 / (23.330) = 99.87$ (1.5 marks)

iii) After PCA the result will be $[-3 \quad -2 \quad 2 \quad 1 \quad 2 \quad 4] * \begin{pmatrix} 0.115 & 0.205 \\ 0.106 & 0.215 \\ 0.118 & 0.315 \\ 0.082 & 0.428 \\ 0.013 & 0.238 \\ 0.023 & 0.034 \end{pmatrix} =$
 $[-0.1210 \quad 0.6250]$ (2 marks)