

Central Ohio Orthopedic Center

4900 Reed Road, Columbus, OH 43220

Operative Report

Patient	Henry J. McMillan
MRN	0048-77-215
Date of procedure	March 5, 2019
Surgeon	Mark T. Holloway, MD
Anesthesia	General
Preoperative diagnosis	Left knee medial meniscus tear
Postoperative diagnosis	Left knee medial meniscus tear, confirmed
Procedure	Left knee arthroscopy with partial medial meniscectomy

Indication

The patient is a 49 year old man, a former collegiate offensive lineman and current high school football coach, with several months of left knee pain, catching, and swelling that failed conservative care. MRI showed a tear of the medial meniscus. He elected to proceed with arthroscopic treatment. This is his second knee arthroscopy. The right knee was treated in 2016.

Findings

Diagnostic arthroscopy revealed a complex tear of the posterior horn of the medial meniscus. Articular cartilage showed grade 2 changes in the medial compartment consistent with early wear. The anterior cruciate ligament was intact. The lateral compartment was unremarkable.

Procedure

Standard arthroscopic portals were established. The unstable meniscal fragment was debrided to a stable rim with an arthroscopic shaver and biters. The joint was irrigated. Portals were closed with nylon suture and a sterile dressing was applied. The patient tolerated the procedure well and was transferred to recovery in stable condition.

Plan

Weight bearing as tolerated with crutches for comfort. Physical therapy to begin within one week. Follow up in the office in 10 to 14 days for suture removal. Expect return to full activity over 4 to 6 weeks.

Dictated and electronically signed by Mark T. Holloway, MD, March 5, 2019.