## TUTORIAL 05

1. Let X be a random variable with probability density function

$$f(x) = \begin{cases} c(1-x)^2 ; -1 < x < 1 \\ 0 ; \text{ otherwise} \end{cases}$$

- (a) What is the value of c?
- (b) What is the cumulative distribution function of X?
- 2. Let X be uniformly distributed over  $(\alpha, \beta)$ . Find (a) E[X] and (b) Var(X).
- 3. Buses arrive at a specified stop at 15-minute intervals starting at 7 A.M. That is, they arrive at 7, 7:15, 7:30, 7:45, and so on. If a passenger arrives at the stop at a time that is uniformly distributed between 7 and 7:30, find the probability that he waits, (a) less than 5 minutes for a bus and (b) more than 10 minutes for a bus.
- 4. Consider the function

$$f(x) = \begin{cases} C(2x - x^3) & ; \ 0 < x < \frac{5}{2} \\ 0 & ; \ \text{otherwise} \end{cases}$$

Could f be a probability density function? If so, determine C. Repeat if f(x) were given by

$$f(x) = \begin{cases} C(2x - x^2) & ; \ 0 < x < \frac{5}{2} \\ 0 & ; \ \text{otherwise} \end{cases}$$

5. The probability density function of X, the lifetime of a certain type of electronic device (measured in hours), is given by

$$f(x) = \begin{cases} \frac{10}{x^2} & : x > 10\\ 0 & x \le 10 \end{cases}$$

- (a) Find  $P\{X > 20\}$ .
- (b) What is the cumulative distribution function of X?
- (c) What is the probability that, of 6 such types of devices, at least 3 will function for at least 15 hours? What assumptions are you making?
- 6. Suppose that if you are s minutes early for an appointment, then you incur the cost cs, and if you are s minutes late, then you incur the cost ks. Suppose also that the travel time from where you presently are to the location of your appointment is a continuous random variable having the probability density function f. Determine the time at which you should depart if you want to minimize your expected cost.

1

2

7. If X is a normal variable with parameters  $\mu = 10$  and  $\sigma^2 = 36$ , compute (a)  $P\{X > 5\}$ , (b)  $P\{4 < X < 16\}$ , (c)  $P\{X < 8\}$ , (d)  $P\{X < 2\}$  and (e)  $P\{X > 16\}$ .

TUTORIAL 05

- 8. If X is a normal random variable with parameters  $\mu = 3$  and  $\sigma^2 = 9$ , find (a)  $P\{2 < X < 5\}$ , (b)  $P\{X > 0\}$  and (c)  $P\{|X 3| > 6\}$ .
- 9. A randomly chosen IQ test taker obtains a score that is approximately a normal random variable with mean 100 and standard deviation 15. What is the probability that the score of such a person is (a) above 125; (b) between 90 and 110?
- 10. Suppose that the travel time from your home to your office is normally distributed with mean 40 minutes and standard deviation 7 minutes. If you want to be 95 percent certain that you will not be late for an office appointment at 1 P.M., what is the latest time that you should leave home?
- 11. The life of a certain type of automobile tire is normally distributed with mean 34,000 miles and standard deviation 4000 miles.
  - (1) What is the probability that such a tire lasts over 40,000 miles?
  - (b) What is the probability that if lasts between 30,000 and 35,000 miles?
  - (c) Given that it has survived 30,000 miles, what is the conditional probability that the tire survives another 10,000 miles?
- 12. The annual rainfall in Cleveland, Ohio is approximately a normal random variable with mean 40.2 inches and standard deviation 8.4 inches. What is the probability that
  - (a) next year's rainfall will exceed 44 inches?
  - (b) the yearly rainfall in exactly 3 of the next 7 years will exceed 44 inches?