

The Threshold Theorem of Epidemiology

1. A small group of people introduce an infectious disease in a large population.
2. The disease has a short incubation period.
3. Recovered individuals gain permanent immunity.

There are three classes of population. They are:

- i) $x \rightarrow$ The infected class, ii) $y \rightarrow$ The susceptible class.
- iii) $z \rightarrow$ The removed class (recovered class).

Rule 1: $x(t) + y(t) + z(t) = N$, where N is the fixed total number of population.
(Conserved Condition)

Rule 2: $\frac{dy}{dt} \propto xy \Rightarrow \frac{dy}{dt} = -Axy$ $A \rightarrow$ The infection rate.

Rule 3: $\frac{dz}{dt} \propto x \Rightarrow \frac{dz}{dt} = Bx$ $B \rightarrow$ The removal rate.

Using Rule 1 we can write $\frac{dx}{dt} = -\frac{dy}{dt} - \frac{dz}{dt}$, which gives $\frac{dx}{dt} = Axy - Bx$. In all the three $\frac{dx}{dt}$, $\frac{dy}{dt}$ and $\frac{dz}{dt}$ equations, the right hand side does not depend on z . This is a pseudo-third-order system.

I.) The x-y equation:

$$\frac{dx/dt}{dy/dt} = \frac{dx}{dy} = \frac{Axy - Bx}{-Axy} = -1 + \frac{B}{Ay}$$

$$\Rightarrow \int dx = \int \left(\frac{B}{Ay} - 1 \right) dy \Rightarrow \boxed{x = \frac{B}{A} \ln y - y + C_1}$$

Initial Condition: At $\boxed{t=0}$ (initially), $\boxed{x=x_0}$,
 $\boxed{y=y_0}$ and $\boxed{z=0}$. $\Rightarrow \boxed{x_0 + y_0 = N}$.

$$\therefore \boxed{C_1 = x_0 + y_0 - \frac{B}{A} \ln y_0 = N - \frac{B}{A} \ln y_0}$$

$$\Rightarrow \boxed{x = (x_0 + y_0) - y + \frac{B}{A} \ln(y/y_0)} \quad \boxed{x \equiv x(y)} \text{ in close form.}$$

II.) The y-z equation:

$$\frac{dy/dt}{dz/dt} = \frac{dy}{dz} = \frac{-Axy}{Bx} = -\frac{Ay}{B}$$

$$\Rightarrow \int \frac{dy}{y} = -\int \frac{A}{B} dz \Rightarrow \boxed{\ln y = -\frac{Az}{B} + C_2}$$

At $\boxed{t=0}$, $\boxed{y=y_0}$ and $\boxed{z=0} \Rightarrow \boxed{C_2 = \ln y_0}$

$$\Rightarrow \boxed{\ln\left(\frac{y}{y_0}\right) = -\frac{A}{B} z} \text{ for } t > 0, \boxed{y < y_0}$$

$$\Rightarrow \boxed{\ln\left(\frac{y}{y_0}\right) < 0} \text{ for } t > 0 \Rightarrow \boxed{z = -\frac{B}{A} \ln\left(\frac{y}{y_0}\right) > 0} \text{ for } t > 0.$$

And $\boxed{y = y_0 \exp\left(-\frac{A}{B} z\right)} \quad \boxed{y \equiv y(z)} \text{ in close form.}$

III.) The z-x equation :

$$\boxed{\frac{dx/dt}{dz/dt} = \frac{dx}{dz} = \frac{Axy - Bx}{Bx} = \frac{Ay}{B} - 1}$$

$$\text{Now } \boxed{y = y_0 e^{-\frac{Az}{B}}} \Rightarrow \boxed{\frac{dx}{dz} = \frac{A}{B} y_0 e^{-\frac{Az}{B}} - 1}$$

$$\Rightarrow x = \int \frac{A}{B} y_0 e^{-\frac{A}{B} z} dz - \int 1 dz + C_3$$

$$\Rightarrow x = \frac{A}{B} y_0 \frac{e^{-\frac{Az}{B}}}{-A/B} - z + C_3$$

$$\Rightarrow \boxed{x = -y_0 e^{-Az/B} - z + C_3}$$

When (at $t=0$), $\boxed{x = x_0}$ and $\boxed{z = 0}$,

$$\boxed{C_3 = x_0 + y_0} \Rightarrow \boxed{x = x_0 + y_0(1 - e^{-Az/B}) - z}$$

OR $\boxed{x = (x_0 + y_0) - y_0 \exp(-\frac{Az}{B}) - z}$ But z is not written in a closed form for x .

Plot of $x-y$: $\boxed{x = (x_0 + y_0) - y + \frac{B}{A} \ln\left(\frac{y}{y_0}\right)}$

i.) When $y \rightarrow 0$, $x \rightarrow -\infty$.


ii.) $\boxed{\frac{dx}{dy} = x(Ay - B)}$ $\Rightarrow \boxed{\frac{dx}{dy} = 0}$, when either $\boxed{x = 0}$ or $\boxed{y = B/A}$.

iii.) $\boxed{\frac{dx}{dy} = \frac{B}{Ay} - 1}$ $\Rightarrow \boxed{\frac{dx}{dy} = 0}$, when $\boxed{y = \frac{B}{A} = p(\text{say})}$

iv.) $\boxed{\frac{d^2x}{dy^2} = -\frac{B}{Ay^2}}$ At $\boxed{y = \frac{B}{A}}$, $\boxed{\frac{d^2x}{dy^2} = -\frac{A}{B} < 0}$. Hence, $y = B/A$ is a maximum.

v) When $x=0$, write $y=y_0$ (say).

With $x=0$, the point becomes an equilibrium point, since both $\frac{dx}{dt} = \frac{dy}{dt} = 0$.



Conclusion 1: An epidemic will breakout, if $(y_0 > p)$, i.e. the initial number of susceptibles are above the threshold p . This is the Threshold Theorem of Epidemiology (Kermack & McKendrick).

Conclusion 2: The spread of the disease stops ($x=0$) because the infective population is reduced to zero, even though there ~~may~~ may be some susceptibles left.

Practically speaking epidemics breakout due to overcrowding of a susceptible population in an unhygienic environment.

Case of Initial Number of Susceptibles Slightly Higher than the Threshold

$$\boxed{\frac{dx}{dt} = x(Ay - B)} \text{ and } \boxed{x = (x_0 + y_0) - y + \frac{B}{A} \ln\left(\frac{y}{y_0}\right)}$$

When $\boxed{\frac{dx}{dt} = 0}$ Either $\boxed{y = B/A = p}$ or $\boxed{x = 0}$.

i) The former case is a turning point in which $y = p$ is a maximum $\Rightarrow \boxed{d^2x/dt^2 < 0}$.

ii) $\boxed{\frac{d^2x}{dt^2} = \frac{dx}{dt}(Ay - B) + x \frac{d}{dt}(Ay - B)}$. When $\boxed{x = 0}$

at $\boxed{\frac{dx}{dt} = 0} \Rightarrow \boxed{\frac{d^2x}{dt^2} = 0}$. This is an equilibrium point. (at $x = 0$).

When $\boxed{x = 0}$, we write $\boxed{y = y_\infty}$. With these,

We get $\boxed{0 = x_0 + y_0 - y_\infty + \frac{B}{A} \ln\left(\frac{y_\infty}{y_0}\right)}$. Usually

$\boxed{x_0 \ll y_0}$, and so we neglect x_0 above.

Now y_0 is the initial value of the Susceptibles, and y_∞ is the final value of the Susceptibles.

$\therefore y_0 - y_\infty$ is the number of people who have contracted the infectious disease. Further,

$\boxed{y_\infty < y_0}$, which implies that $\boxed{y_0 - y_\infty > 0}$.

Hence we can write $\boxed{(y_0 - y_\infty) + \frac{B}{A} \ln\left(\frac{y_0 - y_0 + y_\infty}{y_0}\right) \approx 0}$

$\Rightarrow \boxed{(y_0 - y_\infty) + p \ln\left(1 - \frac{y_0 - y_\infty}{y_0}\right) \approx 0}$ in which we have neglected x_0 .

We now consider the case where the initial number of susceptibles is slightly greater than the threshold, i.e. $y_0 = p + \epsilon$, in which $\epsilon \ll p$. $\therefore \frac{(y_0 - p)}{p} = \frac{\epsilon}{p} \ll 1$.

From the plot of $x-y$ we can see that in this ~~case~~ case $(y_0 - y_\infty) \ll y_0$. Hence

using the formula $\ln(1+u) = u - \frac{u^2}{2} + \frac{u^3}{3} - \dots$,

we can write $\ln\left[1 - \left(\frac{y_0 - y_\infty}{y_0}\right)\right] = -\left(\frac{y_0 - y_\infty}{y_0}\right) - \frac{1}{2}\left(\frac{y_0 - y_\infty}{y_0}\right)^2$

Going only up to the second order term.

Hence, $(y_0 - y_\infty) + p\left[-\left(\frac{y_0 - y_\infty}{y_0}\right) - \frac{1}{2}\left(\frac{y_0 - y_\infty}{y_0}\right)^2\right] \approx 0$

$$\Rightarrow (y_0 - y_\infty) \left[1 - \frac{p}{y_0} - \frac{p}{2} \frac{(y_0 - y_\infty)}{y_0^2}\right] \approx 0 \quad (\text{Now, } y_0 - y_\infty \neq 0)$$

$$\Rightarrow y_0 - y_\infty \approx \frac{2y_0^2}{p} \left(1 - \frac{p}{y_0}\right) = 2y_0 \left(\frac{y_0}{p} - 1\right)$$

But $y_0 = p + \epsilon$ and $\left(\frac{y_0}{p} - 1\right) = \frac{\epsilon}{p}$. Using these

we get, $y_0 - y_\infty \approx 2y_0 \epsilon = 2(p + \epsilon) \epsilon / p$

Neglecting ϵ in $[p + \epsilon \ll p]$, we finally get.

$$y_0 - y_\infty \approx 2p \frac{\epsilon}{p} \Rightarrow y_0 - y_\infty = 2\epsilon$$

$\therefore y_0 - y_\infty \approx 2(y_0 - p)$, an approximate result that is valid only when y_0 is slightly greater than p .

$Z(t)$ as a Function of Time : $Z \equiv Z(t)$.

$$\boxed{\frac{dz}{dt} = Bx} , \quad \boxed{x = (x_0 + y_0) - y + \frac{B}{A} \ln\left(\frac{y}{y_0}\right)} \text{ and}$$

$$\boxed{y = y_0 \exp\left(-\frac{AZ}{B}\right)} \Rightarrow \boxed{Z = -\frac{B}{A} \ln\left(y/y_0\right)}.$$

Hence, $\boxed{\frac{dz}{dt} = B \left[x_0 + y_0 - y_0 \exp\left(-\frac{A}{B} Z\right) - Z \right]}$,

an equation in an autonomous form $\boxed{\frac{dz}{dt} = f(z)}$.

Noting $\boxed{\frac{B}{A} = p}$, we consider the case of $\boxed{Z \ll p}$.

$$\therefore \boxed{e^{-Z/p} \approx 1 - \frac{Z}{p} + \frac{1}{2} \left(\frac{Z}{p}\right)^2 + \dots}$$
, bring up to the second order.

$$\Rightarrow \frac{dz}{dt} \approx B \left[x_0 + \cancel{y_0} - \cancel{y_0} + y_0 \frac{Z}{p} - \frac{y_0}{2} \left(\frac{Z}{p}\right)^2 - Z \right]$$

$$\Rightarrow \frac{dz}{dt} = B \left[x_0 + \left(\frac{y_0}{p} - 1\right) Z - \frac{y_0}{2} \left(\frac{Z}{p}\right)^2 \right].$$

We define $\boxed{\beta = Z/p}$, by which ~~we~~ we set,

$$\boxed{\frac{p}{B} \frac{d\beta}{dt} = x_0 + (y_0 - p)\beta - \frac{y_0}{2} \beta^2}.$$

Multiplying throughout by ~~we~~ $\frac{2}{y_0}$ we get.

$$\boxed{\frac{2}{y_0} \frac{p}{B} \frac{d\beta}{dt} = \frac{2x_0}{y_0} + 2\left(1 - \frac{p}{y_0}\right)\beta - \beta^2}.$$

We further define $\boxed{\alpha^2 = 2x_0/y_0}$ and $\boxed{\beta^* = 1 - \frac{p}{y_0}}$.

Hence $\frac{2}{y_0} \frac{\ell}{B} \frac{d\beta}{dt} = -(\beta^2 - 2\beta\beta + \beta^2 - \beta^2) + \alpha^2$

$\Rightarrow \boxed{\frac{2}{y_0 A} \frac{d\beta}{dt} = (\beta^2 + \alpha^2) - (\beta - \beta)^2} \because \frac{\ell}{B} = \frac{B}{A}$

Further define $\boxed{k^2 = \alpha^2 + \beta^2}$ and $\boxed{\xi = \beta - \beta}$,

which gives $\boxed{\frac{d\beta}{dt} = \frac{d\xi}{dt}}$. Using this we get,

$\boxed{\frac{2}{Ay_0} \frac{d\xi}{dt} = k^2 - \xi^2} \Rightarrow \frac{2}{Ay_0 k} \frac{d}{dt} \left(\frac{\xi}{k} \right) = 1 - \left(\frac{\xi}{k} \right)^2$

Again, defining $\boxed{\psi = \xi/k}$ and $\boxed{\tau = \frac{Ay_0 k}{2} t}$,

we obtain finally, $\boxed{\frac{d\psi}{d\tau} = 1 - \psi^2}$. This

Equation can be integrated to obtain.

$\boxed{\ln \left(\frac{1+\psi}{1-\psi} \right) = 2\tau + 2C}$, where C is an integration constant.

(Check for the solution of $\boxed{\frac{dx}{d\tau} = 1 - x^2}$ equation)

Writing $\boxed{T = \tau + C}$, we get, $\boxed{\frac{1+\psi}{1-\psi} = e^{2T}}$, from

which we get $1+\psi = (1-\psi)e^{2T} \Rightarrow \psi(1+e^{2T}) = e^{2T} - 1$

$\Rightarrow \boxed{\psi = \frac{e^{2T} - 1}{e^{2T} + 1}} \Rightarrow \boxed{\psi = \frac{e^T - e^{-T}}{e^T + e^{-T}} = \tanh(T)}$

$\therefore \boxed{\frac{\xi}{k} = \tanh(\tau + C)} \Rightarrow \boxed{\frac{\beta - \beta}{k} = \tanh(\tau + C)}$

$$\Rightarrow \beta = \beta + k \tanh(\tau + c)$$

$$\Rightarrow \frac{Z}{p} = \beta + k \tanh\left(\frac{A y_0 k}{2} t + c\right)$$

$$\Rightarrow \boxed{Z = \beta p + k p \tanh\left(\frac{A y_0 k}{2} t + c\right)} \quad \boxed{Z \equiv Z(t)}$$


Hence, $\frac{dZ}{dt} = k p \cdot \left(\frac{A y_0 k}{2}\right) \operatorname{sech}^2\left(\frac{A y_0 k}{2} t + c\right)$

$$\Rightarrow \boxed{\frac{dZ}{dt} = \frac{B y_0 k^2}{2} \operatorname{sech}^2\left(\frac{A y_0 k}{2} t + c\right)}$$

The integration constant c can be fixed from the initial condition, $\boxed{Z=0}$ at $\boxed{t=0}$.

$$\Rightarrow \boxed{0 = \beta p + k p \tanh(c)} \Rightarrow \boxed{\tanh(c) = -\frac{\beta}{k}}$$

Now $\boxed{\beta = 1 - \frac{p}{y_0}} \therefore \boxed{\tanh(c) = \frac{1}{k} \left(\frac{p}{y_0} - 1\right)}$



i) The epidemic curve reaches a maximum when $\boxed{\frac{A y_0 k}{2} t + c = 0}$.

$$\Rightarrow \boxed{t = -2c / A k y_0} \quad (c < 0)$$

ii) The removal rate dZ/dt climbs to a peak value and dies out.

iii) Data matched by the Bombay plague of 1905.