Class 10

Shivani Khosla (PID: A59010433)

10/29/2021

```
candy_file <- "https://raw.githubusercontent.com/fivethirtyeight/data/master/candy-power-ranking/candy-
candy = read.csv(candy_file, row.names=1)
head(candy)
```

```
##              chocolate fruity caramel peanutyalmondy nougat crispedricewafer
## 100 Grand            1      0       1              0      0                1
## 3 Musketeers         1      0       0              0      1                0
## One dime             0      0       0              0      0                0
## One quarter          0      0       0              0      0                0
## Air Heads            0      1       0              0      0                0
## Almond Joy           1      0       0              1      0                0
##              hard bar pluribus sugarpercent pricepercent winpercent
## 100 Grand       0   1        0        0.732        0.860   66.97173
## 3 Musketeers    0   1        0        0.604        0.511   67.60294
## One dime        0   0        0        0.011        0.116   32.26109
## One quarter     0   0        0        0.011        0.511   46.11650
## Air Heads       0   0        0        0.906        0.511   52.34146
## Almond Joy      0   1        0        0.465        0.767   50.34755
```

```
rownames(candy) <- gsub("Õ", "'", rownames(candy))
```

Q1. How many different candy types are in this dataset?

```
nrow(candy)
```

```
## [1] 85
```

Q2. How many fruity candy types are in the dataset?

```
sum(candy$fruity)
```

```
## [1] 38
```

```
candy["Twix", ]$winpercent
```

```
## [1] 81.64291
```

Q3. What is your favorite candy in the dataset and what is it's winpercent value?

```
candy["Almond Joy", ]$winpercent
```

```
## [1] 50.34755
```

Q4. What is the winpercent value for “Kit Kat”?

```
candy["Kit Kat", ]$winpercent
```

```
## [1] 76.7686
```

Q5. What is the winpercent value for “Tootsie Roll Snack Bars”?

```
candy["Tootsie Roll Snack Bars", ]$winpercent
```

```
## [1] 49.6535
```

```
library("skimr")
skim(candy)
```

Table 1: Data summary

Name	candy
Number of rows	85
Number of columns	12
Column type frequency:	
numeric	12
Group variables	None

Variable type: numeric

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100	hist
chocolate	0	1	0.44	0.50	0.00	0.00	0.00	1.00	1.00	
fruity	0	1	0.45	0.50	0.00	0.00	0.00	1.00	1.00	
caramel	0	1	0.16	0.37	0.00	0.00	0.00	0.00	1.00	
peanutyalmondy	0	1	0.16	0.37	0.00	0.00	0.00	0.00	1.00	
nougat	0	1	0.08	0.28	0.00	0.00	0.00	0.00	1.00	
crispedricewafer	0	1	0.08	0.28	0.00	0.00	0.00	0.00	1.00	
hard	0	1	0.18	0.38	0.00	0.00	0.00	0.00	1.00	
bar	0	1	0.25	0.43	0.00	0.00	0.00	0.00	1.00	
pluribus	0	1	0.52	0.50	0.00	0.00	1.00	1.00	1.00	
sugarpercent	0	1	0.48	0.28	0.01	0.22	0.47	0.73	0.99	
pricepercent	0	1	0.47	0.29	0.01	0.26	0.47	0.65	0.98	
winpercent	0	1	50.32	14.71	22.45	39.14	47.83	59.86	84.18	

Q6. Is there any variable/column that looks to be on a different scale to the majority of the other columns in the dataset?

Columns 10-12 are not binary values since they are percentages

Q7. What do you think a zero and one represent for the candy$chocolate column?

They are yes(1) or no(0) values.

Q8. Plot a histogram of winpercent values

```
hist(candy$winpercent)
```



Q9. Is the distribution of winpercent values symmetrical?

The distribution is skewed right

Q10. Is the center of the distribution above or below 50%?

The center is below 50%.

Q11. On average is chocolate candy higher or lower ranked than fruit candy?

```
chocolate <- candy[as.logical(candy$chocolate), ]$winpercent
mean(chocolate)
```

```
## [1] 60.92153
```

```
fruity <- candy[as.logical(candy$fruity), ]$winpercent
mean(fruity)
```

```
## [1] 44.11974
```

Q12. Is this difference statistically significant?

```
t.test(chocolate, fruity)
```

```
##
##  Welch Two Sample t-test
##
## data:  chocolate and fruity
## t = 6.2582, df = 68.882, p-value = 2.871e-08
## alternative hypothesis: true difference in means is not equal to 0
## 95 percent confidence interval:
##  11.44563 22.15795
## sample estimates:
## mean of x mean of y
##  60.92153  44.11974
```

p = 2.871e-08 so the difference is statistically significant

Q13. What are the five least liked candy types in this set?

```
head(candy[order(candy$winpercent), ], n = 5)
```

```
##                    chocolate fruity caramel peanutyalmondy nougat
## Nik L Nip                  0      1       0              0      0
## Boston Baked Beans         0      0       0              1      0
## Chiclets                   0      1       0              0      0
## Super Bubble               0      1       0              0      0
## Jawbusters                 0      1       0              0      0
##                    crispedricewafer hard bar pluribus sugarpercent pricepercent
## Nik L Nip                         0    0   0        1        0.197        0.976
## Boston Baked Beans                0    0   0        1        0.313        0.511
## Chiclets                          0    0   0        1        0.046        0.325
## Super Bubble                      0    0   0        0        0.162        0.116
## Jawbusters                        0    1   0        1        0.093        0.511
##                    winpercent
## Nik L Nip            22.44534
## Boston Baked Beans   23.41782
## Chiclets             24.52499
## Super Bubble         27.30386
## Jawbusters           28.12744
```

Q14. What are the top 5 all time favorite candy types out of this set?

```
head(candy[order(candy$winpercent, decreasing = TRUE), ], n = 5)
```

```
##                          chocolate fruity caramel peanutyalmondy nougat
## Reese's Peanut Butter cup         1      0       0              1      0
## Reese's Miniatures                1      0       0              1      0
## Twix                              1      0       1              0      0
## Kit Kat                           1      0       0              0      0
## Snickers                          1      0       1              1      1
##                          crispedricewafer hard bar pluribus sugarpercent
## Reese's Peanut Butter cup                0    0   0        0        0.720
## Reese's Miniatures                       0    0   0        0        0.034
## Twix                                     1    0   1        0        0.546
## Kit Kat                                  1    0   1        0        0.313
## Snickers                                 0    0   1        0        0.546
##                          pricepercent winpercent
## Reese's Peanut Butter cup        0.651   84.18029
## Reese's Miniatures               0.279   81.86626
## Twix                             0.906   81.64291
## Kit Kat                          0.511   76.76860
## Snickers                         0.651   76.67378
```

Reese's Peanut Butter cup, Reese's Miniatures, Twix, Kit Kat, Snickers

Q15. Make a first barplot of candy ranking based on winpercent values.

```
library(ggplot2)
rownames(candy) <- gsub("Õ", "'", rownames(candy))
ggplot(candy) +
  aes(winpercent, rownames(candy)) +
  geom_col()
```



> Q16. This is quite ugly, use the reorder() function to get the bars sorted by winpercent?

```
library(ggplot2)

ggplot(candy) +
  aes(winpercent, reorder(rownames(candy),winpercent)) +
  geom_col()
```



for colors:

```
my_cols=rep("black", nrow(candy))
my_cols[as.logical(candy$chocolate)] = "chocolate"
my_cols[as.logical(candy$bar)] = "brown"
my_cols[as.logical(candy$fruity)] = "pink"
```

incorporate colors

```
ggplot(candy) +
  aes(winpercent, reorder(rownames(candy),winpercent)) +      geom_col(fill=my_cols)
```



> Q17. What is the worst ranked chocolate candy?

Sixlets

Q18. What is the best ranked fruity candy?

Starburst

```
library(ggrepel)

# How about a plot of price vs win
ggplot(candy) +
  aes(winpercent, pricepercent, label=rownames(candy)) +
  geom_point(col=my_cols) +
  geom_text_repel(col=my_cols, size=3.3, max.overlaps = 5)
```

```
## Warning: ggrepel: 54 unlabeled data points (too many overlaps). Consider
## increasing max.overlaps
```



Q19. Which candy type is the highest ranked in terms of winpercent for the least money - i.e. offers the most bang for your buck?

Reese's miniatures

Q20. What are the top 5 most expensive candy types in the dataset and of these which is the least popular?

```
head(candy[order(candy$pricepercent, decreasing = TRUE), ], n = 5)
```

```
##                         chocolate fruity caramel peanutyalmondy nougat
## Nik L Nip                       0      1       0              0      0
## Nestle Smarties                 1      0       0              0      0
## Ring pop                        0      1       0              0      0
## Hershey's Krackel               1      0       0              0      0
## Hershey's Milk Chocolate        1      0       0              0      0
##                         crispedricewafer hard bar pluribus sugarpercent
## Nik L Nip                              0    0   0        1        0.197
## Nestle Smarties                        0    0   0        1        0.267
## Ring pop                               0    1   0        0        0.732
## Hershey's Krackel                      1    0   1        0        0.430
## Hershey's Milk Chocolate               0    0   1        0        0.430
##                         pricepercent winpercent
## Nik L Nip                      0.976   22.44534
```

```
## Nestle Smarties                  0.976   37.88719
## Ring pop                         0.965   35.29076
## Hershey's Krackel                0.918   62.28448
## Hershey's Milk Chocolate         0.918   56.49050
```

Nik L Nip, Nestle Smarties, Ring pop, Hershey's Krackel, Hershey's Milk Chocolate

least popular: Nik L Nip

```
library(corrplot)
```

```
## corrplot 0.90 loaded
```

```
cij <- cor(candy)
corrplot(cij)
```



Q22. Examining this plot what two variables are anti-correlated (i.e. have minus values)?

chocolate and fruity

Q23. Similarly, what two variables are most positively correlated?

chocolate and winpercent

```
pca <- prcomp(candy, scale = TRUE)
summary(pca)
```

```
## Importance of components:
##                           PC1    PC2    PC3     PC4    PC5     PC6     PC7
## Standard deviation     2.0788 1.1378 1.1092 1.07533 0.9518 0.81923 0.81530
## Proportion of Variance 0.3601 0.1079 0.1025 0.09636 0.0755 0.05593 0.05539
## Cumulative Proportion  0.3601 0.4680 0.5705 0.66688 0.7424 0.79830 0.85369
##                            PC8     PC9    PC10    PC11    PC12
## Standard deviation     0.74530 0.67824 0.62349 0.43974 0.39760
## Proportion of Variance 0.04629 0.03833 0.03239 0.01611 0.01317
## Cumulative Proportion  0.89998 0.93832 0.97071 0.98683 1.00000
```

```
plot(pca$x[,1:2])
```




```
plot(pca$x[,1:2], col=my_cols, pch=16)
```




```
# Make a new data-frame with our PCA results and candy data
my_data <- cbind(candy, pca$x[,1:3])
```

```
p <- ggplot(my_data) +
        aes(x=PC1, y=PC2,
            size=winpercent/100,
            text=rownames(my_data),
            label=rownames(my_data)) +
        geom_point(col=my_cols)

p
```




```
library(ggrepel)

#p + geom_text_repel(size=3.3, col=my_cols, max.overlaps = 7)  +
#  theme(legend.position = "none") +
#  labs(title="Halloween Candy PCA Space",
#       subtitle="Colored by type: chocolate bar (dark brown), chocolate other (light brown), fruity (r
#       caption="Data from 538")
```

```
library(plotly)
```

```
##
## Attaching package: 'plotly'

## The following object is masked from 'package:ggplot2':
##
##     last_plot

## The following object is masked from 'package:stats':
##
##     filter

## The following object is masked from 'package:graphics':
##
##     layout
```

```
#ggplotly(p)
```

```
par(mar=c(8,4,2,2))
barplot(pca$rotation[,1], las=2, ylab="PC1 Contribution")
```



> Q24. What original variables are picked up strongly by PC1 in the positive direction? Do these make sense to you?

fruity, hard, pluribus. We discussed that this makes sense since fruity candies tend to be hard and come packaged with many other fruity candies.