

ACC SANDOZ (200 MG)

ACETYLCYSTEINE 200 MG EFFERVESCENT TABLETS

.. HOW TO USE : ONE TABLETSS DISSOLVE IN 100 ML OF WATER ..

- TO DISSOLVE MUCUS IN THE RESPIRATORY TRACT. RELIEVE COUGH WITH MUCUS
- ANTIDOTE FOR PARACETAMOL POISONING



DOSAGE REGIMEN

- > AGE OF 14 : 1 TAB 2-3 TIMES DAILY
- AGE OF 6 – 14 : 1 TAB TWICE DAILY
- AGE OF 2 – 5 : ½ TAB 2-3 TIMES DAILY

NO ADJUSTMENT NEEDED IN RENAL/HEPATIC IMPAIRED PATIENT





INCRAESE BRONCHIAL SECRETION

HEADACHE VASOMOTOR RHINITIS

ITCHING LYELL'S SYNDROME

STEVENS-JOHNSON SYNDROME



PRECAUTIONS



- AVOID USING FOR LONG TERM THERAPY IN PATIENT WITH HISTAMINE INTOLERANCE
- 20 MG OF SORBITOL AND 70 MG OF LACTOSE PER TABLET

AVOID USING IN HFI, TOTAL LACTASE DEFICIENCY

POSSIBLE DRUG INTERACTIONS

- TAKE 2 HRS APART FROM TETRACYCLINES, AMINOGLYCOSIDES, PENICILLINS, ACTIVATED CHARCOAL, NITROGLYCERINE

CONTRAINDICATION



- HYPERSENSITIVITY TO ACETYLCYSTEINE OR TO ANY OF THE EXCIPIENTS
- MUST NOT BE USED IN CHILDREN < 2 YEARS OLD

PREGNANCY & LACTATION

- NO ADEQUATE CLINICAL DATA SUPPORT

