OLICOXIB

PARECOXIB

(40MG INJECTION)





Generic name : Parecoxib 40 mg

Route : Injection

INDICATION

- Management of acute pain and for short-term treatment of post-operative pain
- For patient with need for parenteral therapy and for whom a similar benefit could not be obtained from alternative oral therapy

ADVERSE REACTION

-Verry common: Nausea
-Common: Abdominal pain, constipation, dyspepsia, vomitting, Insomia, Dizziness, Oliguria, Hypotension,Sweating increased, pruritits
-Uncommon: Flatulence, dry mouth, Bradycardia, Back pain, Pharyngitis,Rash, Hypertension
-Rare: Myocadial infarction, Erache, GERD



PREGNANCY AND LACTATION

- Category C: Teratogenic effects ; Embryo-fetal development studies conducted with IV parecoxib sodium in rats and rabbits have not shown any evidence of development malformations

CAUTION

RENAL/HEPATIC IMPAIRED

- Hepatic Impairment: No dasgae adjustment
- Renal Impairment: In severe renal impairment (CrCl <30 ml/min) should be initiated at lowest recommended dose and closely monitored

DOSAGE

40 mg administered either IV or IM , followed by 20 mg , or 40 mg every 6-12 hours as required, up to a maximum daily dosage of 80 mg

PRECAUTIONS

- Increase risk of cardiovascular and thombotic adverse events when take long term

