

# Problems 21 to 30

Shiv Shankar Dayal

August 28, 2019

## Problem 21

21. Find the first negative term of the sequence  $2000, 1995, 1990, \dots$

## Solution of problem 21

**Solution:** Clearly,  $a = 2000$ ,  $d = 2000 - 1995 = 1995 - 1900 = 5$

Let  $t_n$  be the first negative term, then we have

$$a + (n - 1)d < 0 \Rightarrow 2000 + (n - 1)5 < 0 \Rightarrow -5n + 2005 < 0$$

$$\Rightarrow n > 401 \therefore n = 402 \text{ least value for which } n > 401$$

Thus,  $402^{nd}$  term will be first negative term.

$$t_{402} = 2000 + (402 - 1)5 = -5$$

## Problem 22

22. How many terms are identical in two arithmetic progressions  $2, 4, 6, 8, \dots$  up to 100 terms and  $3, 6, 9, \dots$  up to 80 terms.

## Solution of problem 22

**Solution:** Let  $r$  terms be identical. Now the sequence of identical terms is 6, 12, 18, ...

$$t_r = a + (r-1)d = 6 + (r-1)6 = 6r$$

$$100^{\text{th}} \text{ term of the sequence } 2, 4, 6, \dots = 2 + (100-1)2 = 200$$

$$80^{\text{th}} \text{ term of the sequence } 3, 6, 9, \dots = 3 + (80-1)3 = 240$$

Thus,  $r^{\text{th}}$  term of the sequence of identical terms cannot be greater than 200

$$6r \leq 200 \Rightarrow r \leq 33\frac{1}{3} \Rightarrow r = 33$$

Hence, 33 terms are identical.

## Problem 23

**23.** Find the number of all positive integers of 3 digits which are divisible by 5.

## Solution of problem 23

**Solution.** Smallest 3 digit number divisible by 5 is 100 and largest is 995.

Clearly, we have  $a = 100$ ,  $d = 5$ ,  $t_n = 995$

$$t_n = 100 + (n - 1)5 \Rightarrow 995 = 95 + 5n \Rightarrow 900 = 5n \Rightarrow n = 180$$

## Problem 24

24. Is 105 a term of the arithmetic progression 4, 9, 14, ...?



## Solution of problem 24

**Solution:**  $a = 4, d = 9 - 4 = 14 - 9 = 5$

Let 105 be  $n^{th}$  term of the arithmetic progression.

$$t_n = a + (n - 1)d \Rightarrow 105 = 4 + (n - 1)5 \Rightarrow 106 = 5n$$

Since  $n$  is not an integer 105 is not a member of the given arithmetic progression.

## Problem 25

25. Find the first negative term of the sequence  $999, 995, 991, \dots$

## Solution of problem 25

**Solution.**  $a = 999, d = 995 - 999 = 991 - 995 = -4$

Let  $n^{th}$  term is first negative number.

$$t_n = a + (n - 1)d = 999 + (n - 1)(-4) = 1003 - 4n < 0 \Rightarrow n > \frac{1003}{4}$$

Least integral value of  $n$  is 251.

$$\therefore t_n = 999 + (251 - 1)(-4) = -1$$

## Problem 26

**26.** Each of the series  $3 + 5 + 7 + \dots$  and  $4 + 7 + 10 + \dots$  is continued to 100 term. Find how many terms are identical?

## Solution of problem 26

**Solution:**  $a = 7$  and  $d = 6$

Last term of first series  $t_{100} = 3 + (100 - 1)2 = 201$

Last term of second series  $t_{100} = 4 + (100 - 1)3 = 301$

Clearly, last term of series of common terms  $t_n < 201$

$$7 + (n - 1)6 < 201 \Rightarrow 6n < 200 \Rightarrow n < 33\frac{1}{3}$$

Least integral value of  $n = 33$

## Problem 27

27. If  $m$  times the  $m^{\text{th}}$  term of an A.P. is equal to  $n$  times the  $n^{\text{th}}$  term, find its  $(m + n)^{\text{th}}$  term.

## Solution of problem 27

**Solution:** Let  $a$  be the first term and  $d$  be the common difference of the A.P.

Given  $nt_n = mt_m$

$$\therefore n[a + (n-1)d] = m[a + (m-1)d]$$

$$(m-n)a = d[n(n-1) - m(m-1)]$$

$$(m-n)a = [n^2 - n - m^2 + m]d$$

$$(m-n)a = -(m-n)(m+n-1)d$$

$$a = -(m+n-1)d$$

$$\text{Now, } t_{m+n} = a + (m+n-1)d = a - a = 0$$

## Problem 28

28. If  $a, b, c$  be the  $p^{th}, q^{th}$  and  $r^{th}$  terms respectively of an A.P., prove that  $a(q - r) + b(r - p) + c(p - q) = 0$



## Solution of problem 28

**Solution:** Let  $x$  be the first term and  $d$  be the common difference of A.P.

$$t_p = a = x + (p - 1)d$$

$$t_q = b = x + (q - 1)d$$

$$t_r = c = x + (r - 1)d$$

$$t_p - t_r = a - c = (p - r)d$$

$$t_q - t_p = b - a = (q - p)d$$

$$t_r - t_q = c - b = (r - q)d$$

$$\text{Now, } a(q - r) + b(r - p) + c(p - q) = q(a - c) + r(b - a) + c(p - q)$$

$$= q(p - r)d + r(q - p)d + p(r - q)d$$

$$= 0$$

## Problem 29

**29.** Find the number of integers between 100 and 1000 that are divisible by 7 and not divisible by 7.

## Solution of problem 29

**Solution:** First number between 100 and 1000 divisible by 7 = 105 and last number = 994

Hence  $a = 105, d = 7, t_n = 994$

$$t_n = 994 = 105 + (n - 1)7$$

$$889 = 7n \therefore n = 127$$

Numbers not divisible by 7 = Total number of numbers between 100 and 1000 - number of numbers between 100 and 1000 divisible by 7  
= 899 - 127 = 772

## Problem 30

30. If  $a, b, c$  be the  $p^{th}, q^{th}$  and  $r^{th}$  terms respectively of an A.P., prove that  $(a - b)r + (b - c)p + (c - a)q = 0$

## Splution of problem 30

**Solution:** Let  $x$  be the first term and  $d$  be the common difference of the A.P.

$$t_p = a = x + (p - 1)d$$

$$t_q = b = x + (q - 1)d$$

$$t_r = c = x + (r - 1)d$$

$$t_p - t_q = a - b = (p - q)d$$

$$t_q - t_r = b - c = (q - r)d$$

$$t_r - t_p = c - a = (r - p)d$$

$$\text{Now, } (a - b)r + (b - c)p + (c - a)q$$

$$= d[(p - q)r + (q - r)p + (r - p)q]$$

$$= 0$$