Arithmetic Progression Problems 71 to 75

Shiv Shankar Dayal

April 10, 2020

71. If the sum of m terms of an A.P.is n and the sum of its n terms is m, show that sum of (m+n) terms is -(m+n)

Solution: Let *a* be the first term and *d* be the common difference of the A.P. Given, $S_m = n$ and $S_n = m$

$$\therefore \frac{m}{2}[2a+(m-1)d] = n \Rightarrow 2a+(m-1)d = \frac{2n}{m}$$

and,

$$\frac{n}{2}[2a+(n-1)d]=m\Rightarrow 2a+(n-1)d=\frac{2m}{n}$$

Subtracting, we get

$$(m-n)d=\frac{2(n^2-m^2)}{mn}\Rightarrow d=-\frac{2(m+n)}{mn}$$

Substituting d in S_m , we get

$$\frac{m}{2}[2a + (m-1)d] = n$$

$$2a = \frac{2n}{m} + (m-1)\frac{2(m+n)}{mn}$$

$$2a = \frac{2n^2 + 2m^2 + 2mn - 2m - 2n}{mn}$$

Thus,

$$S_{m+n} = \frac{m+n}{2}[2a+(m+n-1)d]$$

Substituting for 2a and d, we get



$$S_{m+n} = \frac{m+n}{2} \left[\frac{2n^2 + 2m^2 + 2mn - 2m - 2n}{mn} - (m+n-1) \frac{2(m+n)}{mn} \right]$$

$$S_{m+n} = \frac{m+n}{2} \left[\frac{2n^2 + 2m^2 + 2mn - 2m - 2n - 2m^2 - 2mn - 2mn - 2n^2 + 2m + 2n}{mn} \right]$$

$$S_{m+n} = -(m+n)$$

72. If S be the sum of 2n+1 terms of an A.P., and S_1 that of alternate terms beginning with the first, then show that $\frac{S}{S_1} = \frac{2n+1}{n+1}$

Solution. Let a be the first term and d be the common difference.

$$S = \frac{2n+1}{2}[2a+2nd]$$

$$S_1 = \frac{n+1}{2}[2a+(n+1-1)2d]$$

$$\therefore \frac{S}{S_1} = \frac{2n+1}{n+1}$$

73. If a, b, c be the 1st, 3rd, nth terms respectively of an A.P., prove that the sum of n terms is $\frac{c+a}{2} + \frac{c^2 - a^2}{b-a}$.

Solution: Let *d* be the common difference, then we have

$$t_1 = a, b = a + 2d, c = a + (n-1)d$$

Thus,

$$d = \frac{b-a}{2} \text{ and } (n-1)d = c-a \Rightarrow n = \frac{2(c-a)}{b-a} + 1$$

$$S_n = \frac{n}{2}[2a + (n-1)d]$$

$$= \left(\frac{2(c-a)}{(b-a)} + 1\right)[2a + c - a]$$

$$= \frac{c+a}{2} + \frac{c^2 - a^2}{b-a}$$

73. If the pth term of an A.P. is x and qth term is y. Show that the sum of (p+q) terms is $\frac{p+q}{2}\left(x+y+\frac{x-y}{p-q}\right)$

Solution:Let a be the first term and d be the common difference.

$$t_p = a + (p-1)d = x$$
$$t_q = a + (q-1)d = y$$

Subtracting, we get

$$(p-q)d = x - y \Rightarrow d = \frac{x-y}{p-1}$$

Substituting value of d in t_p , we get

$$a + (p-1)\frac{x-y}{p-q} = x$$

$$a = x - (p-1)\frac{x-y}{p-q}$$

$$S_{p+q} = \frac{p+q}{2}[2a + (p+q-1)d]$$

$$= \frac{p+q}{2}\left[2x - 2(p-1)\frac{x-y}{p-q} + (p+q-1)\frac{x-y}{p-q}\right]$$

$$= \frac{p+q}{2}\left[2x + \frac{x-y}{p-q}(-2p+2+p+q-1)\right]$$

$$= \frac{p+q}{2}\left[2x + \frac{x-y}{p-q}(q-p+1)\right]$$

$$= \frac{p+q}{2}\left[2x - x + y + \frac{x-y}{p-q}\right]$$

$$= \frac{p+q}{2}\left[x - x + y + \frac{x-y}{p-q}\right]$$

4□ > 4回 > 4 亘 > 4 亘 > □ 9 Q @

74. The sum of n terms of two series in A.P. are in ratio (3n+8): (7n+15). Find the ratio of their 12th terms.

Solution: Let a_1, a_2 and d_1, d_2 be first terms and common differences of the two A.P. respectively. Then, we have

$$\frac{\frac{n}{2}[2a_1 + (n-1)d_1]}{\frac{n}{2}[2a_2 + (n-1)d_2]} = \frac{3n+8}{7n+15}$$

$$\frac{2a_1 + (n-1)d_1}{2a_2 + (n-1)d_2} = \frac{3n+8}{7n+15}$$

Substituting n = 23 in the above equation, we get

$$\frac{2a_1 + 22d_1}{2a_2 + 22d_2} = \frac{77}{176}$$

$$\frac{a_1+11d_1}{a_2+11d_2}=\frac{7}{16}$$

Thus, the ratio of 12th terms is $\frac{7}{16}$

75. If the ratio of the sum of m terms and n terms of an A.P. is $m^2 : n^2$, prove that the ratio of its mth and nth term will be (2m-1):(2n-1)

Solution: Let a be the first term and d be the common difference of the A.P. Given,

$$\frac{\frac{m}{2}[2a + (m-1)d]}{\frac{n}{2}[2a + (n-1)d]} = \frac{m^2}{n^2}$$
$$\frac{2a + (m-1)d}{2a + (n-1)d} = \frac{m}{n}$$

Ratio of mth and nth terms is

$$\frac{a + (m-1)d}{a + (n-1)d} = \frac{2a + (2m-1-1)d}{2a + (2n-1-1)d}$$

Substituting m = 2m - 1 and n = 2n - 1 in the equation for ratio of sums, we get

$$\frac{2a + (2m - 1 - 1)d}{2a + (2n - 1 - 1)d} = \frac{2m - 1}{2n - 1}$$

Thus, ratio of mth and nth term is (2m-1): (2n-1)