


Trigonometry with Geometry

A problem-oriented approach



Build Date: March 12, 2023

Copyright, © Shiv Shankar Dayal, 2011-2023. All rights reserved.

Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.3 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled “GNU Free Documentation License”.

*Dedicated to my family
and Free Software Community*

When you take an action think of the poorest person you have seen and evaluate how your action will benefit that person.

स्वध्याय परमो तपः (Self study is ultimate penance.)

CONTENTS

Contents	i
Preface	iii
I Theory and Problems	1
1 Measurement of Angles	2
1.1 Angles in Geometry	2
1.2 Angles in Trigonometry	3
1.3 Angles Exceeding 360°	3
1.4 Quadrants	4
1.5 Units of Measurement	5
1.6 Problems	5
2 Trigonometric Ratios	11
2.1 Relationship between Trigonometric Functions or Ratios	12
2.2 Problems	13
3 Trigonometric Ratios of Any Angle and Sign	18
3.1 Angle of 45°	18
3.2 Angles of 30° and 60°	19
3.3 Angle of 0°	19
3.4 Angle of 90°	20
3.5 Complementary Angles	20
3.6 Supplementary Angles	21
3.7 Angles of $-\theta$	22
3.8 Angles of $90^\circ + \theta$	22
3.9 Angles of $360^\circ + \theta$	23
3.10 Problems	23
4 Compound Angles	27
4.1 The Addition Formula	27
4.2 The Subtraction Formula	28
4.3 Important Deductions	29
4.4 To express $a \cos \theta + b \sin \theta$ in the form of $k \cos \phi$ or $k \sin \phi$	30
4.5 Problems	31

5	Transformation Formulae	35
5.1	Transformation of products into sums or differences	35
5.2	Transformation of sums or differences into products	35
5.3	Problems	36
6	Multiple and Submultiple Angles	41
6.1	Multiple Angles	41
II	Solutions	42
	GNU Free Documentation License	43

PREFACE

This is a book on trigonometry, which, covers basics of trigonometry till high school level. It covers the most essential topics to take up a bachelor's course where knowledge of trigonometry is required. There is no specific purpose for writing this book. This is a book for self study and is not recommended for courses in schools and universities. I will try to cover as much as I can and will keep adding new material over a long period. I have no interest in writing a book in a fixed way which serves a university or college course as I have always loved freedom. Life, freedom and honor in that order are important. That was one of the reasons why I never did a masters course. Enough of personal rant. Let us get on with the actual work.

Trigonometry is probably one of the most fundamental subjects in Mathematics as further study of subjects like coordinate geometry, 3D and 2D geometry, engineering and rest all depend on it. It is very important to understand trigonometry for the readers if they want to advance further in mathematics.

Why I Wrote This Book?

I wrote this book for myself! I did not write it for anybody else. My knowledge, which I have acquired by reading books written by many great mathematicians and authors and interactions with many intelligent people, is what has been put in the book. I have just tried to add my flavor to it. Think of it as notes for me. Just that I like to organize my notes so it has taken form of a book and nothing more. If you benefit from this then that is a pure coincidence and not intentional at all.

How to Read This Book?

No I will not simply tell that you must solve the problems. My advice would be more detailed. Every chapter will have theory. Read that first. Make sure you understand that. Of course, you have to meet the prerequisites for the book. Then, go on and try to solve the problems. In this book, there are no pure problems. Almost all have answers except those which are of similar kind and repetitive in nature for the sake of practice. If you can solve the problem then all good else look at the answer and try to understand that.

Then, few days later take on the problem again. If you fail to understand the answer you can always email me with your work and I will try to answer to the best of my ability. However, if you have a local expert seek his/her advice first. Just that email is bad for mathematics.

Note that mathematics is not only about solving problems. If you understand the theory well, then you will be able to solve problems easily. However, problems do help enforce with the enforcement of theory in your mind.

I am a big fan of old MIR publisher's problem books, so I emphasize less on theory and more on problems. I hope that you find this style much more fun as a lot of theory is boring. Mathematics is about problem solving as that is the only way to enforce theory and find innovative techniques for problem solving.

Some of the problems in certain chapters rely on other chapters which you should look ahead or you can skip those problems and come back to it later. Since this book is meant for self study answers of most of the problems have been given which you can make use of. However, do not use for just copying but rather to develop understanding.

Who Should Read This Book?

Since this book is written for self study anyone with interest in trigonometry can read it. That does not mean that school or college students cannot read it. You need to be selective as to what you need for your particular requirements. This is mostly high school course with a little bit of lower classes' course thrown in with a bit of detail here and there.

Prerequisite

You should have knowledge till grade 10th course. Attempt has been made to keep it simple and give as much as background to the topic which is reasonable and required. However, not everything will be covered below grade 10.

Goals for Readers

The goal of for reading this book is becoming proficient in solving simple and basic problems of trigonometry. Another goal would be to be able to study other subjects which require this knowledge like trigonometry or calculus or physics or chemistry or other subjects. If you can solve 95% problems after 2 years of reading this book then you have achieved this goal.

All of us possess a certain level of intelligence. At average any person can read this book. But what is most important is you have to have interest in the subject. Your interest gets multiplied with your intelligence and thus you will be more capable than you think you

can be. One more point is focus and effort. It is not something new which I am telling but I am saying it again just to emphasize the point. Trust me if you are reading this book for just scoring a nice grade in your course then I have failed in my purpose of explaining my ideas.

Also, if you find this book useful feel free to share it with others without hesitation as it is free as in freedom. There are no conditions to share it.

Acknowledgements

I am in great debt of my family and free software community because both of these groups have been integral part of my life. Family has provided direct support while free software community has provided the freedom and freed me from the slavery which comes as a package with commercial software. I am especially grateful to my wife, son and parents because it is their time which I have borrowed to put in the book. To pay my thanks from free software community I will take one name and that is Richard Stallman who started all this and is still fighting this never-ending war. When I was doing the algebra book then I realized how difficult it is to put Math on web in HTML format and why Donald Knuth wrote T_EX. Also, T_EX was one of the first softwares to be released as a free software.

Now as this book is being written using L^AT_EX so obviously Leslie Lamport and all the people involved with it have my thanks along with Donald Knuth. I use Emacs with Auctex and hope that someday I will use it in a much more productive way someday.

I have used Asymptote and tikz for drawing all the diagrams. Both are wonderful packages and work very nicely. Asymptote in particular is very nice for 3d-drawings and linear equation solving.

I would like to thank my parents, wife and son for taking out their fair share of time and the support which they have extended to me during my bad times. After that I would like to pay my most sincere gratitude to my teachers particularly H. N. Singh, Yogendra Yadav, Satyanand Satyarthi, Kumar Shailesh and Prof. T. K. Basu. Now is the turn of people from software community. I must thank the entire free software community for all the resources they have developed to make computing better. However, few names I know and here they go. Richard Stallman is the first, Donald Knuth, Edger Dijkstra, John von Neumann after that as their lives have strong influence in how I think and base my life on. Cover graphics has been done by Koustav Halder so much thanks to him.

I am not a native English speaker and this book has just gone through one pair of eyes therefore chances are high that it will have lots of errors (particularly with commas and spelling mistakes). At the same time it may contain lots of technical errors. Please feel free to drop me an email at shivshankar.dayal@gmail.com where I will try to respond to each mail as much as possible. Please use your real names in email not something like coolguy. If you have more problems which you want to add it to the book please send those by email or create a PR on github. The github url is <https://github.com/shivshankardayal/Trigonometry-Latex>.

Shiv Shankar Dayal

Part I


Theory and Problems

MEASUREMENT OF ANGLES

The word trigonometry comes from means measurement of triangles. The word originally comes from Greek language. measurement. The objective of studying plane trigonometry is to develop a method of solving plane triangles. However, as time changes everything it has changed the scope of trigonometry to include polygons and circles as well. A lot of concepts in this book will come from your geometry classes in lower classes. It is a good idea to review the concepts which you have studied till now without which you are going to struggle while studying trigonometry in this book.


1.1 Angles in Geometry

If we consider a line extending to infinity in both directions, and a point O which divides this line in two parts one on each side of the point then each part is called a ray or half-line. Thus O divides the line into two rays OA and OA' .



The point O is called vertex or origin for these rays. An angle is a figure formed by two rays or half lines meeting at a common vertex. These half lines are called *sides of the angle*.

An angle is denoted by the symbol \angle followed by three capital letters of which the middle one represents the vertex and remaining two points point to two sides. Otherwise the angle is simply written as one letter representing the vertex of the angle.




The angle in above image is written as $\angle AOB$ or $\angle BOA$ or $\angle O$.

Each angle can be measured and there are different units for the measurement. In Geometry, an angle always lie between 0° and 360° and negative angles are meaningless. Measure of an angle is the smallest amount of rotation from the direction of one ray of the angle to the direction of the other.

1.2 Angles in Trigonometry

Angles are more generalized in Trigonometry. They can have positive or negative values. As was the case in geometry, similarly angles are measured in Trigonometry. The starting and ending positions of revolving rays are called initial side and terminal side respectively. The revolving half line is called the generating line or the radius vector. For example, if OA and OB are the initial and final position of the radius vector then angle formed will be $\angle AOB$.

1.3 Angles Exceeding 360°




In Geometry, angles are limited to 0° to 360° . However, when multiple revolutions are involved angles are more than 360° . For example, the revolving line starts from the initial

position and makes n complete revolutions in anticlockwise direction and also further angle α in the same direction. We then have a certain angle β_n given by $\beta_n = n \times 360^\circ + \alpha$, where $0^\circ < \alpha < 360^\circ$ and n is zero or positive integer. Thus, there are infinite possible angles.

Angles formed by anticlockwise rotation of the radius vector are taken as positive and angles formed by clockwise rotation of the radius vector are taken as negative.

1.4 Quadrants



Let XOX' and YOY' be two mutually perpendicular lines in a plane and OX be the initial half line. The lines divide the whole plane into four quadrants. XOY , YOX' , $X'OY'$ and $Y'OX$ are respectively called 1st, 2nd, 3rd and 4th quadrants. According to terminal side lying in 1st, 2nd, 3rd and 4th quadrants the angles are said to be in 1st, 2nd, 3rd and 4th quadrants respectively. A *quadrant angle* is an angle formed if terminal side coincides with one of the axes.

For any angle \angle which is not a quadrant angle and when number of revolutions is zero and radius vector rotates in anticlockwise directions:

1. $0^\circ < \alpha < 90^\circ$ if α lies in first quadrant
2. $90^\circ < \alpha < 180^\circ$ if α lies in second quadrant
3. $180^\circ < \alpha < 270^\circ$ if α lies in third quadrant
4. $270^\circ < \alpha < 360^\circ$ if α lies in fourth quadrant
5. when terminal side lies on OY , angle formed = 90°
6. when terminal side lies on OX' , angle formed = 180°
7. when terminal side lies on OY' , angle formed = 270°
8. when terminal side lies on OX , angle formed = 360°

1.5 Units of Measurement

In Geometry, angles are usually measured in terms of right angles, however, that is an inconvenient system for smaller angles. So we introduce different systems of measurements. There are three system of units for this:

1. Sexagesimal or British system: In British system, a right angle is divided into 90 equal parts called degrees. Each degree is then divided into 60 equal parts called minutes and each minute is further is divided into 60 parts called seconds.
A degree, a minute and a second are denoted by 1° , $1''$, and 1 respectively.
2. Centesimal or French System: In French system, a right angle is divided into 100 equal parts called grades. Each grade is then divided into 100 equal parts called minutes and each minute is further is divided into 100 parts called seconds.
A degree, a grade and a second are denoted by 1^g , $1''$, and 1 respectively.
3. Radian or Circular Measure: An arc equal to radius of a circle when subtends an anngle on the center then that angle is 1 radian and is denoted by 1^c . The angle made by half of perimeter is π radians. Also, from Geometry we know that angle subtended is the ratio between length of cord and radius. This ratio is in radians. Since both length or chord and radius have same unit radian is a constant.

Relationship between Systems of Measurements

If measure of an angle if D degrees, G grades and C radians then upon elementary manipulation we find that $\frac{D}{180} = \frac{G}{200} = \frac{C}{\pi}$.

Meaning of π

The ratio of circumference and diameter of a circle is always constant and this constant is denoted by gree letter π .

π is an irrational number. In general, we use the value of $\frac{22}{7}$ but $\frac{355}{113}$ is more accurate though not exact. If r be the radius of a circle and c be the circumference then $\frac{c}{2r} = \pi$ leading circumference to be $c = 2\pi r$.

1.6 Problems

1. Reduce $63^\circ 14' 51''$ to centesimal measure.
2. Reduce $45^\circ 20' 10''$ to centesimal and radian measure.

3. Reduce $94^g23'27''$ to Sexagecimal measure.
4. Reduce 1.2 radians in Sexageciaml measure.

Express in terms of right angle; the angles

5. 60°
6. $75^\circ15'$
7. $63^\circ17'25''$
8. $130^\circ30'$
9. $210^\circ30'30''$
10. $370^\circ20'48''$

Express in grades, minutes and degrees

11. 30°
12. 81°
13. $138^\circ30'$
14. $35^\circ47'15''$
15. $235^\circ12'36''s$
16. $475^\circ13'48''$

Express in terms of right angles and also in degrees, minutes and seconds; the angles

17. 120^g
18. $45^g35'24''$
19. $39^g45'36''$
20. $255^g8'9''$
21. $759^g0'5''$
22. Reduce $55^\circ12'36''$ to centesimal measure.
23. Reduce $18^\circ33'45''$ to circular measure.
24. Reduce $196^g35'24''$ to sexagecimal measure.
25. How many degrees, minutes and seconds are respectively passed over in $11\frac{1}{9}$ minutes by the hour and minute hand of a watch.

26. The number of degrees in one acute angle of a right-angled triangle is equal to the number of grades in the other; express both angles in degrees.
27. Prove that the number of Sexagesimal minutes in any angle is to the number of Centesimal minutes in the same angle as 27 : 50.
28. Divide $44^{\circ}8'$ into two parts such that the number of Sexagesimal seconds in one part may be equal to number of Centesimal seconds in the other part.
29. The angles of a triangle are in the ratio of 3 : 4 : 5, find the smallest angle in degrees and greatest angle in radians.
30. Find the angle between the hour hand and the minute hand in circular measure at half past four.
31. If p, q and r denote the grade measure, degree measure and the radian measure of the same angle, prove that
- a) $\frac{p}{10} = \frac{q}{9} = \frac{20r}{\pi}$
- b) $p - q = \frac{20r}{\pi}$
32. Two angles of a triangle are $72^{\circ}53'51''$ and $41^{\circ}22'50''$ respectively. Find the third angle in radians.
33. The angles of triangle are in A.P. and the number of radians in the greatest angle is to the number of degrees in the least one as $\pi : 60$; find the angles in degrees.
34. The angles of a triangle are in A.P. and the number of grades in the least is to the number of radians in the greatest is $40 : \pi$; find the angles in degrees.
35. Three angles are in G.P. The number of grades in the greatest angle is to the number of circular units in the least is $800 : \pi$; and the sum of angles is 126° . Find the angles in grades.
36. Find the angle between the hour-hand and minute-hand in circular measure at 4 o'clock.
37. Express in sexagesimal system the angle between the minute-hand and hour-hand of a clock at quarter to twelve.
38. The diameter of a wheel is 28 cm; through what distance does its center move during one rotation of wheel along the ground?
39. What must be the radius of a circular running path, round which an athlete must run 5 time in order to describe 1760 meters?
40. The wheel of a railway carriage is 90 cm in diameter and it makes 3 revolutions per second; how fast is the train going?
41. A mill sail whose length is 540 cm makes 10 revolutions per minute. What distance does its end travel in one hour?
42. Assuming that the earth describes in one year a circle, or 149,700,000 km. radius, whose center is the sun, how many miles does earth travel in a year?


43. The radius of a carriage wheel is 50 cm, and in $\frac{1}{9}$ th of a second it turns through 80° about its center, which is fixed; how many km. does a point on the rim of the wheel travel in one hour?
44. Express in terms of three systems of angular measurements the magnitude of an angle of a regular decagon.
45. One angle of a triangle is $\frac{2}{3}x$ grades and another is $\frac{3}{2}x$ degrees, while the third is $\frac{\pi x}{75}$ radians; express them all in degrees.
46. The circular measure of two angles of a triangle are $\frac{1}{2}$ and $\frac{1}{3}$. What is the number of degrees of the third angle?
47. The angles of a triangle are in A.P. The number of radians in the least angle is to the number of degree in the mean angle is 1 : 120. Find the angles in radians.
48. Find the magnitude, in radians and degrees, of the interior angle of 1. a regular pentagon 2. a regular heptagon 3. a regular octagon 4. a regular duodecagon 5. a polygon with 17 sides
49. The angle in one regular polygon is to that in another is 3 : 2, also the number of sides in the first is twice that in the second. How many sides are there in the polygons?
50. The number of sides in two regular polygons are as 5 : 4, and the difference between their angles is 9° ; find the number of sides in the polygons.
51. Find two regular polygons such that the number of their sides may be 3 to 4 and the number of degrees of an angle of the first to the number of grades of the second as 4 to 5.
52. The angles of a quadrilateral are in A.P. and the greatest is double the least; express the least angle in radians.
53. Find in radians, degrees, and grades the angle between hour-hand and minute-hand of a clock at 1. half-past three 2. twenty minutes to six 3. a quarter past eleven.
54. Find the times 1. between four and five o'clock when the angle between the minute hand and the hour-hand is 78° , 2. between seven and eight o'clock when the angle is 54°
55. The interior angles of a polygon are in A.P. The smallest angle is 120° and the common difference is 5° . Find the number of sides of the polygon.
56. The angles of quadrilateral are in A.P. and the number of grades in the least angle is to the number of radians in the greatest is $100 : \pi$. Find the angles in degrees.
57. The angles of a polygons are in A.P. The least angle is $\frac{5\pi}{12}$ common difference is 10° , find the number of sides in the polygon.
58. Find the angle subtended at the center of a circle of radius 3 cm. by an arc of length 1 cm.

59. In a circle of radius 5 cm., what is the length of the arc which subtends an angle of $33^{\circ}15'$ at the center.
60. Assuming the average distance of sun from the earth to be 149,700,000 km., and the angle subtended by the sun at the eye of a person on the earth is $32'$, find the sun's diameter.
61. Assuming that a person of normal sight can read print at such a distance that the letter subtends an angle of $5'$ at his eye, find what is the height of the letters he can read at a distance of 1. 12 meters 2. 1320 meters.
62. Find the number of degrees subtended at the center of a circle by an arc whose length is 0.357 times the radius.
63. Express in radians and degrees the angle subtended at the center of a circle by an arc whose length is 15 cm., the radius of the circle being 25 cm.
64. The value of the divisions on the outer rim of a graduated circle is $5'$ and the distance between successive graduations is .1 cm. Find the radius of the circle.
65. The diameter of a graduated circle is 72 cm., and the graduations on the rim are $5'$ apart; find the distance of one graduation to another.
66. Find the radius of a globe which is such that the distance between two places on the same meridian whose latitude differs by $1^{\circ}10'$ may be 0.5 cm.
67. Taking the radius of earth to be 6400 km., find the difference in latitude of two places, one of which is 100 km. north of another.
68. Assuming the earth to be a sphere and the difference between two parallels of latitude, which subtends an angle of 1° at the earth's center, to be $69\frac{1}{2}$ km., find the radius of the earth.
69. What is the ratio of radii of the circles at the center of which two arcs of same length subtend angles of 60° and 75° ?
70. If an arc, of length 10 cm., on a circle of 8 cm. diameter subtend at the center of circle an angle of $143^{\circ}14'22''$, find the value of π to 4 places of decimals.
71. If the circumference of a circle be divided into five parts which are in A.P., and if the greatest part be six times the least find in radians the magnitude of the angles the parts subtend at the center of the circle.
72. The perimeter of a certain sector of a circle is equal to the length of the arc of a semicircle having the same radius; express the angle of the sector in degrees.
73. At what distance a man, whose height is 2 m., subtend an angle of $10'$.
74. Find the length which at a distance of 5280 m., will subtend an angle of $1'$ at the eye.
75. Assuming the distance of the earth from the moon to be 38400 km., and the angle subtended by the moon at the eye of a person on earth to be $31'$, find the diameter of the moon.

-
76. The wheel of a railway carriage is 4 ft. in diameter and makes 6 revolutions in a second; how fast is the train going?
77. Assuming that moon subtends an angle of $30'$ at the eye of an observer, find how far from the eye a coin of one inch diameter must be held so as just to hide the moon.
78. A wheel make 30 revolutions per minute. Find the circular measure of the angle described by spoke in half a second.
79. A man running along a circular track at the rate of 10 miles per hour, traverses in 36 seconds, an arc which subtends an angle of 56° at the center. Find the diameter of the circle.

TRIGONOMETRIC RATIOS

From Geometry, we know that an acute angle is an angle whose measure is between 0° and 90° . Consider the following figure:



This picture contains two similar triangles $\triangle OMP$ and $\triangle OM'P'$. We are interested in $\angle MOP$ or $\angle M'OP'$. In the $\triangle MOP$ and $\triangle M'OP'$, OP, OP' are called the hypotenuses i.e. sides opposite to the right angle, $PM, P'M'$ are called perpendiculars i.e. sides opposite to the angle of interest and OM, OM' are called bases i.e. the third angle.

Hypotenuses are usually denoted by h , perpendiculars by p and bases by b . Let $OM = b, OM' = b', PM = p, P'M' = p', OP = h, OP' = h'$. Since the two triangles are similar $\therefore \frac{p}{p'} = \frac{b}{b'} = \frac{h}{h'}$. Thus the ratio of any two sides is dependent purely on $\angle O$ or $\angle MOP$ or $\angle M'OP'$.

Since there are three sides, we can choose 2 in 3C_2 i.e. 3 ways and for each combination there will be two permutations where a side can be in either numerator or denominator.

From this we can conclude that there will be six ratios (these are called trigonometric ratios). These six trigonometric ratios or functions are given below:

$\frac{MP}{OP}$ or $\frac{p}{h}$ is called the **Sine** of the $\angle MOP$.

$\frac{OM}{OP}$ or $\frac{b}{h}$ is called the **Cosine** of the $\angle MOP$.

$\frac{MP}{OM}$ or $\frac{p}{b}$ is called the **Tangent** of the $\angle MOP$.

$\frac{OP}{MP}$ or $\frac{h}{p}$ is called the **Cosecant** of the $\angle MOP$.

$\frac{OP}{OM}$ or $\frac{h}{b}$ is called the **Secant** of the $\angle MOP$.

$\frac{OM}{MP}$ or $\frac{b}{p}$ is called the **Cotangent** of the $\angle MOP$.

$1 - \cos MOP$ is called the **Versed Sine** of $\angle MOP$ and $1 - \sin MOP$ is called the **Covered Sine** of $\angle MOP$. These two are rarely used in trigonometry. It should be noted that the trigonometric ratios are all numbers. The name of the trigonometric ratios are written for brevity $\sin MOP$, $\cos MOP$, $\tan MOP$, $\cot MOP$, $\sec MOP$, $\csc MOP$, $\text{vers } MP$, $\text{coverse } MOP$.

2.1 Relationship between Trigonometric Functions or Ratios

Let us represent the $\angle MOP$ with θ , we observe from previous section that

$$\sin \theta = \frac{1}{\csc \theta}, \cos \theta = \frac{1}{\sec \theta}, \tan \theta = \frac{1}{\cot \theta}, \csc \theta = \frac{1}{\sin \theta}, \sec \theta = \frac{1}{\cos \theta}, \cot \theta = \frac{1}{\tan \theta}$$

$$\text{We also observe that } \tan \theta = \frac{\sin \theta}{\cos \theta} \text{ and } \cot \theta = \frac{\cos \theta}{\sin \theta}$$

From Pythagora theorem in geometry, we know that $\text{hypotenuse}^2 = \text{perpendicular}^2 + \text{base}^2$ or $h^2 = p^2 + b^2$

1. Dividing both side by h^2 , we get

$$\frac{p^2}{h^2} + \frac{b^2}{h^2} = 1$$

$$\sin^2 \theta + \cos^2 \theta = 1$$

We can rewrite this as $\sin^2 \theta = 1 - \cos^2 \theta$, $\cos^2 \theta = 1 - \sin^2 \theta$, $\sin \theta = \sqrt{1 - \cos^2 \theta}$, $\cos \theta = \sqrt{1 - \sin^2 \theta}$.

2. If we divide both sides by b^2 , then we get

$$\frac{h^2}{b^2} = \frac{p^2}{b^2} + 1$$

$$\sec^2 \theta = \tan^2 \theta + 1$$

We can rewrite this as $\sec^2 \theta - \tan^2 \theta = 1$, $\tan^2 \theta = \sec^2 \theta - 1$, $\sec \theta = \sqrt{1 + \tan^2 \theta}$, $\tan \theta = \sqrt{\sec^2 \theta - 1}$

3. Similarly, if we divide by p^2 , then we get

$$\frac{h^2}{p^2} = 1 + \frac{b^2}{p^2}$$

$$\operatorname{cosec}^2 \theta = 1 + \cot^2 \theta$$

We can rewrite this as $\operatorname{cosec}^2 \theta - \cot^2 \theta = 1$, $\cot^2 \theta = \operatorname{cosec}^2 \theta - 1$, $\operatorname{cosec} \theta = \sqrt{1 + \cot^2 \theta}$, $\cot \theta = \sqrt{\operatorname{cosec}^2 \theta - 1}$

2.2 Problems

Prove the following:

1. $\sqrt{\frac{1 - \cos A}{1 + \cos A}} = \operatorname{cosec} A - \cot A.$
2. $\sqrt{\sec^2 A + \operatorname{cosec}^2 A} = \tan A + \cot A.$
3. $(\operatorname{cosec} A - \sin A)(\sec A - \cos A)(\tan A + \cot A) = 1.$
4. $\cos^4 A - \sin^4 A + 1 = 2 \cos^2 A.$
5. $(\sin A + \cos A)(1 - \sin A \cos A) = \sin^3 A + \cos^3 A.$
6. $\frac{\sin A}{1 + \cos A} + \frac{1 + \cos A}{\sin A} = 2 \operatorname{cosec} A.$
7. $\sin^6 A - \cos^6 A = 1 - 3 \cos^2 A \sin^2 A.$
8. $\sqrt{\frac{1 - \sin A}{1 + \sin A}} = \sec A - \tan A.$
9. $\frac{\operatorname{cosec} A}{\operatorname{cosec} A - 1} + \frac{\operatorname{cosec} A}{\operatorname{cosec} A + 1} = 2 \sec^2 A.$
10. $\frac{\operatorname{cosec} A}{\tan A + \cot A} = \cos A.$
11. $(\sec A + \cos A)(\sec A - \cos A) = \tan^2 A + \sin^2 A.$

12. $\frac{1}{\tan A + \cot A} = \sin A \cos A.$
13. $\frac{1 - \tan A}{1 + \tan A} = \frac{\cot A - 1}{\cot A + 1}.$
14. $\frac{1 + \tan^2 A}{1 + \cot^2 A} = \frac{\sin^2 A}{\cos^2 A}.$
15. $\frac{\sec A - \tan A}{\sec A + \tan A} = 1 - 2 \sec A \tan A + 2 \tan^2 A.$
16. $\frac{1}{\sec A - \tan A} = \sec A + \tan A.$
17. $\frac{\tan A}{1 - \cot A} + \frac{\cot A}{1 - \tan A} = \sec A \operatorname{cosec} A + 1.$
18. $\frac{\cos A}{1 - \tan A} + \frac{\sin A}{1 - \cot A} = \sin A + \cos A.$
19. $(\sin A + \cos A)(\tan A + \cot A) = \sec A + \operatorname{cosec} A.$
20. $\sec^4 A - \sec^2 A = \tan^4 A + \tan^2 A.$
21. $\cot^4 A + \cot^2 A = \operatorname{cosec}^4 A - \operatorname{cosec}^2 A.$
22. $\sqrt{\operatorname{cosec}^2 A - 1} = \cos A \operatorname{cosec} A.$
23. $\sec^2 A \operatorname{cosec}^2 A = \tan^2 A + \cot^2 A + 2.$
24. $\tan^2 A - \sin^2 A = \sin^4 A \sec^2 A.$
25. $(1 + \cot A - \operatorname{cosec} A)(1 + \tan A + \sec A) = 2.$
26. $\frac{\cot A \cos A}{\cot A + \cos A} = \frac{\cot A - \cos A}{\cot A \cos A}.$
27. $\frac{\cot A + \tan B}{\cot B + \tan A} = \cot A \tan B.$
28. $\left(\frac{1}{\sec^2 A - \cos^2 A} + \frac{1}{\operatorname{cosec}^2 A - \sin^2 A} \right) \cos^2 A \sin^2 A = \frac{1 - \cos^2 A \sin^2 A}{2 + \cos^2 A \sin^2 A}.$
29. $\sin^8 A - \cos^8 A = (\sin^2 A - \cos^2 A)(1 - 2 \sin^2 A \cos^2 A).$
30. $\frac{\cos A \operatorname{cosec} A - \sin A \sec A}{\cos A + \sin A} = \operatorname{cosec} A - \sec A.$
31. $\frac{1}{\operatorname{cosec} A - \cot A} - \frac{1}{\sin A} = \frac{1}{\sin A} - \frac{1}{\operatorname{cosec} A + \cot A}.$
32. $\frac{\tan A + \sec A - 1}{\tan A - \sec A + 1} = \frac{1 + \sin A}{\cos A}.$
33. $(\tan A + \operatorname{cosec} B)^2 - (\cot B - \sec A)^2 = 2 \tan A \cot B (\operatorname{cosec} A + \sec B).$
34. $2 \sec^2 A - \sec^4 A - 2 \operatorname{cosec}^2 A + \operatorname{cosec}^4 A = \cot^4 A - \tan^4 A.$


35. $(\sin A + \operatorname{cosec} A)^2 + (\cos A + \sec A)^2 = \tan^2 A + \cot^2 A + 7$.
36. $(\operatorname{cosec} A + \cot A)(1 - \sin A) - (\sec A + \tan A)(1 - \cos A) = (\operatorname{cosec} A - \sec A)[2 - (1 - \cos A)(1 - \sin A)]$.
37. $(1 + \cot A + \tan A)(\sin A - \cos A) = \frac{\sec A}{\operatorname{cosec}^2 A} - \frac{\operatorname{cosec} A}{\sec^2 A}$.
38. $\frac{1}{\sec A - \tan A} - \frac{1}{\cos A} = \frac{1}{\cos A} - \frac{1}{\sec A + \tan A}$.
39. $3(\sin A - \cos A)^4 + 4(\sin^6 A + \cos^6 A) + 6(\sin A + \cos A)^2 = 13$.
40. $\sqrt{\frac{1 + \cos A}{1 - \cos A}} = \operatorname{cosec} A + \cot A$.
41. $\frac{\cos A}{1 + \sin A} + \frac{\cos A}{1 - \sin A} = 2 \sec A$.
42. $\frac{\tan A}{\sec A - 1} + \frac{\tan A}{\sec A + 1} = 2 \operatorname{cosec} A$.
43. $\frac{1}{1 - \sin A} - \frac{1}{1 + \sin A} = 2 \sec A \tan A$.
44. $\frac{1 + \tan^2 A}{1 + \cot^2 A} = \left(\frac{1 - \tan A}{1 - \cot A} \right)^2$.
45. $1 + \frac{2 \tan^2 A}{\cos^2 A} = \tan^4 A + \sec^4 A$.
46. $(1 - \sin A - \cos A)^2 = 2(1 - \sin A)(1 - \cos A)$.
47. $\frac{\cot A + \operatorname{cosec} A - 1}{\cot A - \operatorname{cosec} A + 1} = \frac{1 + \cos A}{\sin A}$.
48. $(\sin A + \sec A)^2 + (\cos A + \operatorname{cosec} A)^2 = (1 + \sec A \operatorname{cosec} A)^2$.
49. $\frac{2 \sin A \tan A(1 - \tan A) + 2 \sin A \sec^2 A}{(1 + \tan A)^2} = \frac{2 \sin A}{1 + \tan A}$.
50. If $2 \sin A = 2 - \cos A$, find $\sin A$.
51. If $8 \sin A = 4 + \cos A$, find $\sin A$.
52. If $\tan A + \sec A = 1.5$, find $\sin A$.
53. If $\cot A + \operatorname{cosec} A = 5$, find $\cos A$.
54. If $3 \sec^4 A + 8 = 10 \sec^2 A$, find the value of $\tan A$.
55. If $\tan^2 A + \sec A = 5$, find $\cos A$.
56. If $\tan A + \cot A = 2$, find $\sin A$.
57. If $\sec^2 A = 2 + 2 \tan A$, find $\tan A$.

58. If $\tan A = \frac{2x(x+1)}{2x+1}$, find $\sin A$ and $\cos A$.
59. If $3 \sin A + 5 \cos A = 5$, show that $5 \sin A - 3 \cos A = \pm 3$.
60. If $\sec A + \tan A = \sec A - \tan A$ prove that each side is ± 1 .
61. If $\frac{\cos^4 A}{\cos^2 B} + \frac{\sin^4 A}{\sin^2 B} = 1$, prove that
- $\sin^4 A + \sin^4 B = 2 \sin^2 A \sin^2 B$,
 - $\frac{\cos^4 B}{\cos^2 A} + \frac{\sin^4 B}{\sin^2 A} = 1$.
62. If $\cos A + \sin A = \sqrt{2} \cos A$, prove that $\cos A - \sin A = \pm \sqrt{2} \sin A$.
63. If $a \cos A - b \sin A = c$, prove that $a \sin A + b \cos A = \sqrt{a^2 + b^2 - c^2}$.
64. If $1 - \sin A = 1 + \sin A$, then prove that value of each side is $\pm \cos A$.
65. If $\sin^4 A + \sin^2 A = 1$, prove that
- $\frac{1}{\tan^4 A} + \frac{1}{\tan^2 A} = 1$,
 - $\tan^4 A - \tan^2 A = 1$.
66. If $\cos^2 A - \sin^2 A = \tan^2 B$, prove that $2 \cos^2 B - 1 = \cos^2 B - \sin^2 B = \tan^2 A$.
67. If $\sin A + \operatorname{cosec} A = 2$, then prove that $\sin^n A + \operatorname{cosec}^n A = 2$.
68. If $\tan^2 A = 1 - e^2$, prove that $\sec A + \tan^3 A \operatorname{cosec} A = (2 - e^2) \frac{3}{2}$.
69. Eliminate A between the equations $a \sec A + b \tan A + c = 0$ and $p \sec A + q \tan A + r = 0$.
70. If $\operatorname{cosec} A - \sin A = m$ and $\sec A - \cos A = n$, eliminate A .
71. Is the equation $\sec^2 A = \frac{4xy}{(x+y)^2}$ possible for real values of x and y ?
72. Show that the equation $\sin A = x + \frac{1}{x}$ is impossible for real values of x .
73. If $\sec A - \tan A = p, p \neq 0$, find $\tan A, \sec A$ and $\sin A$.
74. If $\sec A = p + \frac{1}{4p}$, show that $\sec A + \tan A = 2p$ or $\frac{1}{2p}$.
75. If $\frac{\sin A}{\sin B} = p, \frac{\cos A}{\cos B} = q$, find $\tan A$ and $\tan B$.
76. If $\frac{\sin A}{\sin B} = \sqrt{2}, \frac{\tan A}{\tan B} = \sqrt{3}$, find A and B .
77. If $\tan A + \cot A = 2$, find $\sin A$.
78. If $m = \tan A + \sin A$ and $n = \tan A - \sin A$, prove that $m^2 - n^2 = 4\sqrt{mn}$.

79. If $\sin A + \cos A = m$ and $\sec A + \operatorname{cosec} A = n$, prove that $n(m^2 - 1) = 2m$.
80. If $x \sin^3 A + y \cos^3 A = \sin A \cos A$ and $x \sin A - y \cos A = 0$, prove that $x^2 + y^2 = 1$.
81. Prove that $\sin^2 A = \frac{(x+y)^2}{4xy}$ is possible for real values of x and y only when $x = y$ and $x, y \neq 0$.

TRIGONOMETRIC RATIOS OF ANY ANGLE AND SIGN

3.1 Angle of 45°



Consider the above figure, which is a right-angle triangle, drawn so that $\angle OMP = 90^\circ$ and $\angle MOP = 45^\circ$. We know that the sum of all angles of a triangle is 180° . Thus,

$$\angle OPM = 180^\circ - \angle MOP - \angle OMP = 180^\circ - 90^\circ - 45^\circ = 45^\circ$$


$\therefore OM = MP$. Let $OP = 2a$, then from Pythagora theorem, we can write

$$4a^2 = OP^2 = OM^2 + MP^2 = 2OM^2 \Rightarrow Om = a\sqrt{2} = MP$$

$$\sin 45^\circ = \frac{MP}{OP} = \frac{a\sqrt{2}}{2a} = \frac{1}{\sqrt{2}}.$$

Other trigonometric ratios can be deduced similarly for this angle.

3.2 Angles of 30° and 60°



Consider an equilateral $\triangle OMP$. Let the sides OM, OP, MP be each $2a$. We draw a bisector of $\angle MOP$, which will be a perpendicular bisector of MP at X because the triangle is equilateral. Thus, $MX = a$. In $\triangle OMX$, $OM = 2a$, $\angle MOX = 30^\circ$, $\angle OXM = 90^\circ$ because each angle in an equilateral triangle is 60° .

$$\sin MOX = \frac{MX}{OM} = \frac{1}{2} \Rightarrow \sin 30^\circ = \frac{1}{2}$$

Similarly, $\angle OMX = 60^\circ$ because the sum of all angles of a triangle is 180° .

$$\cos OMX = \frac{MX}{OM} = \frac{1}{2} \Rightarrow \cos 60^\circ = \frac{1}{2}$$

All other trigonometric ratios can be found from these two.

3.3 Angle of 0°



Consider the $\triangle MOP$ such that side MP is smaller than any quantity we can assign i.e. what we denote by 0. Thus, $\angle MOP$ is what is called approaching 0 or $\lim_{x \rightarrow 0}$ in terms of calculus. Why we take such a value is because if any angle of a triangle is equal to 0° then the triangle won't exist. Thus these values are limiting values as you will learn in calculus.

However, in this case, $\sin 0^\circ = \frac{MP}{OP} = \frac{0}{OP} = 0$. Other trigonometric ratios can be found from this easily.

3.4 Angle of 90°


In the previous figure, as $\angle OMP$ will approach 0° , the $\angle OPM$ will approach 90° . Also, OP will approach the length of OM . Similar to previous case, in right-angle triangle if one angle (other than right angle) approaches 0° the other one will approach 90° and at that value the triangle will cease to exist.

Thus, $\sin 90^\circ = \frac{OM}{OP} = \frac{OP}{OP} = 1$. Now other angles can be found easily from this.

Given below is a table of most useful angles:

Angle	0°	30°	45°	60°	90°
sin	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
tan	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	∞
cot	∞	$\sqrt{3}$	1	$\frac{1}{\sqrt{3}}$	0
sec	1	$\frac{2}{\sqrt{3}}$	$\sqrt{2}$	2	∞
cosec	∞	2	$\sqrt{2}$	$\frac{2}{\sqrt{3}}$	1

3.5 Complementary Angles



Angles are said to be complementary if their sum is equal to one right angle i.e. 90° . Thus, if measure of one angle is θ the other will automatically be $90^\circ - \theta$.

Consider the figure. $\triangle OMP$ is a right-angle triangle, whose $\angle OMP$ is a right angle. Since the sum of all angles is 180° , therefore sum of $\angle MOP$ and $\angle MPO$ will be equal to one right angle or 90° i.e. they are complementary angles.

Let $\angle MPO = \theta$ then $\angle MOP = 90^\circ - \theta$. When $\angle MPO$ is considered MP becomes the base and OM becomes the perpendicular.


$$\text{Thus, } \sin(90^\circ - \theta) = \sin MOP = \frac{MP}{OP} = \cos MPO = \cos \theta$$

$$\cos(90^\circ - \theta) = \sin MPO = \frac{MO}{OP} = \sin \theta$$

$$\tan(90^\circ - \theta) = \tan MOP = \frac{PM}{OM} = \cot MPO = \cot \theta$$

Similarly, $\cot(90^\circ - \theta) = \tan \theta$, $\operatorname{cosec}(90^\circ - \theta) = \sec \theta$, $\sec(90^\circ - \theta) = \operatorname{cosec} \theta$.

3.6 Supplementary Angles



Angles are said to be supplementary if their sum is equal to two right angles i.e. 180° . Thus, if measure of one angle is θ , the other will automatically be $180^\circ - \theta$.

Consider the above figure which includes the angles of $180^\circ - \theta$. In each figure OM and OM' are drawn in different directions, while MP and $M'P'$ are drawn in the same direction so that $OM' = -OM$ and $M'P' = MP$. Hence we can say that


$$\sin(180^\circ - \theta) = \sin MOP' = \frac{M'P'}{OP'} = \frac{MP}{OP} = \sin \theta$$

$$\cos(180^\circ - \theta) = \cos MOP' = \frac{OM'}{OP'} = -\frac{OM}{OP} = -\cos \theta$$

$$\tan(180^\circ - \theta) = \tan MOP' = \frac{OM'}{M'P'} = -\frac{OM}{MP} = -\tan \theta$$

$$\text{Similarly, } \cot(180^\circ - \theta) = -\cot \theta, \sec(180^\circ - \theta) = -\sec \theta, \operatorname{cosec}(180^\circ - \theta) = \operatorname{cosec} \theta$$

3.7 Angles of $-\theta$



Consider the above diagram which plots the angles of θ and $-\theta$. Note that MP and MP' are equal in magnitude but opposite in sign. Thus, we have

$$\sin(-\theta) = \frac{MP'}{OP'} = -\frac{MP}{OP} = -\sin \theta.$$

$$\cos(-\theta) = \frac{OM}{MP'} = \frac{OM}{OP} = \cos \theta.$$

$$\tan(-\theta) = \frac{MP'}{OM} = \frac{-MP}{OM} = -\tan \theta.$$

$$\text{Similarly, } \cot(-\theta) = -\cot \theta, \sec(-\theta) = \sec \theta, \operatorname{cosec}(-\theta) = -\operatorname{cosec} \theta.$$

3.8 Angles of $90^\circ + \theta$

The diagram has been left as an exercise. Similarly, it can be proven that $\sin(90^\circ + \theta) = \cos \theta$, $\cos(90^\circ + \theta) = -\sin \theta$, $\tan(90^\circ + \theta) = -\cot \theta$, $\cot(90^\circ + \theta) = -\tan \theta$, $\sec(90^\circ + \theta) = -\operatorname{cosec} \theta$, $\operatorname{cosec}(90^\circ + \theta) = \sec \theta$.

Angles of $180^\circ + \theta$, $270^\circ - \theta$, $270^\circ + \theta$ can be found using previous relations.

3.9 Angles of $360^\circ + \theta$

For angles of θ the radius vector makes an angle of θ with initial side. For angles of $360^\circ + \theta$ it will complete a full revolution and then make an angle of θ with initial side. Thus, the trigonometrical ratios for an angle of $360^\circ + \theta$ are the same as those for θ .

It is clear that angle will remain θ for any multiple of 360° .

3.10 Problems

1. If $A = 30^\circ$, verify that
 - a) $\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1$
 - b) $\sin 2A = 2 \sin A \cos A$
 - c) $\cos 3A = 4 \cos^3 A - 3 \cos A$
 - d) $\sin 3A = 3 \sin A - 4 \sin^3 A$
 - e) $\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$
2. If $A = 45^\circ$, verify that
 - a) $\sin 2A = 2 \sin A \cos A$
 - b) $\cos 2A = 1 - 2 \sin^2 A$
 - c) $\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$

Verify that

3. $\sin^2 30^\circ + \sin^2 45^\circ + \sin^2 60^\circ = \frac{3}{2}$
4. $\tan^2 30^\circ + \tan^2 45^\circ + \tan^2 60^\circ = 4\frac{1}{3}$
5. $\sin 30^\circ \cos 60^\circ + \sin 60^\circ \cos 30^\circ = 1$
6. $\cos 45^\circ \cos 60^\circ - \sin 45^\circ \sin 60^\circ = -\frac{\sqrt{3}-1}{2\sqrt{2}}$
7. $\operatorname{cosec}^2 45^\circ \cdot \sec^2 30^\circ \cdot \sin^2 90^\circ \cdot \cos 60^\circ = 1\frac{1}{3}$
8. $4 \cot^2 45^\circ - \sec^2 60^\circ + \sin^2 30^\circ = \frac{1}{4}$

Prove that

9. $\sin 420^\circ \cos 390^\circ + \cos(-300^\circ) \sin(-330^\circ) = 1$

10. $\cos 570^\circ \sin 510^\circ - \sin 330^\circ \cos 390^\circ = 0$

What are the values of $\cos A - \sin A$ and $\tan A + \cot A$ when A has the values

11. $\frac{\pi}{3}$

12. $\frac{2\pi}{3}$

13. $\frac{5\pi}{4}$

14. $\frac{7\pi}{4}$

15. $\frac{11\pi}{3}$

What values between 0° and 360° may A have when

16. $\sin A = \frac{1}{\sqrt{2}}$

17. $\cos A = -\frac{1}{2}$

18. $\tan A = -1$

19. $\cot A = -\sqrt{3}$

20. $\sec A = -\frac{2}{\sqrt{3}}$

21. $\operatorname{cosec} A = -2$

Express in terms of the ratios of a positive angle, which is less than 45° , the quantities

22. $\sin(-65^\circ)$

23. $\cos(-84^\circ)$

24. $\tan 137^\circ$

25. $\sin 168^\circ$

26. $\cos 287^\circ$

27. $\tan(-246^\circ)$

28. $\sin 843^\circ$

- 29. $\cos(-928^\circ)$
- 30. $\tan 1145^\circ$
- 31. $\cos 1410^\circ$
- 32. $\cot(-1054^\circ)$
- 33. $\sec 1327^\circ$
- 34. $\operatorname{cosec}(-756^\circ)$

What sign has $\sin A + \cos A$ for the following values of A ?

- 35. 140°
- 36. 278°
- 37. -356°
- 38. -1125°

What sign has $\sin A - \cos A$ for the following values of A ?

- 39. 215°
- 40. 825°
- 41. -634°
- 42. -457°

- 43. Find the sine and cosine of all angles in the first four quadrants whose tangents are equal to $\cos 135^\circ$.

Prove that

- 44. $\sin(270^\circ + A) = -\cos A$ and $\tan(270^\circ + A) = -\cot A$
- 45. $\cos(270^\circ - A) = -\sin A$ and $\cot(270^\circ - A) = \tan A$
- 46. $\cos A + \sin(270^\circ + A) - \sin(270^\circ - A) + \cos(180^\circ + A) = 0$
- 47. $\sec(270^\circ - A) \sec(90^\circ - A) - \tan(270^\circ - A) \tan(90^\circ + A) + 1 = 0$
- 48. $\cot A + \tan(180^\circ + A) + \tan(90^\circ + A) + \tan(360^\circ - A) = 0$
- 49. Find the value of $3 \tan^2 45^\circ - \sin^2 60^\circ - \frac{1}{2} \cot^2 30^\circ + \frac{1}{8} \sec^2 45^\circ$
- 50. Simplify $\frac{\sin 300^\circ \cdot \tan 330^\circ \cdot \sec 420^\circ}{\tan 135^\circ \cdot \sin 210^\circ \cdot \sec 315^\circ}$

51. Show that $\tan 1^\circ \tan 2^\circ \dots \tan 89^\circ = 1$

52. Show that $\sin^2 5^\circ + \sin^2 10^\circ + \sin^2 15^\circ + \dots + \sin^2 90^\circ = 9\frac{1}{2}$

53. Find the value of $\cos^2 \frac{\pi}{16} + \cos^2 \frac{3\pi}{16} + \cos^2 \frac{5\pi}{16} + \cos^2 \frac{7\pi}{16}$

Find the value of the following:

54. $\sec^2 \frac{\pi}{6} \sec^2 \frac{\pi}{4} + \tan^2 \frac{\pi}{3} \sin^2 \frac{\pi}{2}$

55. $\cot^2 30^\circ - 2 \cos^2 60^\circ - \frac{3}{4} \sin^2 45^\circ - 4 \sin^2 30^\circ$

56. $\frac{\sec 480^\circ \operatorname{cosec} 570^\circ \cdot \tan 330^\circ}{\sin 600^\circ \cdot \cos 660^\circ \cdot \cot 405^\circ}$

57. If $A = 30^\circ$, show that $\cos^6 A + \sin^6 A = 1 - \sin^2 A \cos^2 A$

58. Show that $\left(\tan \frac{\pi}{4} + \cot \frac{\pi}{4} + \sec \frac{\pi}{4} \right) \left(\tan \frac{\pi}{4} + \cot \frac{\pi}{4} - \sec \frac{\pi}{4} \right) = \operatorname{cosec}^2 \frac{\pi}{4}$

59. Show that $\sin^2 6^\circ + \sin^2 12^\circ + \sin^2 18^\circ + \dots + \sin^2 84^\circ + \sin^2 90^\circ = 8$

60. Show that $\tan 9^\circ \cdot \tan 27^\circ \cdot \tan 45^\circ \cdot \tan 63^\circ \cdot \tan 81^\circ = 1$

61. Show that $\sum_{r=1}^9 \sin^2 \frac{r\pi}{18} = 5$

62. If $4n\alpha = \pi$, show that $\tan \alpha \tan 2\alpha \tan 3\alpha \dots \tan(2n-2)\alpha \tan(2n-1)\alpha = 1$

COMPOUND ANGLES


Algebraic sum of two or more angles is called a *compound angle*. If A, B, C are any angle then $A + B, A - B, A - B + C, A + B + C, A - B - C, A + B - C$ etc. are all compound angles.

4.1 The Addition Formula

$$\sin(A + B) = \sin A \cos B + \sin B \cos A$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$



Consider the diagram above. PM and PN are perpendicular to OQ and ON . RN is parallel to OQ and NQ is perpendicular to OQ . The left diagram represents the case

when sum of angles is an acute angle while the right diagram represents the case when sum of angles is an obtuse angle.

$$\angle RPN = 90^\circ - \angle PNR = \angle RNO = \angle NOQ = \angle A$$

$$\begin{aligned} \text{Now we can write, } \sin(A + B) &= \sin QOP = \frac{MP}{OP} = \frac{MR + RP}{OP} = \frac{QN}{OP} + \frac{RP}{OP} \\ &= \frac{QN}{ON} \frac{ON}{OP} + \frac{RP}{NP} \frac{NP}{OP} = \sin A \cos B + \cos A \sin B \end{aligned}$$

$$\begin{aligned} \text{Also, } \cos(A + B) &= \cos QOP = \frac{OM}{OP} = \frac{OQ - MQ}{OP} = \frac{OQ}{ON} \frac{ON}{OP} - \frac{RN}{NP} \frac{NP}{OP} \\ &= \cos A \cos B - \sin A \sin B \end{aligned}$$

$$\text{These two results lead to } \tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

We have shown that addition formula is true when angles involved are acute angles. The same proof can be applied to prove the results for all values of A and B .

$$\text{Consider } A' = 90^\circ + A \therefore \sin A' = \cos A \text{ and } \cos A' = \sin A$$

$$\sin(A' + B) = \cos(A + B) = \cos A \cos B - \sin A \sin B = \sin A' \cos B + \cos A' \sin B$$

$$\text{Similarly } \cos(A' + B) = -\sin(A + B) = -\sin A \cos B - \sin B \cos A = \cos A' \cos B - \sin A' \sin B$$


We can prove it again for $B' = 90^\circ + B$ and so on by increasing the values of A and B . Then we can again increase values by 90° and proceeding this way we see that the formula holds true for all values of A and B .

4.2 The Subtraction Formula

$$\sin(A - B) = \sin A \cos B - \sin B \cos A$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$



Consider the diagram above. The angle MOP is $A - B$. We take a point P , and draw PM and PN perpendicular to OM and ON respectively. From N we draw NQ and NR perpendicular to OQ and MP respectively.

$$\angle RPN = 90^\circ - \angle PNR = \angle QON = A$$

$$\text{Thus, we can write } \sin(A - B) = \sin MOP = \frac{MP}{OP} = \frac{MR - PR}{OP} = \frac{QN}{ON} \frac{ON}{OP} - \frac{PR}{PN} \frac{PN}{OP}$$

$$\text{Thus, } \sin(A - B) = \sin A \cos B - \cos A \sin B$$

$$\begin{aligned} \text{Also, } \cos(A - B) &= \frac{OM}{OP} = \frac{OQ + QM}{OP} = \frac{OQ}{ON} \frac{ON}{OP} + \frac{RN}{NP} \frac{NP}{OP} \\ &= \cos A \cos B + \sin A \sin B \end{aligned}$$

We have shown that subtraction formula is true when angles involved are acute angles. The same proof can be applied to prove the results for all values of A and B .

$$\text{From the results obtained we find upon division that } \tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

4.3 Important Deductions

- $\sin(A + B) \sin(A - B) = \sin^2 A - \sin^2 B = \cos^2 B - \cos^2 A$
 L.H.S. $= (\sin A \cos B + \sin B \cos A)(\sin A \cos B - \sin B \cos A)$
 $= \sin^2 A \cos^2 B - \sin^2 B \cos^2 A = \sin^2 A(1 - \sin^2 B) - \sin^2 B(1 - \sin^2 A)$
 $= \sin^2 A - \sin^2 A \sin^2 B - \sin^2 B + \sin^2 B \sin^2 A$
 $= \sin^2 A - \sin^2 B = (1 - \cos^2 A) - (1 - \cos^2 B)$
 $= \cos^2 B - \cos^2 A$
- $\cos(A + B) \cos(A - B) = \cos^2 A - \sin^2 B = \cos^2 B - \sin^2 A$
 L.H.S. $= (\cos A \cos B - \sin A \sin B)(\cos A \cos B + \sin A \sin B)$

$$\begin{aligned}
&= \cos^2 A \cos^2 B - \sin^2 A \sin^2 B \\
&= \cos^2 A (1 - \sin^2 B) - (1 - \cos^2 A) \sin^2 B \\
&= \cos^2 A - \cos^2 A \sin^2 B - \sin^2 B + \cos^2 A \sin^2 B \\
&= \cos^2 A - \sin^2 B = \cos^2 B - \sin^2 A
\end{aligned}$$

$$3. \cot(A + B) = \frac{\cot A \cot B - 1}{\cot B + \cot A}$$

$$\text{L.H.S.} = \cot(A + B) = \frac{\cos(A + B)}{\sin(A + B)}$$

$$= \frac{\cos A \cos B - \sin A \sin B}{\sin A \cos B + \cos A \sin B}$$

Dividing numerator and denominator by $\sin A \sin B$

$$= \frac{\cot A \cot B - 1}{\cot B + \cot A}$$

$$4. \cot(A - B) = \frac{\cot A \cot B + 1}{\cot B - \cot A}$$

$$\text{L.H.S.} = \cot(A - B) = \frac{\cos(A - B)}{\sin(A - B)}$$

$$= \frac{\cos A \cos B + \sin A \sin B}{\sin A \cos B - \cos A \sin B}$$

Dividing numerator and denominator by $\sin A \sin B$

$$= \frac{\cot A \cot B + 1}{\cot B - \cot A}$$

$$5. \tan(A + B + C) = \frac{\tan A + \tan B + \tan C - \tan A \tan B \tan C}{1 - \tan A \tan B - \tan B \tan C - \tan C \tan A}$$

$$\text{L.H.S.} = \tan[(A + B) + C] = \frac{\tan(A + B) + \tan C}{1 - \tan(A + B) \tan C}$$

$$= \frac{\frac{\tan A + \tan B}{1 - \tan A \tan B} + \tan C}{1 - \frac{\tan A + \tan B}{1 - \tan A \tan B} \tan C}$$

$$= \frac{\frac{\tan A + \tan B + \tan C - \tan A \tan B \tan C}{1 - \tan A \tan B}}{1 - \tan A \tan B - \tan B \tan C - \tan C \tan A}$$

$$= \frac{\tan A + \tan B + \tan C - \tan A \tan B \tan C}{1 - \tan A \tan B - \tan B \tan C - \tan C \tan A}$$

4.4 To express $a \cos \theta + b \sin \theta$ in the form of $k \cos \phi$ or $k \sin \phi$

$$a \cos \theta + b \sin \theta = \sqrt{a^2 + b^2} \left(\frac{a}{\sqrt{a^2 + b^2}} \cos \theta + \frac{b}{\sqrt{a^2 + b^2}} \sin \theta \right)$$

Let $\cos \alpha = \frac{a}{\sqrt{a^2 + b^2}}$ then $\sin \alpha = \frac{b}{\sqrt{a^2 + b^2}}$

Thus, $a \cos \theta + b \sin \theta = \sqrt{a^2 + b^2}(\cos \alpha \cos \theta + \sin \alpha \sin \theta)$
 $= \sqrt{a^2 + b^2} \cos(\theta - \alpha) = k \cos \phi$ where $k = \sqrt{a^2 + b^2}$ and $\phi = \theta - \alpha$

Alternatively, if $\frac{a}{\sqrt{a^2 + b^2}} = \sin \alpha$ then $\frac{b}{\sqrt{a^2 + b^2}} = \cos \alpha$

Thus, $a \cos \theta + b \sin \theta = \sqrt{a^2 + b^2}(\sin \alpha \cos \theta + \cos \alpha \sin \theta)$
 $= \sqrt{a^2 + b^2} \sin(\theta + \alpha) = k \sin \phi$ where $k = \sqrt{a^2 + b^2}$ and $\phi = \theta + \alpha$

4.5 Problems

1. If $\sin \alpha = \frac{3}{5}$ and $\cos \beta = \frac{9}{41}$, find the values of $\sin(\alpha - \beta)$ and $\cos(\alpha + \beta)$.
2. If $\sin \alpha = \frac{45}{53}$ and $\sin \beta = \frac{33}{65}$, find the values of $\sin(\alpha - \beta)$ and $\sin(\alpha + \beta)$.
3. If $\sin \alpha = \frac{15}{17}$ and $\cos \beta = \frac{12}{13}$, find the values of $\sin(\alpha + \beta)$, $\cos(\alpha - \beta)$ and $\tan(\alpha + \beta)$.

Prove the following:

1. $\cos(45^\circ - A) \cos(45^\circ - B) - \sin(45^\circ - A) \sin(45^\circ - B) = \sin(A + B)$.
2. $\sin(45^\circ + A) \cos(45^\circ - B) + \cos(45^\circ + A) \sin(45^\circ - B) = \cos(A - B)$.
3. $\frac{\sin(A - B)}{\cos A \cos B} + \frac{\sin(B - C)}{\cos B \cos C} + \frac{\sin(C - A)}{\cos C \cos A} = 0$.
4. $\sin 105^\circ + \cos 105^\circ = \cos 45^\circ$.
5. $\sin 75^\circ - \sin 15^\circ = \cos 105^\circ + \cos 15^\circ$.
6. $\cos \alpha \cos(\gamma - \alpha) - \sin \alpha \sin(\gamma - \alpha) = \cos \gamma$.
7. $\cos(\alpha + \beta) \cos \gamma - \cos(\beta + \gamma) \cos \alpha = \sin \beta \sin(\gamma - \alpha)$.
8. $\sin(n + 1)A \sin(n - 1)A + \cos(n + 1)A \cos(n - 1)A = \cos 2A$.
9. $\sin(n + 1)A \sin(n + 2)A + \cos(n + 1)A \cos(n + 2)A = \cos A$.
10. Find the value of $\cos 15^\circ$ and $\sin 105^\circ$.
11. Find the value of $\tan 105^\circ$.
12. Find the value of $\frac{\tan 495^\circ}{\cot 855^\circ}$.

13. Evaluate $\sin\left(n\pi + (-1)^n \frac{\pi}{4}\right)$, where n is an integer.

Prove the following:

14. $\sin 15^\circ = \frac{\sqrt{3} - 1}{2\sqrt{2}}.$

15. $\cos 75^\circ = \frac{\sqrt{3} - 1}{2\sqrt{2}}.$

16. $\tan 75^\circ = 2 + \sqrt{3}.$

17. $\tan 15^\circ = 2 - \sqrt{3}.$

Find the value of following:

18. $\cos 1395^\circ.$

19. $\tan(-330^\circ).$

20. $\sin 300^\circ \operatorname{cosec} 1050^\circ - \tan(-120^\circ).$

21. $\tan\left(\frac{11\pi}{12}\right).$

22. $\tan\left((-1)^n \frac{\pi}{4}\right).$

Prove the following:

23. $\cos 18^\circ - \sin 18^\circ = \sqrt{2} \sin 27^\circ.$

24. $\tan 70^\circ = 2 \tan 50^\circ + \tan 20^\circ.$

25. $\cot\left(\frac{\pi}{4} + x\right) \cot\left(\frac{\pi}{4} - x\right) = 1.$

26. $\cos(m+n)\theta \cdot \cos(m-n)\theta - \sin(m+n)\theta \sin(m-n)\theta = \cos 2m\theta.$

27. $\frac{\tan(\theta + \phi) + \tan(\theta - \phi)}{1 - \tan(\theta + \phi) \tan(\theta - \phi)} = \tan 2\theta.$

28. $\cos 9^\circ + \sin 9^\circ = \sqrt{2} \sin 54^\circ.$

29. $\frac{\cos 20^\circ - \sin 20^\circ}{\cos 20^\circ + \sin 20^\circ} = \tan 25^\circ.$

30. $\frac{\tan A + \tan B}{\tan A - \tan B} = \frac{\sin(A + B)}{\sin(A - B)}.$

31. $\frac{1}{\tan 3A - \tan A} - \frac{1}{\cot 3A - \cot A} = \cot 2A.$
32. $\frac{1}{\tan 3A + \tan A} - \frac{1}{\cot 3A - \cot A} = \cot 4A.$
33. $\frac{\sin 3\alpha}{\sin \alpha} + \frac{\cos 3\alpha}{\cos \alpha} = 4 \cos 2\alpha.$
34. $\frac{\tan\left(\frac{\pi}{4} + A\right) - \tan\left(\frac{\pi}{4} - A\right)}{\tan\left(\frac{\pi}{4} + A\right) + \tan\left(\frac{\pi}{4} - A\right)} = \sin 2A.$
35. $\tan 40^\circ + 2 \tan 10^\circ = \tan 50^\circ.$
36. $\tan(\alpha + \beta) \tan(\alpha - \beta) = \frac{\sin^2 \alpha - \sin^2 \beta}{\cos^2 \alpha - \sin^2 \beta}.$
37. $\tan^2 \alpha - \tan^2 \beta = \frac{\sin(\alpha + \beta) \sin(\alpha - \beta)}{\cos^2 \alpha \cos^2 \beta}.$
38. $\tan[(2n + 1)\pi + \theta] + \tan[(2n + 1)\pi - \theta] = 0.$
39. $\tan\left(\frac{\pi}{4} + \theta\right) \tan\left(\frac{3\pi}{4} + \theta\right) + 1 = 0.$
40. If $\tan \alpha = p$ and $\tan \beta = q$ prove that $\cos(\alpha + \beta) = \frac{1 - pq}{\sqrt{(1 + p^2)(1 + q^2)}}.$
41. if $\tan \beta = \frac{2 \sin \alpha \sin \gamma}{\sin(\alpha + \gamma)},$ show that $\cot \alpha, \cot \beta, \cot \gamma$ are in A.P.
42. Eliminate θ if $\tan(\theta - \alpha) = a$ and $\tan(\theta + \alpha) = b.$
43. Eliminate α and β if $\tan \alpha + \tan \beta = b, \cot \alpha + \cot \beta = a$ and $\alpha + \beta = \gamma.$
44. If $A + B = 45^\circ,$ show that $(1 + \tan A)(1 + \tan B) = 2.$
45. If $\sin \alpha \sin \beta - \cos \alpha \cos \beta + 1 = 0,$ prove that $1 + \cot \alpha \tan \beta = 0.$
46. If $\tan \beta = \frac{n \sin \alpha \cos \alpha}{1 - n \sin^2 \alpha},$ prove that $\tan(\alpha - \beta) = (1 - n)\alpha.$
47. If $\cos(\beta - \gamma) + \cos(\gamma - \alpha) + \cos(\alpha - \beta) = -\frac{3}{2},$ prove that $\cos \alpha + \cos \beta + \cos \gamma = \sin \alpha + \sin \beta + \sin \gamma = 0.$
48. If $\tan \alpha = \frac{m}{m + 1}, \tan \beta = \frac{1}{2m + 1},$ prove that $\alpha + \beta = \frac{\pi}{4}.$
49. If $A + B = 45^\circ,$ show that $(\cot A - 1)(\cot B - 1) = 2.$
50. If $\tan \alpha - \tan \beta = x$ and $\cot \beta - \cot \alpha = y,$ prove that $\cot(\alpha - \beta) = \frac{x + y}{xy}.$
51. If a right angle be divided into three parts α, β and $\gamma,$ prove that $\cot \alpha = \frac{\tan \beta + \tan \gamma}{1 - \tan \beta \tan \gamma}.$

52. If $2 \tan \beta + \cot \beta = \tan \alpha$, show that $\cot \beta = 2 \tan(\alpha - \beta)$.
53. If in any $\triangle ABC$, $C = 90^\circ$, prove that $\operatorname{cosec}(A - B) = \frac{a^2 + b^2}{a^2 - b^2}$ and $\sec(A - B) = \frac{c^2}{2ab}$.
54. If $\cot A = \sqrt{ac}$, $\cot B = \sqrt{\frac{c}{a}}$, $\tan C = \sqrt{\frac{c}{a^3}}$ and $c = a^2 + a + 1$, prove that $A = B + C$.
55. If $\frac{\tan(A - B)}{\tan A} + \frac{\sin^2 C}{\sin^2 A} = 1$, prove that $\tan A \tan B = \tan^2 C$.
56. If $\sin \alpha \sin \beta - \cos \alpha \cos \beta = 1$ show that $\tan \alpha + \tan \beta = 0$.
57. If $\sin \theta = 3 \sin(\theta + 2\alpha)$, prove that $\tan(\theta + \alpha)$, prove that $\tan(\theta + \alpha) + 2 \tan \alpha = 0$.
58. If $3 \tan \theta \tan \phi = 1$, prove that $2 \cos(\theta + \phi) = \cos(\theta - \alpha)$.
59. Find the sign of the expression $\sin \theta + \cos \theta$ when $\theta = 100^\circ$.
60. Prove that the value of $5 \cos \theta + 3 \cos\left(\theta + \frac{\pi}{3}\right) + 3$ lies between -4 and 10 .
61. If $m \tan(\theta - 30^\circ) = n \tan(\theta + 120^\circ)$, show that $\cos 2\theta = \frac{m + n}{2(m - n)}$.
62. if $\alpha + \beta = \theta$ and $\tan \alpha : \tan \beta = x : y$, prove that $\sin(\alpha - \beta) = \frac{x - y}{x + y} \sin \theta$.
63. Find the maximum and minimum value of $7 \cos \theta + 24 \sin \theta$.
64. Show that $\sin 100^\circ - \sin 10^\circ$ is positive.

TRANSFORMATION FORMULAE

5.1 Transformation of products into sums or differences

We know that $\sin(A + B) = \sin A \cos B + \cos A \sin B$ and $\sin(A - B) = \sin A \cos B - \cos A \sin B$

Adding these, we get $2 \sin A \cos B = \sin(A + B) + \sin(A - B)$

Subtracting, we get $2 \cos A \sin B = \sin(A + B) - \sin(A - B)$

We also know that $\cos(A + B) = \cos A \cos B - \sin A \sin B$ and $\cos(A - B) = \cos A \cos B + \sin A \sin B$

Adding, we get $2 \cos A \cos B = \cos(A + B) + \cos(A - B)$

Subtrating we get $2 \sin A \sin B = \cos(A - B) - \cos(A + B)$

5.2 Transformation of sums or differences into products

We have $2 \sin A \cos B = \sin(A + B) \sin(A - B)$

Substituting for $A + B = C, A - B = D$ so that $A = \frac{C + D}{2}$ and $B = \frac{C - D}{2}$

$$\sin C + \sin D = 2 \sin \frac{C + D}{2} \cos \frac{C - D}{2}$$

We also have $2 \cos A \sin B = \sin(A + B) - \sin(A - B)$

Following similarly $\sin C - \sin D = 2 \cos \frac{C+D}{2} \sin \frac{C-D}{2}$

For $2 \cos A \cos B = \cos(A+B) + \cos(A-B)$, we get $\cos C + \cos D = 2 \cos \frac{C+D}{2} \cos \frac{C-D}{2}$

For $2 \sin A \sin B = \cos(A-B) - \cos(A+B)$, we get $\cos C - \cos D = 2 \sin \frac{C+D}{2} \sin \frac{D-C}{2}$

5.3 Problems

- Find the value of $\frac{\sin 75^\circ - \sin 15^\circ}{\cos 75^\circ + \cos 15^\circ}$.
- Simplify the expression $\frac{(\cos \theta - \cos 3\theta)(\sin 8\theta + \sin 2\theta)}{(\sin 5\theta - \sin \theta)(\cos 4\theta - \cos 6\theta)}$.

Prove that

- $\frac{\sin 7\theta - \sin 5\theta}{\cos 7\theta + \cos 5\theta} = \tan \theta$.
- $\frac{\cos 6\theta - \cos 4\theta}{\sin 6\theta + \sin 4\theta} = -\tan \theta$.
- $\frac{\sin A + \sin 3A}{\cos A + \cos 3A} = \tan 2A$.
- $\frac{\sin 7A - \sin A}{\sin 8A - \sin 2A} = \cos 4A \sec 5A$.
- $\frac{\cos 2B + \cos 2A}{\cos 2B - \cos 2A} = \cot(A+B) \cot(A-B)$.
- $\frac{\sin 2A + \sin 2B}{\sin 2A - \sin 2B} = \frac{\tan(A+B)}{\tan(A-B)}$.
- $\frac{\sin A + \sin 2A}{\cos A - \cos 2A} = \cot \frac{A}{2}$.
- $\frac{\sin 5A - \sin 3A}{\cos 3A + \cos 5A} = \tan A$.
- $\frac{\cos 2B - \cos 2A}{\sin 2B + \sin 2A} = \tan(A-B)$.
- $\cos(A+B) + \sin(A-B) = 2 \sin(45^\circ + A) \cos(45^\circ + B)$.
- $\frac{\cos 3A - \cos A}{\sin 3A - \sin A} + \frac{\cos 2A - \cos 4A}{\sin 4A - \sin 2A} = \frac{\sin A}{\cos 2A \cos 3A}$.

14. $\frac{\sin(4A - 2B) + \sin(4B - 2A)}{\cos(4A - 2B) + \cos(4B - 2A)} = \tan(A + B).$
15. $\frac{\tan 5\theta + \tan 3\theta}{\tan 5\theta - \tan 3\theta} = 4 \cos 2\theta \cos 4\theta.$
16. $\frac{\cos 3\theta + 2 \cos 5\theta + \cos 7\theta}{\cos \theta + 2 \cos 3\theta + \cos 5\theta} = \cos 2\theta - \sin 2\theta \tan 3\theta.$
17. $\frac{\sin A + \sin 3A + \sin 5A + \sin 7A}{\cos A + \cos 3A + \cos 5A + \cos 7A} = \tan 4A.$
18. $\frac{\sin(\theta + \phi) - 2 \sin \theta + \sin(\theta - \phi)}{\cos(\theta + \phi) - 2 \cos \theta + \cos(\theta - \phi)} = \tan \theta.$
19. $\frac{\sin A + 2 \sin 3A + \sin 5A}{\sin 3A + 2 \sin 5A + \sin 7A} = \frac{\sin 3A}{\sin 5A}.$
20. $\frac{\sin(A - C) + 2 \sin A + \sin(A + C)}{\sin(B - C) + 2 \sin B + \sin(B + C)} = \frac{\sin A}{\sin B}.$
21. $\frac{\sin A - \sin 5A + \sin 9A - \sin 13A}{\cos A - \cos 5A - \cos 9A + \cos 13A} = \cot 4A.$
22. $\frac{\sin A + \sin B}{\sin A - \sin B} = \tan \frac{A + B}{2} \cot \frac{A - B}{2}.$
23. $\frac{\cos A + \cos B}{\cos B - \cos A} = \cot \frac{A + B}{2} \cot \frac{A - B}{2}.$
24. $\frac{\sin A + \sin B}{\cos A + \cos B} = \tan \frac{A + B}{2}.$
25. $\frac{\sin A - \sin B}{\cos B - \cos A} = \cot \frac{A + B}{2}.$
26. $\frac{\cos(A + B + C) + \cos(-A + B + C) + \cos(A - B + C) + \cos(A + B - C)}{\sin(A + B + C) + \sin(-A + B + C) - \sin(A - B + C) + \sin(A + B - C)} = \cot B.$
27. $\cos 3A + \cos 5A + \cos 7A + \cos 15A = 4 \cos 4A \cos 5A \cos 6A.$
28. $\cos(-A + B + C) + \cos(A - B + C) + \cos(A + B - C) + \cos(A + B + C) = 4 \cos A \cos B \cos C.$
29. $\sin 50^\circ - \sin 70^\circ + \sin 10^\circ = 0.$
30. $\sin 10^\circ + \sin 20^\circ + \sin 40^\circ + \sin 50^\circ = \sin 70^\circ + \sin 80^\circ.$
31. $\sin \alpha + \sin 2\alpha + \sin 4\alpha + \sin 5\alpha = 4 \cos \frac{\alpha}{2} \cos \frac{3\alpha}{2} \sin 3\alpha.$

Simplify:

$$32. \cos \left[\theta + \left(n - \frac{3}{2} \right) \phi \right] - \cos \left[\theta + \left(n + \frac{3}{2} \right) \phi \right].$$

$$33. \sin \left[\theta + \left(n - \frac{3}{2} \right) \phi \right] + \sin \left[\theta + \left(n + \frac{3}{2} \right) \phi \right].$$

Express as a sum or difference the following:

$$34. 2 \sin 5\theta \sin 7\theta.$$

$$35. 2 \cos 7\theta \sin 5\theta.$$

$$36. 2 \cos 11\theta \cos 3\theta.$$

$$37. 2 \sin 54^\circ \sin 66^\circ.$$

Prove that

$$38. \sin \frac{\theta}{2} \sin \frac{7\theta}{2} + \sin \frac{3\theta}{2} \sin \frac{11\theta}{2} = \sin 2\theta \sin 5\theta.$$

$$39. \cos 2\theta \cos \frac{\theta}{2} - \cos 3\theta \cos \frac{9\theta}{2} = \sin 5\theta \sin \frac{5\theta}{2}.$$

$$40. \sin A \sin(A + 2B) - \sin B \sin(B + 2A) = \sin(A - B) \sin(A + B).$$

$$41. (\sin 3A + \sin A) \sin A + (\cos 3A - \cos A) \cos A = 0.$$

$$42. \frac{2 \sin(A - C) \cos C - \sin(A - 2C)}{2 \sin(B - C) \cos C - \sin(B - 2C)} = \frac{\sin A}{\sin B}.$$

$$43. \frac{\sin A \sin 2A + \sin 3A \sin 6A + \sin 4A \sin 13A}{\sin A \cos 2A + \sin 3A \cos 6A + \sin 4A \cos 13A} = \tan 9A.$$

$$44. \frac{\cos 2A \cos 3A - \cos 2A \cos 7A + \cos A \cos 10A}{\sin 4A \sin 3A - \sin 2A \sin 5A + \sin 4A \sin 7A} = \cot 6A \cot 5A.$$

$$45. \cos(36^\circ - A) \cos(36^\circ + A) + \cos(54^\circ + A) \cos(54^\circ - A) = \cos 2A.$$

$$46. \cos A \sin(B - C) + \cos B \sin(C - A) + \cos C \sin(A - B) = 0.$$

$$47. \sin(45^\circ + A) \sin(45^\circ - A) = \frac{1}{2} \cos 2A.$$

$$48. \sin(\beta - \gamma) \cos(\alpha - \delta) + \sin(\gamma - \alpha) \cos(\beta - \delta) + \sin(\alpha - \beta) \cos(\gamma - \delta) = 0.$$

$$49. 2 \cos \frac{\pi}{13} \cos \frac{9\pi}{13} + \cos \frac{3\pi}{13} + \cos \frac{5\pi}{13} = 0.$$

$$50. \cos 55^\circ + \cos 65^\circ + \cos 175^\circ = 0.$$

$$51. \cos 18^\circ - \sin 18^\circ = \sqrt{2} \sin 27^\circ.$$

$$52. \frac{\sin A + \sin 2A + \sin 4A + \sin 5A}{\cos A + \cos 2A + \cos 4A + \cos 5A} = \tan 3A.$$

53. $\left(\frac{\cos A + \cos B}{\sin A - \sin B}\right)^n + \left(\frac{\sin A + \sin B}{\cos A - \cos B}\right)^n = 2 \cot^n \frac{A-B}{2}$ or 0 according as n is even or odd.
54. If α, β, γ are in A.P., show that $\cos \beta = \frac{\sin \alpha - \sin \gamma}{\cos \gamma - \cos \alpha}$.
55. If $\sin \theta + \sin \phi = \sqrt{3}(\cos \phi - \cos \theta)$ prove that $\sin 3\theta + \sin 3\phi = 0$.
56. $\sin 65^\circ + \cos 65^\circ = \sqrt{2} \cos 20^\circ$.
57. $\sin 47^\circ + \cos 77^\circ = \cos 17^\circ$.
58. $\frac{\cos 10^\circ - \sin 10^\circ}{\cos 10^\circ + \sin 10^\circ} = \tan 35^\circ$.
59. $\cos 80^\circ + \cos 40^\circ - \cos 20^\circ = 0$.
60. $\cos \frac{\pi}{5} + \cos \frac{2\pi}{5} + \cos \frac{6\pi}{5} + \cos \frac{7\pi}{5} = 0$.
61. $\cos \alpha + \cos \beta + \cos \gamma + \cos(\alpha + \beta + \gamma) = 4 \cos \frac{\alpha + \beta}{2} \cos \frac{\beta + \gamma}{2} \cos \frac{\gamma + \alpha}{2}$.
62. If $\sin \alpha - \sin \beta = \frac{1}{3}$ and $\cos \beta - \cos \alpha = \frac{1}{2}$, prove that $\cot \frac{\alpha + \beta}{2} = \frac{2}{3}$.
63. If $\operatorname{cosec} A + \sec A = \operatorname{cosec} B + \sec B$, prove that $\tan A \tan B = \cot \frac{A+B}{2}$.
64. If $\sec(\theta + \alpha) + \sec(\theta - \alpha) = 2 \sec \theta$, show that $\cos^2 \theta = 1 + \cos \alpha$.
65. Show that $\sin 50^\circ \cos 85^\circ = \frac{1 - \sqrt{2} \sin 35^\circ}{2\sqrt{2}}$.
66. Prove that $\sin 20^\circ \sin 40^\circ \sin 80^\circ = \frac{\sqrt{3}}{8}$.
67. Prove that $\sin A \sin(60^\circ - A) \sin(60^\circ + A) = \frac{1}{4} \sin 3A$.
68. If $\alpha + \beta = 90^\circ$, find the maximum value of $\sin \alpha \sin \beta$.
69. Prove that $\sin 25^\circ \cos 115^\circ = \frac{1}{2}(\sin 40^\circ - 1)$.
70. Prove that $\sin 20^\circ \sin 40^\circ \sin 60^\circ \sin 80^\circ = \frac{3}{16}$.
71. Prove that $\cos 20^\circ \cos 40^\circ \cos 80^\circ = \frac{1}{8}$.
72. Prove that $\tan 20^\circ \tan 40^\circ \tan 60^\circ \tan 80^\circ = 3$.
73. Prove that $\cos 10^\circ \cos 30^\circ \cos 50^\circ \cos 70^\circ = \frac{3}{16}$.
74. Prove that $4 \cos \theta \cos \left(\frac{\pi}{3} + \theta\right) \cos \left(\frac{\pi}{3} - \theta\right) = \cos 3\theta$.

75. Prove that $\tan \theta \tan(60^\circ - \theta) \tan(60^\circ + \theta) = \tan 3\theta$.
76. If $\alpha + \beta = 90^\circ$, show that the maximum value of $\cos \alpha \cos \beta$ is $\frac{1}{2}$.
77. If $\cos \alpha = \frac{1}{\sqrt{2}}$, $\sin \beta = \frac{1}{\sqrt{3}}$, show that $\tan \frac{\alpha + \beta}{2} \cot \frac{\alpha - \beta}{2} = 5 + 2\sqrt{6}$ or $5 - 2\sqrt{6}$.
78. If $x \cos \theta = y \cos \left(\theta + \frac{2\pi}{3} \right) = z \cos \left(\theta + \frac{4\pi}{3} \right)$, prove that $xy + yz + xz = 0$.
79. If $\sin \theta = n \sin(\theta + 2\alpha)$, prove that $\tan(\theta + \alpha) = \frac{1+n}{1-n} \tan \alpha$.
80. If $\frac{\sin(\theta + \alpha)}{\cos(\theta - \alpha)} = \frac{1-m}{1+m}$, prove that $\tan \left(\frac{\pi}{4} - \theta \right) \tan \left(\frac{\pi}{4} - \alpha \right) = m$.
81. If $y \sin \phi = x \sin(2\theta + \phi)$, show that $(x + y) \cot(\theta + \phi) = (y - x) \cot \theta$.
82. If $\cos(\alpha + \beta) \sin(\gamma + \delta) = \cos(\alpha - \beta) \sin(\gamma - \delta)$, prove that $\cot \alpha \cot \beta \cot \gamma = \cot \delta$.
83. If $\frac{\cos(A - B)}{\cos(A + B)} + \frac{\cos(C + D)}{\cos(C - D)} = 0$, prove that $\tan A \tan B \tan C \tan D = -1$.
84. If $\tan(\theta + \phi) = 3 \tan \theta$, prove that $\sin(2\theta + \phi) = 2 \sin \phi$.
85. If $\tan(\theta + \phi) = 3 \tan \theta$, prove that $\sin 2(\theta + \phi) + \sin 2\theta = 2 \sin 2\phi$.

MULTIPLE AND SUBMULTIPLE ANGLES

6.1 Multiple Angles

An angle of the form nA , where n is an integer is called a *multiple angle*. For example, $2A, 3A, 4A, \dots$ are multiple angles of A .

Part II

Solutions

GNU FREE DOCUMENTATION LICENSE

Version 1.3, 3 November 2008

Copyright © 2000, 2001, 2002, 2007, 2008 Free Software Foundation, Inc.

`<http://fsf.org/>`

Everyone is permitted to copy and distribute verbatim copies of this license document, but changing it is not allowed.

Preamble

The purpose of this License is to make a manual, textbook, or other functional and useful document “free” in the sense of freedom: to assure everyone the effective freedom to copy and redistribute it, with or without modifying it, either commercially or noncommercially. Secondly, this License preserves for the author and publisher a way to get credit for their work, while not being considered responsible for modifications made by others.

This License is a kind of “copyleft”, which means that derivative works of the document must themselves be free in the same sense. It complements the GNU General Public License, which is a copyleft license designed for free software.

We have designed this License in order to use it for manuals for free software, because free software needs free documentation: a free program should come with manuals providing the same freedoms that the software does. But this License is not limited to software manuals; it can be used for any textual work, regardless of subject matter or whether it is published as a printed book. We recommend this License principally for works whose purpose is instruction or reference.

1. APPLICABILITY AND DEFINITIONS

This License applies to any manual or other work, in any medium, that contains a notice placed by the copyright holder saying it can be distributed under the terms of this

License. Such a notice grants a world-wide, royalty-free license, unlimited in duration, to use that work under the conditions stated herein. The “**Document**”, below, refers to any such manual or work. Any member of the public is a licensee, and is addressed as “**you**”. You accept the license if you copy, modify or distribute the work in a way requiring permission under copyright law.

A “**Modified Version**” of the Document means any work containing the Document or a portion of it, either copied verbatim, or with modifications and/or translated into another language.

A “**Secondary Section**” is a named appendix or a front-matter section of the Document that deals exclusively with the relationship of the publishers or authors of the Document to the Document’s overall subject (or to related matters) and contains nothing that could fall directly within that overall subject. (Thus, if the Document is in part a textbook of mathematics, a Secondary Section may not explain any mathematics.) The relationship could be a matter of historical connection with the subject or with related matters, or of legal, commercial, philosophical, ethical or political position regarding them.

The “**Invariant Sections**” are certain Secondary Sections whose titles are designated, as being those of Invariant Sections, in the notice that says that the Document is released under this License. If a section does not fit the above definition of Secondary then it is not allowed to be designated as Invariant. The Document may contain zero Invariant Sections. If the Document does not identify any Invariant Sections then there are none.

The “**Cover Texts**” are certain short passages of text that are listed, as Front-Cover Texts or Back-Cover Texts, in the notice that says that the Document is released under this License. A Front-Cover Text may be at most 5 words, and a Back-Cover Text may be at most 25 words.

A “**Transparent**” copy of the Document means a machine-readable copy, represented in a format whose specification is available to the general public, that is suitable for revising the document straightforwardly with generic text editors or (for images composed of pixels) generic paint programs or (for drawings) some widely available drawing editor, and that is suitable for input to text formatters or for automatic translation to a variety of formats suitable for input to text formatters. A copy made in an otherwise Transparent file format whose markup, or absence of markup, has been arranged to thwart or discourage subsequent modification by readers is not Transparent. An image format is not Transparent if used for any substantial amount of text. A copy that is not “Transparent” is called “**Opaque**”.

Examples of suitable formats for Transparent copies include plain ASCII without markup, Texinfo input format, LaTeX input format, SGML or XML using a publicly available DTD, and standard-conforming simple HTML, PostScript or PDF designed for human modification. Examples of transparent image formats include PNG, XCF and JPG. Opaque formats include proprietary formats that can be read and edited only by proprietary word processors, SGML or XML for which the DTD and/or processing tools are not generally available, and the machine-generated HTML, PostScript or PDF produced by some word processors for output purposes only.

The “**Title Page**” means, for a printed book, the title page itself, plus such following pages as are needed to hold, legibly, the material this License requires to appear in the title

page. For works in formats which do not have any title page as such, “Title Page” means the text near the most prominent appearance of the work’s title, preceding the beginning of the body of the text.

The “**publisher**” means any person or entity that distributes copies of the Document to the public.

A section “**Entitled XYZ**” means a named subunit of the Document whose title either is precisely XYZ or contains XYZ in parentheses following text that translates XYZ in another language. (Here XYZ stands for a specific section name mentioned below, such as “**Acknowledgements**”, “**Dedications**”, “**Endorsements**”, or “**History**”.) To “**Preserve the Title**” of such a section when you modify the Document means that it remains a section “Entitled XYZ” according to this definition.

The Document may include Warranty Disclaimers next to the notice which states that this License applies to the Document. These Warranty Disclaimers are considered to be included by reference in this License, but only as regards disclaiming warranties: any other implication that these Warranty Disclaimers may have is void and has no effect on the meaning of this License.

2. VERBATIM COPYING

You may copy and distribute the Document in any medium, either commercially or noncommercially, provided that this License, the copyright notices, and the license notice saying this License applies to the Document are reproduced in all copies, and that you add no other conditions whatsoever to those of this License. You may not use technical measures to obstruct or control the reading or further copying of the copies you make or distribute. However, you may accept compensation in exchange for copies. If you distribute a large enough number of copies you must also follow the conditions in section 3.

You may also lend copies, under the same conditions stated above, and you may publicly display copies.

3. COPYING IN QUANTITY

If you publish printed copies (or copies in media that commonly have printed covers) of the Document, numbering more than 100, and the Document’s license notice requires Cover Texts, you must enclose the copies in covers that carry, clearly and legibly, all these Cover Texts: Front-Cover Texts on the front cover, and Back-Cover Texts on the back cover. Both covers must also clearly and legibly identify you as the publisher of these copies. The front cover must present the full title with all words of the title equally prominent and visible. You may add other material on the covers in addition. Copying with changes limited to the covers, as long as they preserve the title of the Document and satisfy these conditions, can be treated as verbatim copying in other respects.

If the required texts for either cover are too voluminous to fit legibly, you should put the first ones listed (as many as fit reasonably) on the actual cover, and continue the rest onto adjacent pages.

If you publish or distribute Opaque copies of the Document numbering more than 100, you must either include a machine-readable Transparent copy along with each Opaque copy, or state in or with each Opaque copy a computer-network location from which the general network-using public has access to download using public-standard network protocols a complete Transparent copy of the Document, free of added material. If you use the latter option, you must take reasonably prudent steps, when you begin distribution of Opaque copies in quantity, to ensure that this Transparent copy will remain thus accessible at the stated location until at least one year after the last time you distribute an Opaque copy (directly or through your agents or retailers) of that edition to the public.

It is requested, but not required, that you contact the authors of the Document well before redistributing any large number of copies, to give them a chance to provide you with an updated version of the Document.

4. MODIFICATIONS

You may copy and distribute a Modified Version of the Document under the conditions of sections 2 and 3 above, provided that you release the Modified Version under precisely this License, with the Modified Version filling the role of the Document, thus licensing distribution and modification of the Modified Version to whoever possesses a copy of it. In addition, you must do these things in the Modified Version:

- A. Use in the Title Page (and on the covers, if any) a title distinct from that of the Document, and from those of previous versions (which should, if there were any, be listed in the History section of the Document). You may use the same title as a previous version if the original publisher of that version gives permission.
- B. List on the Title Page, as authors, one or more persons or entities responsible for authorship of the modifications in the Modified Version, together with at least five of the principal authors of the Document (all of its principal authors, if it has fewer than five), unless they release you from this requirement.
- C. State on the Title page the name of the publisher of the Modified Version, as the publisher.
- D. Preserve all the copyright notices of the Document.
- E. Add an appropriate copyright notice for your modifications adjacent to the other copyright notices.
- F. Include, immediately after the copyright notices, a license notice giving the public permission to use the Modified Version under the terms of this License, in the form shown in the Addendum below.
- G. Preserve in that license notice the full lists of Invariant Sections and required Cover Texts given in the Document's license notice.
- H. Include an unaltered copy of this License.

- I. Preserve the section Entitled “History”, Preserve its Title, and add to it an item stating at least the title, year, new authors, and publisher of the Modified Version as given on the Title Page. If there is no section Entitled “History” in the Document, create one stating the title, year, authors, and publisher of the Document as given on its Title Page, then add an item describing the Modified Version as stated in the previous sentence.
- J. Preserve the network location, if any, given in the Document for public access to a Transparent copy of the Document, and likewise the network locations given in the Document for previous versions it was based on. These may be placed in the “History” section. You may omit a network location for a work that was published at least four years before the Document itself, or if the original publisher of the version it refers to gives permission.
- K. For any section Entitled “Acknowledgements” or “Dedications”, Preserve the Title of the section, and preserve in the section all the substance and tone of each of the contributor acknowledgements and/or dedications given therein.
- L. Preserve all the Invariant Sections of the Document, unaltered in their text and in their titles. Section numbers or the equivalent are not considered part of the section titles.
- M. Delete any section Entitled “Endorsements”. Such a section may not be included in the Modified Version.
- N. Do not retitle any existing section to be Entitled “Endorsements” or to conflict in title with any Invariant Section.
- O. Preserve any Warranty Disclaimers.

If the Modified Version includes new front-matter sections or appendices that qualify as Secondary Sections and contain no material copied from the Document, you may at your option designate some or all of these sections as invariant. To do this, add their titles to the list of Invariant Sections in the Modified Version’s license notice. These titles must be distinct from any other section titles.

You may add a section Entitled “Endorsements”, provided it contains nothing but endorsements of your Modified Version by various parties—for example, statements of peer review or that the text has been approved by an organization as the authoritative definition of a standard.

You may add a passage of up to five words as a Front-Cover Text, and a passage of up to 25 words as a Back-Cover Text, to the end of the list of Cover Texts in the Modified Version. Only one passage of Front-Cover Text and one of Back-Cover Text may be added by (or through arrangements made by) any one entity. If the Document already includes a cover text for the same cover, previously added by you or by arrangement made by the same entity you are acting on behalf of, you may not add another; but you may replace the old one, on explicit permission from the previous publisher that added the old one.

The author(s) and publisher(s) of the Document do not by this License give permission to use their names for publicity for or to assert or imply endorsement of any Modified Version.

5. COMBINING DOCUMENTS

You may combine the Document with other documents released under this License, under the terms defined in section 4 above for modified versions, provided that you include in the combination all of the Invariant Sections of all of the original documents, unmodified, and list them all as Invariant Sections of your combined work in its license notice, and that you preserve all their Warranty Disclaimers.

The combined work need only contain one copy of this License, and multiple identical Invariant Sections may be replaced with a single copy. If there are multiple Invariant Sections with the same name but different contents, make the title of each such section unique by adding at the end of it, in parentheses, the name of the original author or publisher of that section if known, or else a unique number. Make the same adjustment to the section titles in the list of Invariant Sections in the license notice of the combined work.

In the combination, you must combine any sections Entitled “History” in the various original documents, forming one section Entitled “History”; likewise combine any sections Entitled “Acknowledgements”, and any sections Entitled “Dedications”. You must delete all sections Entitled “Endorsements”.

6. COLLECTIONS OF DOCUMENTS

You may make a collection consisting of the Document and other documents released under this License, and replace the individual copies of this License in the various documents with a single copy that is included in the collection, provided that you follow the rules of this License for verbatim copying of each of the documents in all other respects.

You may extract a single document from such a collection, and distribute it individually under this License, provided you insert a copy of this License into the extracted document, and follow this License in all other respects regarding verbatim copying of that document.

7. AGGREGATION WITH INDEPENDENT WORKS

A compilation of the Document or its derivatives with other separate and independent documents or works, in or on a volume of a storage or distribution medium, is called an “aggregate” if the copyright resulting from the compilation is not used to limit the legal rights of the compilation’s users beyond what the individual works permit. When the Document is included in an aggregate, this License does not apply to the other works in the aggregate which are not themselves derivative works of the Document.

If the Cover Text requirement of section 3 is applicable to these copies of the Document, then if the Document is less than one half of the entire aggregate, the Document’s Cover Texts may be placed on covers that bracket the Document within the aggregate, or the electronic equivalent of covers if the Document is in electronic form. Otherwise they must appear on printed covers that bracket the whole aggregate.

8. TRANSLATION

Translation is considered a kind of modification, so you may distribute translations of the Document under the terms of section 4. Replacing Invariant Sections with translations requires special permission from their copyright holders, but you may include translations of some or all Invariant Sections in addition to the original versions of these Invariant Sections. You may include a translation of this License, and all the license notices in the Document, and any Warranty Disclaimers, provided that you also include the original English version of this License and the original versions of those notices and disclaimers. In case of a disagreement between the translation and the original version of this License or a notice or disclaimer, the original version will prevail.

If a section in the Document is Entitled “Acknowledgements”, “Dedications”, or “History”, the requirement (section 4) to Preserve its Title (section 1) will typically require changing the actual title.

9. TERMINATION

You may not copy, modify, sublicense, or distribute the Document except as expressly provided under this License. Any attempt otherwise to copy, modify, sublicense, or distribute it is void, and will automatically terminate your rights under this License.

However, if you cease all violation of this License, then your license from a particular copyright holder is reinstated (a) provisionally, unless and until the copyright holder explicitly and finally terminates your license, and (b) permanently, if the copyright holder fails to notify you of the violation by some reasonable means prior to 60 days after the cessation.

Moreover, your license from a particular copyright holder is reinstated permanently if the copyright holder notifies you of the violation by some reasonable means, this is the first time you have received notice of violation of this License (for any work) from that copyright holder, and you cure the violation prior to 30 days after your receipt of the notice.

Termination of your rights under this section does not terminate the licenses of parties who have received copies or rights from you under this License. If your rights have been terminated and not permanently reinstated, receipt of a copy of some or all of the same material does not give you any rights to use it.

10. FUTURE REVISIONS OF THIS LICENSE

The Free Software Foundation may publish new, revised versions of the GNU Free Documentation License from time to time. Such new versions will be similar in spirit to the present version, but may differ in detail to address new problems or concerns. See <http://www.gnu.org/copyleft/>.

Each version of the License is given a distinguishing version number. If the Document specifies that a particular numbered version of this License “or any later version” applies to it, you have the option of following the terms and conditions either of that specified version or of any later version that has been published (not as a draft) by the Free Software Foundation. If the Document does not specify a version number of this License, you may choose any version ever published (not as a draft) by the Free Software Foundation. If the Document specifies that a proxy can decide which future versions of this License can be used, that proxy’s public statement of acceptance of a version permanently authorizes you to choose that version for the Document.

11. RELICENSING

“Massive Multiauthor Collaboration Site” (or “MMC Site”) means any World Wide Web server that publishes copyrightable works and also provides prominent facilities for anybody to edit those works. A public wiki that anybody can edit is an example of such a server. A “Massive Multiauthor Collaboration” (or “MMC”) contained in the site means any set of copyrightable works thus published on the MMC site.

“CC-BY-SA” means the Creative Commons Attribution-Share Alike 3.0 license published by Creative Commons Corporation, a not-for-profit corporation with a principal place of business in San Francisco, California, as well as future copyleft versions of that license published by that same organization.

“Incorporate” means to publish or republish a Document, in whole or in part, as part of another Document.

An MMC is “eligible for relicensing” if it is licensed under this License, and if all works that were first published under this License somewhere other than this MMC, and subsequently incorporated in whole or in part into the MMC, (1) had no cover texts or invariant sections, and (2) were thus incorporated prior to November 1, 2008.

The operator of an MMC Site may republish an MMC contained in the site under CC-BY-SA on the same site at any time before August 1, 2009, provided the MMC is eligible for relicensing.

ADDENDUM: How to use this License for your documents

To use this License in a document you have written, include a copy of the License in the document and put the following copyright and license notices just after the title page:

Copyright © YEAR YOUR NAME. Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.3 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled “GNU Free Documentation License”.

If you have Invariant Sections, Front-Cover Texts and Back-Cover Texts, replace the “with ... Texts.” line with this:

with the Invariant Sections being LIST THEIR TITLES, with the Front-Cover Texts being LIST, and with the Back-Cover Texts being LIST.

If you have Invariant Sections without Cover Texts, or some other combination of the three, merge those two alternatives to suit the situation.

If your document contains nontrivial examples of program code, we recommend releasing these examples in parallel under your choice of free software license, such as the GNU General Public License, to permit their use in free software.

